

Moisture Effects on Entomopathogenic Nematodes

Jennifer A. Grant and Michael G. Villani

Department of Entomology, NYSAES, Geneva

BACKGROUND

Entomopathogenic nematodes are well adapted to infect larval insect pests living in soil, and have the potential to be important biological control agents in a variety of ornamental and crop production systems. In the mid 1980's, several species of entomopathogenic nematodes became commercially available for insect pest management. Initially, small-scale production and limited marketing resulted in these products being used mainly for home gardens, lawns and landscapes. More recently, a few large companies have attempted mainstream marketing aimed at the commercial turf, vegetable, and fruit industries, but acceptance has been hindered by variability in the success of the nematodes' ability to control target insects.

More consistent results have been achieved by educating users on better application techniques and appropriate selection of nematode species and strains for particular uses. However, the impact of the agronomic environment on nematode ecology must be better understood before nematodes can be a reliable pest management alternative. Because entomopathogenic nematodes need both high relative humidity to survive and a film of free water for movement, moisture conditions have been recognized as one of the most important factors in the soil environment affecting survival, infectivity and persistence of nematodes (Curran 1993, Klein 1990).

Our 1997 work, reported last year, examined the effects of soil moisture on entomopathogenic nematode infectivity, under both laboratory and field conditions. In 1998, we concentrated our research on the ability of nematodes to be reactivated to infect insects after long periods of storage under unfavorable (dry soil) conditions. Our work investigates nematode persistence and reactivation in naturally varying climatic and soil conditions. The project has broad implications for managing soil dwelling insects with entomopathogenic nematodes in turf, ornamental and crop production systems.

For a printed copy of the entire report, please contact the NYS IPM office at:

IPM House
630 W. North St.
New York State Agricultural Experiment Station
Geneva NY 14456
315-878-2353