

THE VEC-PERMUTATION MATRIX, THE VEC OPERATOR AND
KRONECKER PRODUCTS: A REVIEW

by

Harold V. Henderson*

Biometrics Section, Ruakura Agricultural Research Centre,
Hamilton, New Zealand

and

S. R. Searle

Biometrics Unit, Cornell University, Ithaca, New York

ABSTRACT

The vec-permutation matrix $I_{\tilde{m},n}$ is defined by the equation $\text{vec}A_{\tilde{m} \times n} = I_{\tilde{m},n} \text{vec}A'$, where vec is the vec operator such that $\text{vec}A$ is the vector of columns of A stacked one under the other. The variety of definitions, names and notations for $I_{\tilde{m},n}$ are discussed, and its properties are developed by simple proofs in contrast to certain lengthy proofs in the literature that are based on descriptive definitions. For example, the role of $I_{\tilde{m},n}$ in reversing the order of Kronecker products is succinctly derived using the vec operator. The matrix $M_{\tilde{m},n}$ is introduced as $M_{\tilde{m},n} = I_{\tilde{m},n} M$; it is the matrix having for rows, every n 'th row of M , of order $mn \times c$, starting with the first, then every n 'th row starting with the second, and so on. Special cases of $M_{\tilde{m},n}$ are discussed.

Paper No. BU-645-M in the Biometrics Unit, Cornell University.

* Supported by a New Zealand National Research Advisory Council Research Fellowship during his sojourn at Cornell.

1. NOTATION

Let $\underline{\underline{A}}$ and $\underline{\underline{B}}$ be matrices of order $m \times n$ and $p \times q$, respectively. Rows of a matrix will be denoted by Greek letters transposed and columns by Roman letters:

$$\underline{\underline{A}} = \{a_{ij}\} = [a_{\sim 1} \cdots a_{\sim n}] = \begin{bmatrix} \alpha'_{\sim 1} \\ \vdots \\ \alpha'_{\sim m} \end{bmatrix} \text{ and } \underline{\underline{B}} = \{b_{rs}\} = [b_{\sim 1} \cdots b_{\sim q}] = \begin{bmatrix} \beta'_{\sim 1} \\ \vdots \\ \beta'_{\sim p} \end{bmatrix}. \quad (1)$$

$\underline{\underline{I}}_n$ is the identity matrix of order n , with $e_{\sim i}$ denoting its i 'th column and then

$$\underline{\underline{I}}_n = \sum_{i=1}^n e_{\sim i} e'_{\sim i}.$$

2. THE VEC OPERATOR AND KRONECKER PRODUCTS

2.1. The vec operator and its many names and notations

The vec operator stacks the columns of a matrix one underneath the other to form a single vector. Thus for $\underline{\underline{A}}$ of (1)

$$\text{vec} \underline{\underline{A}} = \begin{bmatrix} a_{\sim 1} \\ \vdots \\ a_{\sim n} \end{bmatrix}, \quad (2)$$

with $\text{vec} \underline{\underline{A}}$ (for "vector of columns of $\underline{\underline{A}}$ ") being the notation currently in vogue. The equivalent notations $\text{vec} \underline{\underline{A}}$ and $\text{vec}(\underline{\underline{A}})$ are used interchangeably, the parentheses being employed only when deemed necessary for clarity.

An early reference to this idea of stacking elements of a matrix in a vector is Sylvester [1884a,b,c] who used it in connection with linear equations. Roth [1934], using the notation \underline{A}^c , develops results for using the operation on a product matrix, Aitken [1949] mentions the idea in connection with Jacobians, and Koopmans et al. [1950] introduce the notation vec. More recently, the concept has been exploited in a variety of ways, for example, in solving linear matrix equations and in matrix differentiation, from which it is seen to be useful in deriving Jacobians of matrix transformations. The paper by Henderson and Searle [1979] outlines these applications and highlights several uses in statistics; e.g., in rewriting multivariate linear models in a univariate form, in developing the dispersion matrix of elements of a matrix such as a Wishart matrix and, from this, in deriving fourth moments in a general linear model.

As a result of these applications, the concept has, in recent years, been used by numerous writers. There is vecA or vec(A) used by Neudecker [1968, 1969a,b], Browne [1974] (who also uses a), Swain [1975] (who also uses A), Conlisk [1976], Balestra [1976], Anderson et al. [1977], Anderson [1978], Searle [1978, 1979], Brewer [1978], Magnus and Neudecker [1979] and Henderson and Searle [1979]. Equivalent notations are S(A), for stacking columns of A, used by Nissen [1968] who also uses a; and L(A) used by Conlisk [1969]; also, A for the column-rolled-out form of A, as it is described by Cole [1969]; and csA as the column string of A used by Vetter [1970, 1973, 1975], Kucera [1974] and Mitra [1977]. There is also

\underline{a}_c of Tracy and Dwyer [1969], Singh [1972] and Tracy and Singh [1972a,b], $\text{col}(\underline{A})$ of Hartwig [1972, 1975] and Hartwig and Morris [1975], $\theta(\underline{A})$ for the pack of \underline{A} , in MacRae [1974] and in Rogers and Young [1978]; and $\text{vc}\underline{A}$ as the vector of columns of \underline{A} by Legault-Giguère [1974] and Giguère and Styan [1974] and \underline{a}_c by Nel [1978].

Variations on what we have defined as $\text{vec}\underline{A}$ are also available: for example, having it be the column vector derived from writing each transposed row of \underline{A} one under the other. This stacks the elements of \underline{A} in lexicon order and is, in our notation, equivalent to $\text{vec}\underline{A}'$, where \underline{A}' is the transpose of \underline{A} . Notations for this include \underline{a} used by Lancaster [1969, 1970], \underline{A}_r by Tracy and Dwyer [1969], Singh [1972] and Tracy and Singh [1972a,b], $v(\underline{A})$ by Barnett [1973], $\text{vr}\underline{A}$ as the vector of transposed rows of \underline{A} by Legault-Giguère [1974] and \underline{a}_r by Nel [1978]; and even $\text{vec}\underline{A}$ by McDonald and Swaminathan [1973], Bentler and Lee [1975], McDonald [1976] and Pukelsheim [1977]. Other variations include the row vector of rows of \underline{A} , $(\text{vec}\underline{A}')'$ in our notation, which Roth [1934] denotes by \underline{A}^R and Vetter [1970, 1973, 1975] by $\text{rs}\underline{A}$, as the row string of \underline{A} . A final form, the row vector of transposed columns of \underline{A} , $(\text{vec}\underline{A})'$ has not yet been seen in the literature!

2.2. Origins of the Kronecker product

The Kronecker product of two matrices is defined for \underline{A} and \underline{B} of (1) as the $mp \times nq$ matrix

$$\underline{A} \otimes \underline{B} = \{a_{ij}\underline{B}\} , \quad (3)$$

which is a particular case of the tensor product for transformations as discussed, for example, by Halmos [1958, p. 97]. In its original setting it seems to have been first studied by Zehfuss [1858], and although he dealt only with its determinant, Rutherford [1933] appropriately calls $\underline{\underline{A}} \otimes \underline{\underline{B}}$ the Zehfuss matrix of $\underline{\underline{A}}$ and $\underline{\underline{B}}$. Loewy [1910, pp. 149-150] refers to Zehfuss's determinantal result as Kronecker's theorem which, according to Hensel [1891, p. 319], Kronecker (1823-91) had for sometime given in his algebra lectures in Berlin, which Bell [1937, p. 478] notes, he presented regularly from 1861 "principally on his personal researches, after the necessary introductions". Thus, although the name of Kronecker is now generally associated with the $\underline{\underline{A}} \otimes \underline{\underline{B}}$ operation, with an early use as pointed out by Dr. George Styan being Murnaghan [1938, p. 68], the exact origin of this association is hard to find.

Basic properties of Kronecker products are collected, with references to sources after 1889, by Loewy [1910, pp. 148-151] and by MacDuffee [1933, p. 82] using the names "product transformation" and "direct product", respectively. More recent summaries are Searle [1966], Neudecker [1968, 1969a,b] and Graybill [1969] in the context of statistical applications, and Barnett [1979] with engineering applications.

2.3. The vec and Kronecker product operators are connected

The definitions of vec and of Kronecker product show that $\text{vec}(\underline{ab}') = \underline{b} \otimes \underline{a}$, so that $\text{vec}[(\underline{Aa})(\underline{b}'\underline{C})] = (\underline{C}' \otimes \underline{A})\text{vec}(\underline{ab}')$. This, together with $\underline{B} = \{b_{ij}\} = \sum_{ij} b_{ij} \underline{e}_i \underline{e}_j'$, leads to

$$\text{vec}(\underline{ABC}) = (\underline{C}' \otimes \underline{A})\text{vec}\underline{B}, \quad (4)$$

a result derived by Roth [1934], and hence called Roth's column lemma by Hartwig [1975], and rediscovered by Aitken [1949], Koopmans et al. [1950], Nissen [1968] and Neudecker [1969b].

3. VEC-PERMUTATION MATRICES

The vectors $\text{vec}\underline{A}$ and $\text{vec}\underline{A}'$ contain the same elements, in different sequences. We define $\underline{I}_{m,n}$ as the permutation matrix such that, for \underline{A} of order m by n ,

$$\text{vec}\underline{A} = \underline{I}_{m,n} \text{vec}\underline{A}', \quad (5)$$

and give it the name vec-permutation matrix.

There has been increasing interest over the last decade in vec-permutation matrices in matrix algebra, mathematics, statistics, econometrics and psychometrics. This wide interest in what we call

$\tilde{I}_{m,n}$ of (5) is partly responsible for the diversity of its literature, for duplication of published results, for varying definitions and for a wide range of notation and nomenclature. For example, (5) defines what $\tilde{I}_{m,n}$ does, in contrast to definitions discussed later that describe what $\tilde{I}_{m,n}$ looks like. This is an important distinction because operational definitions of $\tilde{I}_{m,n}$, like (5), lead to succinct derivations of its properties, in contrast to some recent and rather tedious proofs based on descriptive definitions. An example of this is the important role of $\tilde{I}_{m,n}$ in reversing the order of Kronecker products; e.g., for $\tilde{A}_{m \times n}$ and $\tilde{B}_{p \times q}$

$$\tilde{B} \otimes \tilde{A} = \tilde{I}_{m,p} (\tilde{A} \otimes \tilde{B}) \tilde{I}_{q,n} . \quad (6)$$

In these circumstances, and because of the topical nature of the subject, a cohesive account and succinct development of properties of vec-permutation matrices is needed. This is attempted in Section 4. But first we discuss the variety of definitions, names and notations to be found in the literature and summarized in Table 1, for a vec-permutation matrix, using our notation $\tilde{I}_{m,n}$.

(Show Table 1 here)

3.1. Definitions from reversing the order of Kronecker products of matrices and vectors

Ledermann [1936], in material from his Ph.D. thesis supervised by A. C. Aitken, gives an elegant proof of (6), although without explicit development of the vec-permutation matrices. Using column vectors \tilde{r} , \tilde{x} , \tilde{s} and \tilde{y} of order m , n , p and q , respectively, with $\tilde{r} = \tilde{A}\tilde{x}$ and $\tilde{s} = \tilde{B}\tilde{y}$, he proceeds as follows.

"The two products $\underline{\underline{A}} \otimes \underline{\underline{B}}$ and $\underline{\underline{B}} \otimes \underline{\underline{A}}$ are related to each other by an identity

$$\underline{\underline{Q}}(\underline{\underline{A}} \otimes \underline{\underline{B}})\underline{\underline{P}}^{-1} = (\underline{\underline{B}} \otimes \underline{\underline{A}}) ,$$

where $\underline{\underline{P}}$ and $\underline{\underline{Q}}$ are permutation matrices which depend only on the types of $\underline{\underline{A}}$ and $\underline{\underline{B}}$ and not on their elements.

Proof: Apart from the order, the vectors $\underline{\underline{x}} \otimes \underline{\underline{y}}$ and $\underline{\underline{y}} \otimes \underline{\underline{x}}$ contain the same elements ... We can therefore find a permutation matrix $\underline{\underline{P}}$ of degree nq such that

$$(\underline{\underline{y}} \otimes \underline{\underline{x}}) = \underline{\underline{P}}(\underline{\underline{x}} \otimes \underline{\underline{y}})$$

and similarly

$$(\underline{\underline{s}} \otimes \underline{\underline{r}}) = \underline{\underline{Q}}(\underline{\underline{r}} \otimes \underline{\underline{s}}) ,$$

where $\underline{\underline{Q}}$ is a permutation matrix of degree mp . Evidently $\underline{\underline{P}}$ and $\underline{\underline{Q}}$ do not depend on the elements of $\underline{\underline{x}}$, $\underline{\underline{y}}$, $\underline{\underline{r}}$, $\underline{\underline{s}}$ but only on the numbers m , n , p , q . By $[(\underline{\underline{A}} \otimes \underline{\underline{B}})(\underline{\underline{x}} \otimes \underline{\underline{y}}) = \underline{\underline{A}}\underline{\underline{x}} \otimes \underline{\underline{B}}\underline{\underline{y}}]$ we have

$$(\underline{\underline{r}} \otimes \underline{\underline{s}}) = (\underline{\underline{A}} \otimes \underline{\underline{B}})(\underline{\underline{x}} \otimes \underline{\underline{y}}),$$

$$(\underline{\underline{s}} \otimes \underline{\underline{r}}) = (\underline{\underline{B}} \otimes \underline{\underline{A}})(\underline{\underline{y}} \otimes \underline{\underline{x}}).$$

On premultiplying the first equation by $\underline{\underline{Q}}$ and substituting ... we get

$$(\underline{\underline{s}} \otimes \underline{\underline{r}}) = \underline{\underline{Q}}(\underline{\underline{A}} \otimes \underline{\underline{B}})(\underline{\underline{x}} \otimes \underline{\underline{y}}) = (\underline{\underline{B}} \otimes \underline{\underline{A}})\underline{\underline{P}}(\underline{\underline{x}} \otimes \underline{\underline{y}}) .$$

Since there is obviously no linear relation between the elements of $\underline{\underline{x}} \otimes \underline{\underline{y}}$, we obtain

$$\underline{\underline{Q}}(\underline{\underline{A}} \otimes \underline{\underline{B}}) = (\underline{\underline{B}} \otimes \underline{\underline{A}})\underline{\underline{P}}$$

or

$$\underline{\underline{Q}}(\underline{\underline{A}} \otimes \underline{\underline{B}})\underline{\underline{P}}^{-1} = (\underline{\underline{B}} \otimes \underline{\underline{A}}) ,$$

as was to be proved. "

[For consistent notation we have replaced x in the original form of this quotation by \otimes , and changed subscripts to conform with (1); and we do this in all quotations, without further comment.]

Ledermann [1936] then, in our notation, defines $\underline{I}_{\underline{m}, \underline{n}}$ as the permutation matrix such that

$$\underline{I}_{\underline{m}, \underline{n}}(\underline{a} \otimes \underline{b}) = \underline{b} \otimes \underline{a}, \quad (7)$$

for any column vectors \underline{a} and \underline{b} of order m and n , respectively.

Conlisk [1976] has a similar idea in mind for $\underline{I}_{\underline{n}, \underline{n}}$, using the notation \underline{M} . [Actually, he writes (p. 760) $(\underline{a} \otimes \underline{b})\underline{M} = \underline{b} \otimes \underline{a}$, which is clearly incorrect, from dimension considerations alone. However, he correctly claims (p. 763) that $\underline{M}(\underline{a} \otimes \underline{B}_{\underline{n} \times \underline{n}}) = \underline{B} \otimes \underline{a}$.] Defining $\underline{I}_{\underline{m}, \underline{n}}$, as in (7), in terms of reversing the order of Kronecker products of vectors, was also suggested by a referee of Henderson and Searle [1979]. The suggestion has merit, but it emphasizes a feature of $\underline{I}_{\underline{m}, \underline{n}}$ that is second to the more fundamental and more general definition given in (5) in terms of the vec operator. In fact, (7) is the special case of (5) with \underline{A} being $\underline{b}\underline{a}'$, of rank 1.

Murnaghan [1938] was also interested in reversing the order of Kronecker products and, in this connection, gives an explicit formulation of what we call a vec -permutation matrix. He denotes the position of $a_{ij}b_{rs}$ in $\underline{A} \otimes \underline{B}$ by the row index-pair (i, r) and column index-pair (j, s) , and notes (p. 68-69) that for \underline{A} and \underline{B} square of order m and n , respectively,

" $\underline{\underline{B}} \otimes \underline{\underline{A}}$ is obtainable from $\underline{\underline{A}} \otimes \underline{\underline{B}}$ by applying the same permutation to the rows and columns of the latter. Let us denote by $(i,r)^*$ the position of (r,i) when the ordering is dictionary-like, the label with the range n coming first; e.g., if $m = 2, n = 3$ the dictionary order where the label with the range 2 comes first is $(1,1), (1,2), (1,3), (2,1), (2,2), (2,3)$ whilst the dictionary order when the label with the range 3 comes first is $(1,1), (1,2), (2,1), (2,2), (3,1), (3,2)$. Hence $(1,1)^* = (1,1), (1,2)^* = (1,3), (1,3)^* = (2,2), (2,1)^* = (1,2), (2,2)^* = (2,1), (2,3)^* = (2,3)$. Then if the (j,s) column of $\underline{\underline{A}} \otimes \underline{\underline{B}}$ is transferred to the $(j,s)^*$ position and the (i,r) row to the $(i,r)^*$ position we obtain $\underline{\underline{B}} \otimes \underline{\underline{A}}$. In other words

$$(\underline{\underline{B}} \otimes \underline{\underline{A}}) = P(\underline{\underline{A}} \otimes \underline{\underline{B}})P^{-1} \quad (8)$$

where P is the permutation matrix associated with $\begin{pmatrix} (i,r)^* \\ (i,r) \end{pmatrix}$.

Notice that P is quite independent of the elements of $\underline{\underline{A}}$ and $\underline{\underline{B}}$ being completely determined by their dimensions."

A clearer statement of Murnaghan's [1938] development of P , which is our $\underline{\underline{I}}_{m,n}$, is that

$$\text{row } (i, j) \text{ of } \underline{\underline{I}}_{m, n} \text{ is row } (j, i) \text{ of } \underline{\underline{I}}_{mn}, \quad (9)$$

for $i = 1, \dots, n$ and $j = 1, \dots, m$. Note that $\underline{\underline{I}}_{mn}$ is the identity matrix of order mn , to be distinguished from $\underline{\underline{I}}_{m, n}$, a vec-permutation matrix of the same order. Searle [1966, p. 216] makes a more explicit statement of this situation:

"It is apparent from (3) that for $\underline{\underline{A}}_{m \times n} = \{a_{ij}\}$ and $\underline{\underline{B}}_{p \times q} = \{b_{rs}\}$ the elements of both $\underline{\underline{A}} \otimes \underline{\underline{B}}$ and $\underline{\underline{B}} \otimes \underline{\underline{A}}$ consist of all possible products $a_{ij}b_{rs}$. In fact, $\underline{\underline{B}} \otimes \underline{\underline{A}}$ is simply $\underline{\underline{A}} \otimes \underline{\underline{B}}$ with the rows and columns each in a different order. Thus [in general]

$$\underline{\underline{B}} \otimes \underline{\underline{A}} = \underline{\underline{P}}(\underline{\underline{A}} \otimes \underline{\underline{B}})\underline{\underline{Q}} \quad (10)$$

where $\underline{\underline{P}}$ and $\underline{\underline{Q}}$ are each a product of E-type elementary operators [permutation matrices].

For any values of i, j, r and s the element $a_{ij}b_{rs}$ is located in $\underline{\underline{A}} \otimes \underline{\underline{B}}$ in the r 'th row and s 'th column of the i, j 'th sub-matrix $[a_{ij} \underline{\underline{B}}] \dots$. It is therefore in row $[p(i - 1) + r]$ and column $[q(j - 1) + s]$ of $\underline{\underline{A}} \otimes \underline{\underline{B}}$. In $\underline{\underline{B}} \otimes \underline{\underline{A}}$, however, $b_{rs}a_{ij}$ is in row i and column j of the r, s 'th sub-matrix $[b_{rs} \underline{\underline{A}}] \dots$. It is therefore in row $[m(r - 1) + i]$ and column $[n(s - 1) + j]$ of $\underline{\underline{B}} \otimes \underline{\underline{A}}$. Consequently the interchanging of rows and columns implied in (10), to obtain $\underline{\underline{B}} \otimes \underline{\underline{A}}$ from $\underline{\underline{A}} \otimes \underline{\underline{B}}$, can be specified as follows. The $[i + (j - 1)m]$ 'th row of $\underline{\underline{P}}$ is the $[(i - 1)p + j]$ 'th row of $\underline{\underline{I}}_{mp}$, for $i = 1, 2, \dots, m$ and $j = 1, 2, \dots, p$, and the $[i + (j - 1)n]$ 'th column of $\underline{\underline{Q}}$

is the $[(i - 1)q + j]$ 'th column of \underline{I}_{nq} , for $i = 1, 2, \dots, n$ and $j = 1, 2, \dots, q$."

Comparison of (6) with (10) shows that $\underline{P} = \underline{I}_{m,p}$ and $\underline{Q} = \underline{I}_{q,n}$; so that Searle's [1966] expression for \underline{P} may be restated, in our notation, as

$$\text{row } (i - 1)m + j \text{ of } \underline{I}_{m,n} \text{ is row } (j - 1)n + i \text{ of } \underline{I}_{mn}, \quad (11)$$

for $i = 1, \dots, n$ and $j = 1, \dots, m$. This is, of course, equivalent to (9) because row (i,j) of $\underline{I}_{m,n}$ is its $[(i - 1)m + j]$ 'th row. Expressions (9) and (11) show consecutive rows of $\underline{I}_{m,n}$ to be every n 'th row of \underline{I}_{mn} starting with the first, and so on; a fact which Tracy and Dwyer [1969] rediscovered and used as its definition, as is now discussed.

3.2. Defining the row permutations

Tracy and Dwyer [1969, p. 1579] introduce

$\tilde{I}_{(n)}$ as the matrix obtained by rearranging
the rows of \tilde{I}_r by taking
every n 'th row starting with the first, then
every n 'th row starting with the second, and so on. (12)

Tracy and Singh [1972a,b] and Singh [1972], restricting $\tilde{I}_{(n)}$ to be of order mn , establish (5) and (6) and call $\tilde{I}_{(n)}$ a permuted identity matrix, a name which actually applies to any permutation matrix. We call $\tilde{I}_{(n)}$ of (12) a generalized vec-permutation matrix because it is defined for any order, not necessarily a multiple of n , whereas

$$\tilde{I}_{m,n} = \tilde{I}_{(n)} \text{ of order } mn . \tag{13}$$

For example, the rows of

$$\tilde{I}_{2,3} = \begin{bmatrix} 1 & . & . & | & . & . & . \\ . & . & . & | & 1 & . & . \\ \hline . & 1 & . & | & . & . & . \\ . & . & . & | & . & 1 & . \\ \hline . & . & 1 & | & . & . & . \\ . & . & . & | & . & . & 1 \end{bmatrix} \tag{14}$$

(where dots represent zeros), are every third row, namely rows 1, 4, 2, 5, 3 and 6, respectively, of \tilde{I}_6 .

3.3. Descriptive definitions of $\underline{I}_{m,n}$

In contrast to operational definitions like (5) and (7) through (13), there is a number of what can be called descriptive definitions based on describing what $\underline{I}_{m,n}$ looks like. They amount to defining

$\underline{I}_{m,n}$ as a square matrix of order mn , partitioned into an n by m array of submatrices each of order m by n , such that the (i,j) 'th submatrix \underline{S}_{ij} , say, has unity in its (j,i) 'th position and zeros elsewhere. (15)

For example, $\underline{I}_{2,3}$ of (14) is shown partitioned in accord with (15).

Variations of (15) include Vetter's [1970] $\langle \underline{U}_{ij}^{ji} \rangle$, Hartwig's [1972, p. 540] \underline{P} (which should be \underline{P}^{-1}) for his use in $\underline{P}^{-1}(\underline{A}_{m \times m} \otimes \underline{B}_{n \times n})\underline{P} = \underline{B} \otimes \underline{A}$; Vetter's [1973, 1975] $\underline{E}_{n \times m}^{m \times n}$; the "permuted identity matrix" $\underline{I}_{(m,n)}$ of MacRae [1974], who notes (p. 338) that it "is identical to the matrix $\underline{I}_{(n)}$ [of order mn] defined by Tracy and Dwyer [1969]" and the universal flip matrix $\underline{P}_{m,n}$ of Hartwig and Morris [1975]. Balestra [1976, p. 21], using $\underline{P}_{m,n}$, clarifies the partitioning in MacRae's [1974] definition. He also gives a symbolic version of (15):

$$\underline{I}_{m,n} = \underline{P}_{m,n} = \{ \underline{S}_{ij} \} \quad i = 1, \dots, n; j = 1, \dots, m,$$

where

$$\underline{S}_{ij} = \underline{e}_{j\sim} \underline{e}'_{i\sim} = \underline{e}_{\sim j} \otimes \underline{e}'_{\sim i} = \underline{e}'_{\sim i} \otimes \underline{e}_{\sim j},$$

which is a generalization of $\underline{I}_{(n,n)}$ in Anderson et al. [1977], with $\underline{S}_{ij} = \underline{e}_{j\sim} \underline{e}'_{i\sim}$ presented in Anderson [1978, p. 53].

A further formulation of (15) and of (16), where $\underset{\sim}{H}'_{ij}$ is $\underset{\sim}{S}_{ij}$,
is

$$\underset{\sim}{I}_{m,n} = \underset{\sim}{K}_{nm} = \sum_{i=1}^n \sum_{j=1}^m (\underset{\sim}{H}_{ij} \otimes \underset{\sim}{H}'_{ij})$$

where $\underset{\sim}{H}_{ij}$ is the $n \times m$ matrix with a 1 (17)
in position (i,j) and zeros elsewhere.

Vetter [1973, 1975] presents (17) as does Brewer [1977, 1978],
who uses $\underset{\sim}{U}_{n \times m}$ for $\underset{\sim}{I}_{m,n}$. Magnus and Neudecker [1979] also use (17),
which they derive from (5). They call it a "commutation matrix"
because of its role in reversing ("commuting") the order of Kronecker
products, and denote it by $\underset{\sim}{K}_{nm}$.

Definition (15) and its variations impose an artificial par-
titioning on $\underset{\sim}{I}_{m,n}$, which disguises its inherent permutation features.
Only the partitioning into n blocks of m rows or m blocks of n
columns, but not both, is natural. This natural partitioning is
implicit in the operational definitions and is explicit in a further
formulation given by Balestra [1976]:

$$\underset{\sim}{I}_{m,n} = \{ \underset{\sim}{I}_m \otimes \underset{\sim}{e}'_i \}_{i=1 \dots n} = \{ \underset{\sim}{I}_n \otimes \underset{\sim}{e}_j \}_{j=1 \dots m}, \quad (18)$$

the second of which is also given by Swain [1975] who uses $\underset{\sim}{L}_{mn}$ for
 $\underset{\sim}{I}_{m,n}$. Denoting the direct sum $\underset{\sim}{A}_1 \oplus \dots \oplus \underset{\sim}{A}_n$ by $\bigoplus_{i=1}^n \underset{\sim}{A}_i$, we see that
(18) may be written as

$$\underset{\sim}{I}_{m,n} = \left\{ \bigoplus_{j=1}^m \underset{\sim}{e}'_i \right\}_{i=1 \dots n} = \left\{ \bigoplus_{i=1}^n \underset{\sim}{e}_j \right\}_{j=1 \dots m}, \quad (19)$$

where in the first partitioning in (19) the $\underset{\sim}{e}'_i$ have order n , and in
the second the $\underset{\sim}{e}_j$ have order m .

Results (18) and (19) are nicely illustrated by (14). Two further formulations of $\underset{\sim}{I}_{m,n}$ developed by Magnus and Neudecker [1979] are

$$\underset{\sim}{I}_{m,n} = \sum_{i=1}^m (e'_i \otimes \underset{\sim}{I}_n \otimes e_i) \quad \text{and} \quad \underset{\sim}{I}_{m,n} = \sum_{j=1}^n (e_j \otimes \underset{\sim}{I}_m \otimes e'_j). \quad (20)$$

Although Hartwig and Morris [1975] give a block formulation definition of $\underset{\sim}{I}_{m,n}$, they give an ingenious card shuffling interpretation which is in fact based on (5), our definition. They identify the permutation implicit in $\underset{\sim}{I}_{m,n}$ with the "generalized out faro-shuffle" for a one dimensional deck of mn cards. Their description, on p. 451, with further details in Morris and Hartwig [1976], is as follows.

"In this shuffle a deck of mn cards, labeled from top to bottom, is cut into m portions of n cards, and each portion is given in clockwise fashion to one of m players seated at a circular table, starting with the dealer. If, starting with the dealer, in clockwise fashion, each player plays his top card when it is his turn, until all cards have been played, we obtain the permutation Π , labeled from the bottom cards up.

When $m = 2$ this reduces to the classical out faro-shuffle (in which an even deck of cards is cut in halves and then ruffled such that the first and the last cards remain in fixed positions) which is the basis to several remarkable card tricks."

A further application of these permutations, kindly brought to our attention by Dr. Stephen Barnett, is that of Whelchel and Guinn [1970] who are concerned with shuffling data in computer storage. In this context, they refer to $\underset{\sim}{I}_{n,n}$ as the "shuffle matrix", and denote it by $\underset{\sim}{S}_{n^2}$.

3.4. Defining $I_{\underline{m}, \underline{n}}$ as a matrix derivative operator

McDonald and Swaminathan [1973, p. 39] denote $I_{\underline{m}, \underline{n}}$ by E' , and subsequently in McDonald [1976] by $E_{\underline{n}, \underline{m}}$ to eliminate ambiguity, defined as the derivative operator

$$I_{\underline{m}, \underline{n}} = E_{\underline{n}, \underline{m}} = \frac{\partial X'}{\partial X}, \text{ where } X_{\underline{m} \times \underline{n}} \text{ is m.i.v.} \quad (21)$$

[A matrix is said to be mathematically independent and variable (m.i.v.) if no elements are functionally dependent or constant.]

They recognize $E_{\underline{m}, \underline{n}}$ as a permutation matrix of $I_{\underline{m} \times \underline{m}}$, and give (1973, p. 39) its form:

"the general element of $E_{\underline{n}, \underline{m}}$, e_{gh} , is equal to unity if $j = k'$ and $k = j'$ with

$$g = n(j - 1) + k, \quad 0 < j \leq m, \quad 0 < k \leq n,$$

$$h = m(j' - 1) + k', \quad 0 < j' \leq n, \quad 0 < k' \leq m,$$

and is zero otherwise."

This is a less concise statement than each of (9), (11) and (13).

Rather than define $I_{\underline{m}, \underline{n}}$ as the outcome of differentiation, as in (21) and which demands the m.i.v. property, and which fails to highlight the permutation properties of $I_{\underline{m}, \underline{n}}$, we prefer to derive (21) from the more fundamental definition (5) using the standard result $\frac{\partial Ax}{\partial x} = A'$ for x being m.i.v.; and then for X being m.i.v.

$$\frac{\partial X'}{\partial X} \equiv \frac{\partial \text{vec} X'}{\partial \text{vec} X} = \frac{\partial I_{\underline{n}, \underline{m}} \text{vec} X}{\partial \text{vec} X} = (I_{\underline{n}, \underline{m}})' = I_{\underline{m}, \underline{n}}.$$

4. PROPERTIES OF VEC-PERMUTATION MATRICES

Except for (6) which we consider separately, well-known properties of $\underline{I}_{m,n}$ are given without proof by Tracy and Singh [1972a,b], Singh [1972], Vetter [1973, p. 354] who suggests verification "by construction", MacRae [1974, p. 339] who comments they "can be verified by direct examination", Balestra [1976] and Swain [1975], and very recently by Brewer [1978] and Magnus and Neudecker [1979] whose proofs seem somewhat lengthy. Using $\underline{I}_{m,n}$ as defined in (5), we develop considerably shorter and simpler proofs.

Applying (5) to vecA' in (5) itself gives $\text{vecA} = \underline{I}_{m,n} \underline{I}_{n,m} \text{vecA}$ so that

$$\underline{I}_{m,n} \underline{I}_{n,m} = \underline{I}_{mn} . \quad (22)$$

Then, because $\underline{I}_{m,n}$ is a permutation matrix and so is orthogonal, we have

$$(\underline{I}_{m,n})^{-1} = (\underline{I}_{m,n})' = \underline{I}_{n,m} . \quad (23)$$

Also because $\text{vec a} = \text{vec a}'$

$$\underline{I}_{m,1} = \underline{I}_{1,m} = \underline{I}_m . \quad (24)$$

[We use the notation $(\underline{I}_{m,n})'$ and $\underline{I}'_{m,n}$ interchangeably.]

4.1. Reversing the order of Kronecker products of matrices

Vec-permutation matrices are related very directly to Kronecker products through the identity (6):

$$\underline{B}_{p \times q} \otimes \underline{A}_{m \times n} = \underline{I}_{m,p} (\underline{A} \otimes \underline{B}) \underline{I}_{q,n} . \quad (25)$$

Derivation of (25) comes from using the two fundamental properties of the vec operator, (4) and (5), as follows:

$$\begin{aligned} (\underline{B} \otimes \underline{A}) \text{vec} \underline{X} &= \text{vec} \underline{A} \underline{X} \underline{B}' = \underline{I}_{\underline{m}, \underline{p}} \text{vec} \underline{B} \underline{X}' \underline{A}' \\ &= \underline{I}_{\underline{m}, \underline{p}} (\underline{A} \otimes \underline{B}) \text{vec} \underline{X}' \\ &= \underline{I}_{\underline{m}, \underline{p}} (\underline{A} \otimes \underline{B}) \underline{I}_{\underline{q}, \underline{n}} \text{vec} \underline{X}, \end{aligned}$$

for any \underline{X} of order n by q . Letting $\text{vec} \underline{X}$ be in turn the columns of an identity matrix of order nq yields (25).

This concise proof was motivated by Barnett's [1973] use of the vec operator to establish (25) but without explicit development of the permutation matrices. Similar developments are also given by Hartwig and Morris [1975] and more recently by Magnus and Neudecker [1979], even though the latter do not use vec to prove results like (22) and (23).

Result (25) shows the exact form of the relationship between $\underline{A} \otimes \underline{B}$ and $\underline{B} \otimes \underline{A}$, which simply entails a resequencing of rows and of columns of $\underline{A} \otimes \underline{B}$ to obtain $\underline{B} \otimes \underline{A}$. Postmultiplying (25) by $\underline{I}_{\underline{n}, \underline{q}}$ and using (22) gives

$$\underline{I}_{\underline{m}, \underline{p}} (\underline{A} \otimes \underline{B}) = (\underline{B} \otimes \underline{A}) \underline{I}_{\underline{n}, \underline{q}}. \quad (26)$$

An early reference to reversing the order of Kronecker products of rectangular matrices is Ledermann [1936] although without explicit development of permutation matrices. But Murnaghan [1938, pp. 68-69] deals with square matrices, noting for square \underline{A} and \underline{B} , the validity of $\underline{B} \otimes \underline{A} = \underline{P}(\underline{A} \otimes \underline{B})\underline{P}^{-1}$ as in (8). Vartak [1955] generalizes this to (9), namely $\underline{B} \otimes \underline{A} = \underline{P}(\underline{A} \otimes \underline{B})\underline{Q}$, for rectangular \underline{A} and \underline{B} , indicating that a generalization of Murnaghan's proof can be con-

structed. Searle [1966, p. 216] shows that $a_{ij} b_{rs}$ is in row $i + (r - 1)m$ of $\underline{\underline{B}} \otimes \underline{\underline{A}}$ and row $(i - 1)p + r$ of $\underline{\underline{A}} \otimes \underline{\underline{B}}$, with a similar analysis for columns, and is thus able to give the explicit form of $\underline{\underline{P}}$ and $\underline{\underline{Q}}$, as detailed after equation (10). Hartwig [1972, p. 540] gives Murnaghan's result but misprints the form for $\underline{\underline{P}}^{-1}$ as being that for $\underline{\underline{P}}$. Singh [1972, p. 22] identifies columns in (26) and thence establishes (25). Tracy and Singh [1972b], MacRae [1974] and Swain [1975] give (25) without proof; so does Vetter [1973, p. 354] who comments that it is "well-known ... though the explicit transposition relationship is not usually given". In similar vein is the remark that "the proof [not given] ... is detailed and rather tedious" by Bentler and Lee [1975, p. 148], to whom McDonald [1976, p. 90] attributes (25). Balestra [1976, p. 23] uses (18), and Brewer [1978] suggests substituting (17). These developments are tedious in comparison to our proof using the vec operator.

4.2. Trace and determinant of $\underline{\underline{I}}_{m,n}$

The case of $m = n$ is easy. Because $\underline{\underline{I}}_{n,n}$ is real, orthogonal and symmetric, it has eigenvalues ± 1 with multiplicities $\frac{1}{2}n(n \pm 1)$, respectively. The determinant and trace, being the product and sum of the eigenvalues, respectively, are

$$|\underline{\underline{I}}_{n,n}| = (-1)^{\frac{1}{2}n(n-1)} \quad \text{and} \quad \text{tr}(\underline{\underline{I}}_{n,n}) = n.$$

For the more general situation, when m and n are not necessarily equal, the recurrence relation $|\underline{\underline{I}}_{m,n}| = (-1)^{\frac{1}{2}m(m-1)(n-1)} |\underline{\underline{I}}_{m,n-1}|$

yields

$$|\underline{I}_{\underline{m}, \underline{n}}| = (-1)^{\frac{1}{2}m(m-1)n(n-1)} .$$

This derivation is given by Hartwig and Morris [1975, p. 450] and Magnus and Neudecker [1979, p. 383] with an inductive proof presented by Swain [1975, Appendix A].

An expression for the trace is more difficult. Magnus and Neudecker [1979] prove that

$$\text{tr}(\underline{I}_{\underline{m}, \underline{n}}) = 1 + \text{gcd}(m - 1, n - 1) ,$$

where $\text{gcd}(m, n)$ is the greatest common divisor of m and n . A less compact form of this result is available in Hartwig and Morris [1975] who additionally derive expressions for the characteristic and minimal polynomials.

4.3. A generalization: $\underline{M}_{\underline{n}}$ and $\underline{M}_{\underline{m}, \underline{n}}$

Tracy and Dwyer [1969] introduce

$\underline{M}_{\underline{n}}$ as the $r \times c$ matrix formed by rearranging the rows of \underline{M} , of order $r \times c$, by taking every n 'th row starting with the first, then every n 'th row starting with the second, and so on, (27)

so that $\underline{I}_{\underline{n}}$ of (12) performs the row permutations on \underline{M} to obtain $\underline{M}_{\underline{n}}$ as

$$\underline{M}_{\underline{n}} = \underline{I}_{\underline{n}} \underline{M} . \quad (28)$$

For example, with dots denoting zeros, for

$$\underline{\underline{M}} = \begin{bmatrix} a' \\ \underline{\underline{b}}' \\ \underline{\underline{c}}' \end{bmatrix}, \quad \underline{\underline{M}}(2) = \begin{bmatrix} a' \\ \underline{\underline{c}}' \\ \underline{\underline{b}}' \end{bmatrix} = \begin{bmatrix} 1 & . & . \\ . & . & 1 \\ . & 1 & . \end{bmatrix} \begin{bmatrix} a' \\ \underline{\underline{b}}' \\ \underline{\underline{c}}' \end{bmatrix} = \underline{\underline{I}}(2) \underline{\underline{M}}.$$

Papers subsequent to Tracy and Dwyer [1969] have focused on $\underline{\underline{I}}(n)$, rather than on the more general notion of $\underline{\underline{M}}(n)$ that has wider applicability. In this context, we now define $\underline{\underline{M}}_{\underline{\underline{m}}, \underline{\underline{n}}}$ for $\underline{\underline{M}}$ with mn rows as

$$\underline{\underline{M}}_{\underline{\underline{m}}, \underline{\underline{n}}} = \underline{\underline{I}}_{\underline{\underline{m}}, \underline{\underline{n}}} \underline{\underline{M}} \quad (29)$$

$$= \underline{\underline{M}}(\underline{\underline{n}}) \text{ when } \underline{\underline{M}} \text{ has } mn \text{ rows.} \quad (30)$$

Our symbol $\underline{\underline{I}}_{\underline{\underline{m}}, \underline{\underline{n}}}$ for the vec-permutation matrix conforms with $\underline{\underline{M}}_{\underline{\underline{m}}, \underline{\underline{n}}}$ of (29) because it is $\underline{\underline{M}}_{\underline{\underline{m}}, \underline{\underline{n}}}$ with $\underline{\underline{M}} \equiv \underline{\underline{I}}$; i.e.,

$$\underline{\underline{I}}_{\underline{\underline{m}}, \underline{\underline{n}}} = \underline{\underline{I}}_{\underline{\underline{m}}, \underline{\underline{n}}} \underline{\underline{I}}, \quad (31)$$

so motivating use of the letter $\underline{\underline{I}}$ for the matrix $\underline{\underline{I}}_{\underline{\underline{m}}, \underline{\underline{n}}}$ in preference to any other letter. In the special case of $m = n$ there might sometimes be convenience in alternative symbols; e.g., $\underline{\underline{P}}_n \equiv \underline{\underline{I}}_{n, n}$ or $\underline{\underline{K}}_n$ as introduced by Magnus and Neudecker [1979].

Within the framework of the definition (29) of $\underline{\underline{M}}_{\underline{\underline{m}}, \underline{\underline{n}}}$ it can be noted that the defining property, (5), and results (25) and (26) can be rewritten:

$$\text{vec} \underline{\underline{A}} = (\text{vec} \underline{\underline{A}}')_{\underline{\underline{m}}, \underline{\underline{n}}}, \quad (32)$$

$$\underline{\underline{B}} \otimes \underline{\underline{A}} = [(\underline{\underline{A}} \otimes \underline{\underline{B}})'_{\underline{\underline{m}}, \underline{\underline{p}}}]'_{\underline{\underline{n}}, \underline{\underline{q}}} = [(\underline{\underline{A}}' \otimes \underline{\underline{B}}')'_{\underline{\underline{n}}, \underline{\underline{q}}}]'_{\underline{\underline{m}}, \underline{\underline{p}}}, \quad (33)$$

and

$$(\underline{\underline{A}} \otimes \underline{\underline{B}})_{\underline{\underline{m}}, \underline{\underline{p}}} = (\underline{\underline{B}}' \otimes \underline{\underline{A}}')'_{\underline{\underline{q}}, \underline{\underline{n}}}. \quad (34)$$

4.4. Equalities from reversing the order of Kronecker products

Alternative forms of (26) are now developed. In (34), $(\underline{A} \otimes \underline{B})_{\underline{m}, \underline{p}}$ is by definition a reordering of the rows of $\underline{A} \otimes \underline{B}$ by taking every p 'th row starting with the first, and so on. Since \underline{B} has p rows, the i 'th block of rows of $(\underline{A} \otimes \underline{B})_{\underline{m}, \underline{p}}$ is $\underline{A} \otimes \underline{\beta}'_i$, for $\underline{\beta}'_i$ of (1) so that

$$\underline{I}_{\underline{m}, \underline{p}}(\underline{A} \otimes \underline{B}) = (\underline{A} \otimes \underline{B})_{\underline{m}, \underline{p}} = \{\underline{A} \otimes \underline{\beta}'_i\}_{i=1 \dots p} \quad (35)$$

$$= \begin{bmatrix} \underline{A} \otimes \underline{\beta}'_1 \\ \vdots \\ \underline{A} \otimes \underline{\beta}'_p \end{bmatrix} = \{a_{\underline{j}} \otimes \underline{\beta}'_i\}_{\substack{i=1 \dots p \\ j=1 \dots n}} = \{a_{\underline{j}} \underline{\beta}'_i\}_{\substack{i=1 \dots p \\ j=1 \dots n}} \quad (36)$$

Notice that $(\underline{A} \otimes \underline{B})_{\underline{m}, \underline{p}}$ affords a compact notation for $\begin{bmatrix} \underline{A} \otimes \underline{\beta}'_1 \\ \vdots \\ \underline{A} \otimes \underline{\beta}'_p \end{bmatrix}$.

Applying (35) and (36) to the right-hand side of (34) yields the further equalities:

$$\begin{aligned} \underline{I}_{\underline{m}, \underline{p}}(\underline{A} \otimes \underline{B}) &= (\underline{B} \otimes \underline{A})_{\underline{n}, \underline{q}} \\ &= (\underline{B}' \otimes \underline{A}')'_{\underline{q}, \underline{n}} = \{\underline{B}' \otimes \underline{a}'_j\}'_{j=1 \dots n} = \{\underline{B} \otimes \underline{a}_j\}_{j=1 \dots n} \quad (37) \end{aligned}$$

$$= [\underline{B} \otimes \underline{a}_1 \cdots \underline{B} \otimes \underline{a}_n] = \{\underline{\beta}'_i \otimes \underline{a}_j\}_{\substack{i=1 \dots p \\ j=1 \dots n}} = \{a_{\underline{j}} \underline{\beta}'_i\}_{\substack{i=1 \dots p \\ j=1 \dots n}} \quad (38)$$

These equalities lead to the equivalent descriptive formulations of $\underline{I}_{\underline{m}, \underline{n}}$, (15), (16), (17) and (18), on substituting $\underline{A} = \underline{I}_{\underline{m}}$ and $\underline{B} = \underline{I}_{\underline{n}}$ in (35) - (38) to give $\underline{I}_{\underline{m}, \underline{n}} = (\underline{I}_{\underline{m}} \otimes \underline{I}_{\underline{n}})_{\underline{m}, \underline{n}}$.

4.5. Reversing the order of Kronecker products of vectors

The special cases of $\underline{\underline{A}}$ and/or $\underline{\underline{B}}$ being vectors are worthy of note. For column vectors $\underline{\underline{a}}$ and $\underline{\underline{b}}$ of order m and p , respectively, putting $n = 1$, then $q = 1$ and then $n = q = 1$, in (26) gives

$$(\underline{\underline{a}} \otimes \underline{\underline{B}})_{m,p} = \underline{\underline{I}}_{m,p} (\underline{\underline{a}} \otimes \underline{\underline{B}}) = \underline{\underline{B}} \otimes \underline{\underline{a}}, \quad (39)$$

$$(\underline{\underline{A}} \otimes \underline{\underline{b}})_{m,p} = \underline{\underline{I}}_{m,p} (\underline{\underline{A}} \otimes \underline{\underline{b}}) = \underline{\underline{b}} \otimes \underline{\underline{A}}, \quad (40)$$

and

$$(\underline{\underline{a}} \otimes \underline{\underline{b}})_{m,p} = \underline{\underline{I}}_{m,p} (\underline{\underline{a}} \otimes \underline{\underline{b}}) = \underline{\underline{b}} \otimes \underline{\underline{a}}. \quad (41)$$

Transposing (39) - (41), or putting $p = 1$ and $m = 1$ in (26), gives the corresponding results for row vectors. Using $n = 1$ and $p = 1$ in (26), (36) and (38) reveals the familiar commutativity property of Kronecker products of vectors:

$$\underline{\underline{a}} \otimes \underline{\underline{b}}' = \underline{\underline{a}} \underline{\underline{b}}' = \underline{\underline{b}}' \otimes \underline{\underline{a}}.$$

4.6. The vec-permutation matrix and Kronecker products of 3 or more matrices

Extending the vec-permutation matrix to cyclically permute order in Kronecker products of three or more matrices is straightforward, as indicated by Magnus and Neudecker [1979]. Balestra [1976, p. 24] introduced $\underline{\underline{I}}_{mp,s}$, for use in this connection but perhaps surprisingly did not explicitly give its use to cyclically permute a Kronecker product of three matrices. We present this result by immediate application of (25) to $(\underline{\underline{A}} \otimes \underline{\underline{B}}) \otimes \underline{\underline{C}} = \underline{\underline{A}} \otimes (\underline{\underline{B}} \otimes \underline{\underline{C}})$ for $\underline{\underline{A}}$, $\underline{\underline{B}}$ and $\underline{\underline{C}}$ of order $m \times n$, $p \times q$ and $s \times t$, respectively:

$$\underset{\sim}{C} \otimes \underset{\sim}{A} \otimes \underset{\sim}{B} = \underset{\sim}{I}_{\underset{\sim}{m}, \underset{\sim}{p}, \underset{\sim}{s}} [(\underset{\sim}{A} \otimes \underset{\sim}{B}) \otimes \underset{\sim}{C}] \underset{\sim}{I}_{\underset{\sim}{t}, \underset{\sim}{n}, \underset{\sim}{q}} \quad (42)$$

$$= \underset{\sim}{I}_{\underset{\sim}{p}, \underset{\sim}{m}, \underset{\sim}{s}} [\underset{\sim}{B} \otimes (\underset{\sim}{C} \otimes \underset{\sim}{A})] \underset{\sim}{I}_{\underset{\sim}{n}, \underset{\sim}{t}, \underset{\sim}{q}}$$

$$= \underset{\sim}{I}_{\underset{\sim}{p}, \underset{\sim}{m}, \underset{\sim}{s}} \underset{\sim}{I}_{\underset{\sim}{m}, \underset{\sim}{p}, \underset{\sim}{s}} [\underset{\sim}{A} \otimes (\underset{\sim}{B} \otimes \underset{\sim}{C})] \underset{\sim}{I}_{\underset{\sim}{q}, \underset{\sim}{t}, \underset{\sim}{n}} \underset{\sim}{I}_{\underset{\sim}{n}, \underset{\sim}{t}, \underset{\sim}{q}} \quad (43)$$

$$= (\underset{\sim}{I}_{\underset{\sim}{m}, \underset{\sim}{s}} \otimes \underset{\sim}{I}_{\underset{\sim}{p}}) (\underset{\sim}{I}_{\underset{\sim}{m}} \otimes \underset{\sim}{I}_{\underset{\sim}{p}, \underset{\sim}{s}}) [\underset{\sim}{A} \otimes (\underset{\sim}{B} \otimes \underset{\sim}{C})] (\underset{\sim}{I}_{\underset{\sim}{n}} \otimes \underset{\sim}{I}_{\underset{\sim}{t}, \underset{\sim}{q}}) (\underset{\sim}{I}_{\underset{\sim}{t}, \underset{\sim}{n}} \otimes \underset{\sim}{I}_{\underset{\sim}{q}}). \quad (44)$$

Properties of vec-permutation matrices with the same set of three indices, developed by Balestra [1976, pp. 24-25] and Magnus and Neudecker [1979] with lengthy algebra, are now shown to be easy consequences of (42) - (44): put $n = q = t = 1$ in (42) - (44), so that $\underset{\sim}{A} \otimes \underset{\sim}{B} \otimes \underset{\sim}{C}$ becomes $\underset{\sim}{a} \otimes \underset{\sim}{b} \otimes \underset{\sim}{c}$ and the final vec-permutation matrices on the right-hand sides of (42) - (44) reduce to $\underset{\sim}{I}$, giving

$$\begin{aligned} \underset{\sim}{I}_{\underset{\sim}{m}, \underset{\sim}{p}, \underset{\sim}{s}} (\underset{\sim}{a} \otimes \underset{\sim}{b} \otimes \underset{\sim}{c}) &= \underset{\sim}{I}_{\underset{\sim}{p}, \underset{\sim}{m}, \underset{\sim}{s}} \underset{\sim}{I}_{\underset{\sim}{m}, \underset{\sim}{p}, \underset{\sim}{s}} (\underset{\sim}{a} \otimes \underset{\sim}{b} \otimes \underset{\sim}{c}) \\ &= (\underset{\sim}{I}_{\underset{\sim}{m}, \underset{\sim}{s}} \otimes \underset{\sim}{I}_{\underset{\sim}{p}}) (\underset{\sim}{I}_{\underset{\sim}{m}} \otimes \underset{\sim}{I}_{\underset{\sim}{p}, \underset{\sim}{s}}) (\underset{\sim}{a} \otimes \underset{\sim}{b} \otimes \underset{\sim}{c}). \end{aligned} \quad (45)$$

Let $\underset{\sim}{a} \otimes \underset{\sim}{b} \otimes \underset{\sim}{c}$ take in turn the columns of $\underset{\sim}{I}_{\underset{\sim}{m}, \underset{\sim}{p}, \underset{\sim}{s}}$, and hence

$$\underset{\sim}{I}_{\underset{\sim}{m}, \underset{\sim}{p}, \underset{\sim}{s}} = \underset{\sim}{I}_{\underset{\sim}{p}, \underset{\sim}{m}, \underset{\sim}{s}} \underset{\sim}{I}_{\underset{\sim}{m}, \underset{\sim}{p}, \underset{\sim}{s}} = (\underset{\sim}{I}_{\underset{\sim}{m}, \underset{\sim}{s}} \otimes \underset{\sim}{I}_{\underset{\sim}{p}}) (\underset{\sim}{I}_{\underset{\sim}{m}} \otimes \underset{\sim}{I}_{\underset{\sim}{p}, \underset{\sim}{s}}). \quad (46)$$

Since $\underset{\sim}{I}_{\underset{\sim}{m}, \underset{\sim}{p}, \underset{\sim}{s}} = \underset{\sim}{I}_{\underset{\sim}{p}, \underset{\sim}{m}, \underset{\sim}{s}}$, interchanging m and p in (46) yields

$$\underset{\sim}{I}_{\underset{\sim}{m}, \underset{\sim}{p}, \underset{\sim}{s}} = \underset{\sim}{I}_{\underset{\sim}{p}, \underset{\sim}{m}, \underset{\sim}{s}} = \underset{\sim}{I}_{\underset{\sim}{m}, \underset{\sim}{p}, \underset{\sim}{s}} \underset{\sim}{I}_{\underset{\sim}{p}, \underset{\sim}{m}, \underset{\sim}{s}} = (\underset{\sim}{I}_{\underset{\sim}{p}, \underset{\sim}{s}} \otimes \underset{\sim}{I}_{\underset{\sim}{m}}) (\underset{\sim}{I}_{\underset{\sim}{p}} \otimes \underset{\sim}{I}_{\underset{\sim}{m}, \underset{\sim}{s}}). \quad (47)$$

Postmultiplying the middle two equalities in (47) by $\underset{\sim}{I}_{\underset{\sim}{s}, \underset{\sim}{m}, \underset{\sim}{p}}$ gives, using (22),

$$\underset{\sim}{I}_{\underset{\sim}{m}, \underset{\sim}{p}, \underset{\sim}{s}} \underset{\sim}{I}_{\underset{\sim}{p}, \underset{\sim}{m}, \underset{\sim}{s}} \underset{\sim}{I}_{\underset{\sim}{s}, \underset{\sim}{m}, \underset{\sim}{p}} = \underset{\sim}{I}_{\underset{\sim}{m}, \underset{\sim}{p}, \underset{\sim}{s}}. \quad (48)$$

This development, based on (45), is in contrast to the lengthy manipulations of Balestra [1976] who uses (18) to establish (46) - (48), and Magnus and Neudecker [1979] who give (42) and use (20) to develop (48) and all of (47) except its final equality.

A special case of (42) is

$$(\underset{\sim}{a} \otimes \underset{\sim}{b}' \otimes \underset{\sim}{C})_{m,s} = \underset{\sim}{b}' \otimes \underset{\sim}{C} \otimes \underset{\sim}{a} . \quad (49)$$

Using $\underset{\sim}{a} \otimes \underset{\sim}{b}' = \underset{\sim}{ab}'$, (49) becomes

$$(\underset{\sim}{ab}' \otimes \underset{\sim}{C})_{m,s} = \underset{\sim}{b}' \otimes \underset{\sim}{C} \otimes \underset{\sim}{a} \quad (50)$$

which, applied on

$$\underset{\sim}{I}_{m,n} = (\underset{\sim}{I}_m \otimes \underset{\sim}{I}_n)_{m,n} = \left(\sum_{i=1}^m \underset{\sim}{e}_i \underset{\sim}{e}'_i \otimes \underset{\sim}{I}_n \right)_{m,n} , \quad (51)$$

with $\underset{\sim}{C} = \underset{\sim}{I}_n$ and $\underset{\sim}{a} = \underset{\sim}{b} = \underset{\sim}{e}_i$, yields the Magnus and Neudecker [1979] formulations for $\underset{\sim}{I}_{m,n}$ in (20).

Generalization to Kronecker products of four or more matrices and vec-permutation matrices with four or more indices is straightforward.

ACKNOWLEDGMENTS

Grateful thanks go to Robert Anderson, Stephen Barnett, Paul Dwyer, Robert Hartwig and Friedrich Pukelsheim for suggested improvements to early drafts of this paper.

REFERENCES

- Aitken, A. C. [1949]. On the Wishart distribution in statistics. Biometrika, 36, 59-62.
- Anderson, R. D. [1978]. Studies on the estimation of variance components. Ph.D. Thesis, Cornell University. 136 pp.
- Anderson, R. D., Quaas, R. L., and Searle, S. R. [1977]. Fourth moments in the general linear model; and the variance of translation invariant quadratic forms. Paper No. BU-630-M in the Biometrics Unit Mimeo Series, Cornell University. 11 pp.
- Balestra, P. [1976]. La Dérivation Matricielle. Collection de l'Institut de Mathématiques Economiques, No. 12, Sirey, Paris.
- Barnett, S. [1973]. Matrix differential equations and Kronecker products. SIAM J. Appl. Math., 24, 1-5.
- Barnett, Stephen [1979]. Matrix Methods for Engineers and Scientists. McGraw-Hill, U. K.
- Bell, E. T. [1937]. Men of Mathematics. Simon and Schuster, New York.
- Bentler, P. M. and Lee, Sik-Yum [1975]. Some extensions of matrix calculus. General Systems, 20, 145-150.
- Brewer, John W. [1977]. The derivative of the Riccotti Matrix with respect to a matrix. IEEE Trans. Automat. Contr. AC-22, 980-998.
- Brewer, John W. [1978]. Kronecker products and matrix calculus in system theory. IEEE Trans. Cir. Syst. CAS-25, 772-781.
- Browne, M. W. [1974]. Generalized least squares estimators in the analysis of covariance structures. South African Statist. J., 8, 1-24.

- Cole, J. W. L. [1969]. Multivariate analysis of variance using patterned covariance matrices. Ph.D. Thesis, University of North Carolina. 303 pp.
- Conlisk, J. [1969]. The equilibrium covariance matrix of dynamic econometric models. J. Amer. Statist. Assoc., 64, 277-279.
- Conlisk, J. [1976]. A further note on stability in a random coefficients model. Internat. Econom. Rev., 17, 759-764.
- Giguère, M. A. and Styan, G. P. H. [1974]. Multivariate normal estimation with missing data on several variates. Trans. 1974 European Meeting of Statisticians, Prague.
- Graybill, F. A. [1969]. Introduction to Matrices with Applications in Statistics. Wadsworth, Belmont, California.
- Halmos, Paul R. [1958]. Finite-dimensional Vector Spaces. (Second edition). D. Van Nostrand Company, Princeton, New Jersey.
- Hartwig, Robert E. [1972]. The resultant and the matrix equation $\underline{AX} = \underline{XB}$. SIAM J. Appl. Math., 22, 538-544.
- Hartwig, R. E. [1975]. $\underline{AX} - \underline{XB} = \underline{C}$, resultants and generalized inverses. SIAM J. Appl. Math., 28, 154-183.
- Hartwig, R. E. and Morris, S. B. [1975]. The universal flip matrix and the generalized faro-shuffle. Pacific J. Math., 58, 445-455.
- Henderson, H. V. and Searle, S. R. [1979]. Vec and vech operators for matrices, with some uses in Jacobians and multivariate statistics. Canad. J. Statist., 7, 65-81.
- Hensel, K. [1891]. Über die Darstellung der Determinante eines Systems, welches aus Zwei Anderen componirt ist. Acta Math., 14, 317-319.

- Koopmans, T. C., Rubin, H., and Leipnik, R. B. [1950]. Measuring the equation systems of dynamic economics. Statistical Inference in Dynamic Economic Models. (Ed. T. C. Koopmans). John Wiley and Sons, New York.
- Kucera, V. [1974]. The matrix equation $\underline{AX} + \underline{XB} = \underline{C}$. SIAM J. Appl. Math., 26, 15-25.
- Lancaster, Peter [1969]. Theory of Matrices. Academic Press, New York.
- Lancaster, P. [1970]. Explicit solutions of linear matrix equations. SIAM Rev., 12, 544-566.
- Ledermann, W. [1936]. On singular pencils of Zehfuss, compound, and Schläflian matrices. Proc. Roy. Soc. Edinburgh, LVI, 58-89.
- Legault-Giguère, M. A. [1974]. Multivariate normal estimation with missing data. M.Sc. Thesis, Dept. of Math., McGill University, Montreal.
- Loewy, A. [1910]. Kombinatorik, Determinanten und Matrices. Reperitorium der Höheren Mathematik, 1, 43-167. E. Pascal, Teubner.
- MacDuffee, C. C. [1933]. The Theory of Matrices. Verlag von Julius Springer, Berlin.
- MacRae, E. C. [1974]. Matrix derivatives with an application to an adaptive linear decision problem. Ann. Statist., 2, 337-346.
- Magnus, J. R. and Neudecker, H. [1979]. The commutation matrix: Some properties and applications. Ann. Statist., 7, 381-394.
- McDonald, R. P. [1976]. The McDonald-Swaminathan matrix calculus: Clarifications, extensions, and illustrations. General Systems, 21, 87-94.

- McDonald, R. P. and Swaminathan, H. [1973]. A simple matrix calculus with applications to multivariate analysis. General Systems, 18, 37-54.
- Morris, S. B. and Hartwig, R. E. [1976]. The generalized faro shuffle. Discrete Math., 15, 333-346.
- Mitra, S. K. [1977]. The matrix equation $\underline{AXB} + \underline{CXD} = \underline{E}$. SIAM J. Appl. Math., 32, 823-825.
- Murnaghan, Francis D. [1938]. The Theory of Group Representations. Johns Hopkins Press, Baltimore.
- Nel, D. G. [1978]. On the symmetric multivariate normal distribution and asymptotic expansion of a Wishart matrix. South African Statist. J., 12, 145-159.
- Neudecker, H. [1968]. The Kronecker matrix product and some of its applications in econometrics. Statistica Neerlandica, 22, 69-81.
- Neudecker, H. [1969a]. A note on Kronecker matrix products and matrix equation systems. SIAM J. Appl. Math., 17, 603-606.
- Neudecker, H. [1969b]. Some theorems on matrix differentiation with special reference to Kronecker matrix products. J. Amer. Statist. Assoc., 64, 953-963.
- Nissen, D. H. [1968]. A note on the variance of a matrix. Econometrica, 36, 603-604.
- Pukelsheim, Friedrich [1977]. On Hsu's model in regression analysis. Math. Operationsforsch. Statist. Ser. Statist., 8, 323-331.

- Rogers, G. S. and Young, D. L. [1978]. On testing a multivariate linear hypothesis when the covariance matrix and its inverse have the same pattern. J. Amer. Statist. Assoc., 73, 203-207.
- Roth, W. E. [1934]. On direct product matrices. Bull. Amer. Math. Soc., 40, 461-468.
- Rutherford, D. E. [1933]. On the condition that two Zehfuss matrices be equal. Bull. Amer. Math. Soc., 39, 801-808.
- Searle, S. R. [1966]. Matrix Algebra for the Biological Sciences. John Wiley and Sons, New York.
- Searle, S. R. [1978]. A univariate formulation of the multivariate linear model. Contributions to Survey Sampling and Applied Statistics, Papers in Honor of H. O. Hartley (Ed. H. A. David). Academic Press, 181-189.
- Searle, S. R. [1979]. Notes on variance component estimation: A detailed account of maximum likelihood and kindred methodology. Paper No. BU-673-M in the Biometrics Unit Mimeo Series, Cornell University.
- Singh, R. P. [1972]. Some generalizations in matrix differentiation with applications in multivariate analysis. Ph.D. Thesis, University of Windsor, Canada. 209 pp.
- Swain, A. J. [1975]. Analysis of parametric structures for variance matrices. Ph.D. Thesis, Dept. of Statistics, University of Adelaide, Australia. 188 pp.

- Sylvester, J. [1884a]. Sur la solution du cas le plus général des équations linéaires en quantités binaires, c'est-à-dire en quaternions ou en matrices du second ordre. C. R. Acad. Sci. Paris., 99, 117-118.
- Sylvester, J. [1884b]. Sur la résolution générale de l'équation linéaire en matrices d'un ordre quelconque. C. R. Acad. Sci. Paris., 99, 409-412.
- Sylvester, J. [1884c]. Sur la résolution générale de l'équation linéaire en matrices d'un ordre quelconque. C. R. Acad. Sci. Paris., 99, 432-436.
- Tracy, D. S. and Dwyer, P. S. [1969]. Multivariate maxima and minima with matrix derivatives. J. Amer. Statist. Assoc., 64, 1576-1594.
- Tracy, D. S. and Singh, R. P. [1972a]. A new matrix product and its applications in partitioned matrices. Statistica Neerlandica, 26, 143-157.
- Tracy, D. S. and Singh, R. P. [1972b]. Some modifications of matrix differentiation for evaluating Jacobians of symmetric matrix transformations. Symmetric Functions in Statistics, Proceedings of a symposium in honor of Paul S. Dwyer (Ed. Derrick S. Tracy), University of Windsor, Ontario, Canada, 203-224.
- Vartak, Nanohar N. [1955]. On an application of Kronecker product of matrices to statistical designs. Ann. Math. Statist., 26, 420-438.

- Vetter, W. J. [1970]. Derivative operations on matrices. IEEE Trans. Automat. Contr., 15, 241-244.
- Vetter, W. J. [1973]. Matrix calculus operations and Taylor expansions. SIAM Rev., 15, 352-369.
- Vetter, W. J. [1975]. Linear structures and solutions of linear matrix equations. Linear Algebra and Appl., 10, 181-188.
- Whelchel, J. E., Jr. and Guinn, D. F. [1970]. Properties of high radix fast Fourier-Hadamard transforms. Applications of Walsh functions. 1970 Proceedings symposium and workshop held at Naval Research Laboratory, Washington, D. C. (Ed. C. A. Bass). 80-87.
- Zehfuss, G. [1858]. Ueber eine gewisse determinante. Z. Math. Phys., 3, 298-301.

Table 1: Definitions, notations and names for what, in this paper, is called the vec-permutation matrix, $\tilde{I}_{m,n}$.

Definition	Writer	Notation	Name	
Reversing Kronecker products	(a) vectors	Ledermann [1936]	\tilde{P} and \tilde{Q}	Permutation matrix
	(b) matrices	Conlisk [1976] (incorrect)	\tilde{M} for $\tilde{I}_{n,n}$	Permutation matrix
		Murnaghan [1938]	\tilde{P}	Permutation matrix
		Vartak [1955]	\tilde{P} and \tilde{Q}	Permutation matrix
		Searle [1966]	\tilde{P} and \tilde{Q}	Elementary operators
		Whelchel and Guinn [1970]	\tilde{S}_{n^2} for $\tilde{I}_{n,n}$	Shuffle matrix
Defining row permutations	Tracy and Dwyer [1969]	}	$\tilde{I}(n)$	Permuted identity matrix
	Tracy and Singh [1972a,b]			
	Singh [1972]			
The derivative $\partial \tilde{X}' / \partial \tilde{X}$	McDonald and Swaminathan [1973]	\tilde{E}	Permutation matrix	
	Bentler and Lee [1975]	\tilde{E}		
	McDonald [1976]	$\tilde{E}_{n,m}$		
Description of elements	Vetter [1970]	$\langle \tilde{U}_{ij}^{ji} \rangle$		
	Hartwig [1972] (incorrect)	\tilde{P}	Permutation matrix	
	Vetter [1973, 1975]	$\tilde{E}_{n \times m}^{m \times n}$	Permutation matrix	
	MacRae [1974]	$\tilde{I}(m,n)$	Permuted identity matrix	
	Hartwig and Morris [1975]	$\tilde{P}_{m,n}$	Universal flip matrix	
	Swain [1975]	\tilde{L}_{mn}	Permutation matrix	
	Balestra [1976]	$\tilde{P}_{m,n}$	Permuted identity matrix	
	Brewer [1977, 1978]	$\tilde{U}_{n \times m}$	Permutation matrix	
	Magnus and Neudecker [1979]	\tilde{K}_{nm}	Commutation matrix	
	$\text{vec} \tilde{A}$ related to $\text{vec} \tilde{A}'$	Barnett [1973]	\tilde{P} and \tilde{Q}	Permutation matrix
		Henderson and Searle [1979]	$\tilde{I}(m,n)$	vec-permutation matrix
Henderson and Searle (this paper)		$\tilde{I}_{m,n}$	vec-permutation matrix	