SCHOOL OF OPERATIONS RESEARCH
AND INDUSTRIAL ENGINEERING
COLLEGE OF ENGINEERING
CORNELL UNIVERSITY
ITHACA, NEW YORK 14853

TECHNICAL REPORT NO. 814

August 1988

A SINGLE-STAGE SELECTION PROCEDURE
FOR MULTI-FACTOR MULTINOMIAL EXPERIMENTS
WITH MULTIPLICATIVITY

by

Robert E. Bechhofer

David M. Goldsman^{*}

and

Christopher Jennison^{**}

Approved for public release; distributions unlimited

Research partially supported by the U.S. Army Research Office through the Mathematical Sciences Institute of Cornell University.

^{*} School of Industrial and Systems Engineering, Georgia Institute of Technology, Atlanta, GA 30332.

^{**} School of Mathematical Sciences, University of Bath, Bath BA2 7AY, United Kingdom.

A SINGLE-STAGE SELECTION PROCEDURE
FOR MULTI-FACTOR MULTINOMIAL EXPERIMENTS
WITH MULTIPLICATIVITY

Robert E. Bechhofer

School of Operations Research
and Industrial Engineering
Cornell University
Ithaca, New York 14853

David M. Goldsman

School of Industrial
and Systems Engineering
Georgia Institute of Technology
Atlanta, Georgia 30332

Christopher Jennison

School of Mathematical Sciences
University of Bath
Bath, BA2 7AY
UNITED KINGDOM

Key Words and Phrases: multinomial selection problem; multiplicative probability matrix; selection procedures; ranking procedures; single-stage procedures; market research sampling.

ABSTRACT

A single-stage procedure is proposed for selecting the event which has the largest probability in multi-factor multinomial experiments with multiplicativity, i.e., experiments in which the factor-level responses of any one factor are independent of those of all other factors. The procedure is a generalization of one proposed for single-factor multinomial experiments by Bechhofer, Elmaghraby and Morse (B-E-M) (1959). Tables necessary to implement the procedure are provided, and properties of the procedure are obtained. It is shown that if independence holds it is much more efficient in terms of minimizing total sample size to conduct the experiment as a factorial experiment rather than as independent single-factor experiments. Specifically, if an f -factor ($f \geq 2$) experiment is conducted, and certain symmetry conditions hold, the factorial experiment requires $1/f$ as many observations to guarantee the same indifference-zone probability requirement as do f independent single-factor experiments. If multiplicability does not hold the experimenter can use the original B-E-M single-stage, single-factor procedure but for a different goal and probability requirement than the one employed for the 2-factor problem.

1. INTRODUCTION AND SUMMARY

A single-stage procedure for selecting the multinomial event which has the largest probability was proposed by Bechhofer, Elmaghraby and Morse (1959); in that article the so-called indifference-zone approach was adopted, with "distance" between the event probabilities being measured in terms of their ratios. An open fully sequential procedure for this same problem was described in Bechhofer, Kiefer and Sobel (B-K-S) (1968), Section 5.3.3. Cacoullos and Sobel (1966) and Alam (1971) also studied this problem, and proposed closed and open sequential procedures, respectively; later Ramey and Alam (R-A) (1979) devised a closed composite of the C-S and Alam procedures. Bechhofer and Goldsman (B-G) (1985a) studied the R-A procedure in depth, and proposed (B-G (1985b, 1986)) a truncated version of the B-K-S procedure which was shown to have performance characteristics which in most practical circumstances are superior to those of the R-A procedure and, in fact, to all of the procedures proposed earlier.

B-G (1988a) also showed how the B-K-S fully sequential selection procedure could be generalized to accommodate multi-factor experiments involving Koopman-Darmois (K-D) populations with additivity. The rationale for conducting such multi-factor experiments rather than independent single-factor experiments is that in many situations the former will achieve the same goal and guarantee the same probability requirement as the latter, but will do so with a smaller cost of sampling; this was demonstrated conclusively in B-G (1988a) for the normal means selection problem. In that article it also was pointed out (as a special case) that a multinomial population with multiplicativity can, after an appropriate transformation, be represented as a multivariate K-D population with additivity, and hence that the generalized B-K-S sequential selection procedure could be used for multi-factor multinomial experiments with multiplicativity.

However, there are many real-life situations in which sequential procedures are not feasible and/or economical, and single-stage procedures may represent the only option available to the experimenter. Thus it is our objective in the present paper to devise a single-stage selection procedure which would be appropriate for use in multi-factor experiments with

multiplicativity. This procedure would therefore be a generalization of the B-E-M single-stage selection procedure which was designed for use in single-factor experiments. In a later paper we plan to compare the performance characteristics of this new single-stage procedure with those of the generalized B-K-S sequential procedure when both are used for multi-factor multinomial experiments with multiplicativity. We will then be able to determine the relative performance of these two procedures when both are feasible options of the experimenter.

We state the problem in Section 2 and describe our single-stage procedure in Section 3. Section 4 contains some results concerning the probability of a correct selection ($P\{CS\}$) of the procedure, including a key factorization theorem which leads to the calculation of the least-favorable configuration of the cell probabilities. Tables of the $P\{CS\}$ for single-factor experiments, necessary to implement two-factor experiments with multiplicativity, are given in Section 5. The justification for using multi-factor experiments when multiplicativity can be assumed is given in Section 6. The procedure is applied to a practical problem involving market research sampling in Section 7. An alternate method of experimentation when multiplicativity is not present is described in Section 8. Section 9 contains our concluding remarks.

2. STATEMENT OF THE PROBLEM

We pose the problem and illustrate the methods and results only for 2-factor experiments, the generalization to multi-factor experiments being straightforward.

Let $\{X_{ijr} \mid (1 \leq i \leq a, 1 \leq j \leq b)\}$ $r = 1, 2, \dots$ denote independent matrix-observations from a single 2-factor (bivariate) multinomial population Π having an unknown probability matrix $\{p_{ij}\} = \{p_{ij} \mid (1 \leq i \leq a, 1 \leq j \leq b)\}$.

Here p_{ij} ($p_{ij} \geq 0, \sum_{i=1}^a \sum_{j=1}^b p_{ij} = 1$) denotes the probability that $X_{ijr} = 1$ ($r = 1, 2, \dots$) on the r th matrix-observation; if $X_{ijr} = 1$ then $X_{i'j'r} = 0$ for all $i' \neq i$ or $j' \neq j$. Denote the marginal success probabilities for the row and column factors by $\alpha_i = \sum_{j=1}^b p_{ij}$ ($1 \leq i \leq a$) and $\beta_j = \sum_{i=1}^a p_{ij}$ ($1 \leq j \leq b$), respectively. Then $\sum_{i=1}^a \alpha_i = \sum_{j=1}^b \beta_j = 1$. We assume that $p_{ij} = \alpha_i \cdot \beta_j$ (all i, j), i.e., that the factor-level responses for one factor are

independent of those for the other factor. Then $\{\sum_{j=1}^b X_{ijr} \ (1 \leq i \leq a)\}$
 $(\{\sum_{i=1}^a X_{ijr} \ (1 \leq j \leq b)\}) \ r = 1, 2, \dots$ can be regarded as independent
vector-observations from a single-factor a -level ($a \geq 2$) (b -level ($b \geq 2$))
multinomial population with a probability vector $\alpha = (\alpha_1, \dots, \alpha_a)$
 $(\beta = (\beta_1, \dots, \beta_b))$. Let $\alpha_{[1]} \leq \dots \leq \alpha_{[a]}$ and $\beta_{[1]} \leq \dots \leq \beta_{[b]}$ denote the
ordered values of the α_i and β_j , respectively. It is assumed that the values
of the $\alpha_i, \beta_j, \alpha_{[u]}, \beta_{[v]} \ (1 \leq i, u \leq a, \ 1 \leq j, v \leq b)$ are unknown, and that the
pairing of the α_i, β_j with the $\alpha_{[u]}, \beta_{[v]} \ (1 \leq i, u \leq a, \ 1 \leq j, v \leq b)$ is
completely unknown.

The goal of the experimenter is to select the event associated with $\alpha_{[a]}$
and $\beta_{[b]}$. When a procedure selects this event we say that a correct selection
(CS) has been made. Consideration is limited to procedures which guarantee
the following indifference-zone probability requirement:

$$P\{CS\} \geq P^* \text{ whenever } \begin{cases} \alpha_{[a]} \geq \theta_\alpha^* \alpha_{[a-1]} \\ \text{and} \\ \beta_{[b]} \geq \theta_\beta^* \beta_{[b-1]} \end{cases} \quad (2.1)$$

where $\{\theta_\alpha^*, \theta_\beta^*, P^*\}$ with $1 < \theta_\alpha^*, \theta_\beta^* < \infty, \ 1/ab < P^* < 1$ are specified prior
to the start of experimentation. The open sequential procedure $(\mathcal{P}_{B(2,+)}^*)$ of
B-G (1988a), as applied to the present problem (referred to as Case M_2^+ in
Section 4.4 of that article), guarantees (2.1).

After n matrix-observations have been observed let $y_{ijn} = \sum_{r=1}^n x_{ijr}$,
 $A_{in} = \sum_{j=1}^b y_{ijn}$ and $B_{jn} = \sum_{i=1}^a y_{ijn}$. We denote the ordered values of the
 A_{in} and B_{jn} by $A_{[1]n} \leq \dots \leq A_{[a]n}$ and $B_{[1]n} \leq \dots \leq B_{[b]n}$. Equal
 A_{in} -values and equal B_{jn} -values can occur with positive probabilities; when
this happens they should be arranged in any definite order.

3. SINGLE-STAGE PROCEDURE FOR 2-FACTOR MULTINOMIAL EXPERIMENTS WITH MULTIPLICATIVITY

We now describe the procedure.

Sampling rule: Take n independent matrix-observations $\{X_{ijr} \mid (1 \leq i \leq a, 1 \leq j \leq b)\}$, $r = 1, 2, \dots, n$ in a single-stage from Π . (3.1)

Terminal decision rule: Calculate $A_{[a]n}$ and $B_{[b]n}$. If exactly s ($1 \leq s \leq a$) of the A_{in} are tied for largest, i.e., $A_{[a-s]n} < A_{[a-s+1]n} = \dots = A_{[a]n}$, and exactly t ($1 \leq t \leq b$) of the B_{jn} are tied for largest, i.e., $B_{[b-t]n} < B_{[b-t+1]n} = \dots = B_{[b]n}$, then select one of these $s \cdot t$ event combinations at random, and assert that it is the event associated with $\alpha_{[a]}$ and $\beta_{[b]}$. (3.2)

4. PROBABILITY OF A CORRECT SELECTION

4.1 A general expression for the $P\{CS\}$

We now give a general expression for the $P\{CS\}$.

$$P_{(a,b)}^{(n)}\{CS|\alpha,\beta\} = \sum \frac{1}{s} \cdot \frac{1}{t} \frac{n!}{\prod_{i=1}^a \prod_{j=1}^b (y_{ijn}!)} \prod_{i=1}^a \prod_{j=1}^b p_{ij}^{y_{ijn}}. \quad (4.1)$$

Here the sum is over all matrices

$$\{y_n\} = \begin{pmatrix} y_{11n} & \cdots & y_{1bn} \\ \vdots & & \vdots \\ y_{a1n} & \cdots & y_{abn} \end{pmatrix}$$

such that $\sum_{i=1}^a \sum_{j=1}^b y_{ijn} = n$, and

$$A_{(a)n} \geq A_{(i)n} \quad (1 \leq i \leq a-1),$$

$$B_{(b)n} \geq B_{(j)n} \quad (1 \leq j \leq b-1),$$

where the $A_{(i)n}$ ($B_{(j)n}$) are the totals associated with $\alpha_{[i]}$ ($\beta_{[j]}$), and s and t , which are functions of $\{y_n\}$, are the number of $A_{(i)n}$'s and $B_{(j)n}$'s, respectively, tied for largest.

4.2 Calculation of an exact $P\{CS\}$

We now illustrate the calculation of the exact $P\{CS\}$ for a particular (a,b,n) when the single-stage procedure, (3.1) and (3.2), is used.

Example 4.1: $a = 2$, $b = 3$, $n = 3$

The cell probabilities are given in matrices (4.2) and (4.3) where (4.2) represents the general model while (4.3) represents the multiplicative model:

$$\begin{pmatrix} p_{11}, & p_{12}, & p_{13} \\ p_{21}, & p_{22}, & p_{23} \end{pmatrix} \quad (4.2)$$

$$\begin{pmatrix} \alpha_1\beta_1, & \alpha_1\beta_2, & \alpha_1\beta_3 \\ \alpha_2\beta_1, & \alpha_2\beta_2, & \alpha_2\beta_3 \end{pmatrix} \quad (4.3)$$

Suppose that $\alpha_2 = \alpha_{[2]}$ and $\beta_3 = \beta_{[3]}$ so that p_{23} is the largest cell probability. All 12 possible outcomes that can lead to a correct selection using the single-stage rule are enumerated in TABLE I. Also listed is the $P\{CS\}$ associated with each outcome.

Summing the terms in the third column yields a polynomial in the α_i ($i = 1,2$) and β_j ($j = 1,2,3$) which is an expression for the exact $P\{CS\}$ in the multiplicative case for $a = 2$, $b = 3$, $n = 3$. For arbitrary (a,b,n) we denote this $P\{CS\}$ by $P_{(a,b)}^{(n)}\{CS|\alpha,\beta\}$. Thus

TABLE I

Outcomes leading to a correct selection and associated $P\{CS\}$
for $a = 2$, $b = 3$, $n = 3$

Matrices (outcomes) leading to a correct selection	$P\{CS\}$ associated with each outcome expressed in terms of the p_{ij}	$P\{CS\}$ associated with each outcome expressed in terms of the α_i and β_j
$\begin{Bmatrix} 0 & 0 & 0 \\ 0 & 0 & 3 \end{Bmatrix}$	p_{23}^3	$\alpha_2^3 \beta_3^3$
$\begin{Bmatrix} 0 & 0 & 0 \\ 0 & 1 & 2 \end{Bmatrix}$	$3p_{22}p_{23}^2$	$3\alpha_2^3 \beta_2 \beta_3^2$
$\begin{Bmatrix} 0 & 0 & 0 \\ 1 & 0 & 2 \end{Bmatrix}$	$3p_{21}p_{23}^2$	$3\alpha_2^3 \beta_1 \beta_3^2$
$\begin{Bmatrix} 0 & 0 & 1 \\ 0 & 0 & 2 \end{Bmatrix}$	$3p_{13}p_{23}^2$	$3\alpha_1 \alpha_2^2 \beta_3^3$
$\begin{Bmatrix} 0 & 1 & 0 \\ 0 & 0 & 2 \end{Bmatrix}$	$3p_{12}p_{23}^2$	$3\alpha_1 \alpha_2^2 \beta_2 \beta_3^2$
$\begin{Bmatrix} 1 & 0 & 0 \\ 0 & 0 & 2 \end{Bmatrix}$	$3p_{11}p_{23}^2$	$3\alpha_1 \alpha_2^2 \beta_1 \beta_3^2$
$\begin{Bmatrix} 0 & 0 & 0 \\ 1 & 1 & 1 \end{Bmatrix}$	$6p_{21}p_{22}p_{23}^{(1/3)}$	$2\alpha_2^3 \beta_1 \beta_2 \beta_3$
$\begin{Bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \end{Bmatrix}$	$6p_{11}p_{22}p_{23}^{(1/3)}$	$2\alpha_1 \alpha_2^2 \beta_1 \beta_2 \beta_3$
$\begin{Bmatrix} 0 & 0 & 1 \\ 0 & 1 & 1 \end{Bmatrix}$	$6p_{13}p_{22}p_{23}$	$6\alpha_1 \alpha_2^2 \beta_2 \beta_3^2$
$\begin{Bmatrix} 0 & 0 & 1 \\ 1 & 0 & 1 \end{Bmatrix}$	$6p_{13}p_{21}p_{23}$	$6\alpha_1 \alpha_2^2 \beta_1 \beta_3^2$
$\begin{Bmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \end{Bmatrix}$	$6p_{12}p_{21}p_{23}^{(1/3)}$	$2\alpha_1 \alpha_2^2 \beta_1 \beta_2 \beta_3$
$\begin{Bmatrix} 0 & 0 & 1 \\ 1 & 1 & 0 \end{Bmatrix}$	$6p_{21}p_{22}p_{13}^{(1/3)}$	$2\alpha_1 \alpha_2^2 \beta_1 \beta_2 \beta_3$

$$\begin{aligned}
P_{(2,3)}^{(3)}\{CS|\alpha,\beta\} &= \alpha_2^3\beta_3^3 + 3\alpha_2^3\beta_2\beta_3^2 + 3\alpha_2^3\beta_1\beta_3^2 \\
&\quad + 3\alpha_1\alpha_2^2\beta_3^3 + 9\alpha_1\alpha_2^2\beta_2\beta_3^2 + 9\alpha_1\alpha_2^2\beta_1\beta_3^2 \\
&\quad + 2\alpha_2^3\beta_1\beta_2\beta_3 + 6\alpha_1\alpha_2^2\beta_1\beta_2\beta_3.
\end{aligned} \tag{4.4}$$

Now (4.4) can be written as

$$P_{(2,3)}^{(3)}\{CS|\alpha,\beta\} = P_{(2)}^{(3)}\{CS|\alpha\} \times P_{(3)}^{(3)}\{CS|\beta\} \tag{4.5}$$

where

$$P_{(2)}^{(3)}\{CS|\alpha\} = \alpha_2^2(1 + 2\alpha_1) \tag{4.6a}$$

and

$$P_{(3)}^{(3)}\{CS|\beta\} = \beta_3(\beta_3 + 2\beta_1\beta_3 + 2\beta_2\beta_3 + 2\beta_1\beta_2). \tag{4.6b}$$

Here $P_{(a)}^{(n)}\{CS|\alpha\}$ ($P_{(b)}^{(n)}\{CS|\beta\}$) is the exact $P\{CS\}$ associated with a single-factor multinomial experiment, as in B-E-M (1959), with n observations from an a -level (b -level) multinomial distribution with probability vector α (β). It is clear that the calculation of exact probabilities such as (4.4) would be extremely tedious, even for very moderate values of (a,b,n) . The following result on the factorization of the $P\{CS\}$ allows us to address this issue using single-factor calculations in Section 5.

4.3 Factorization of the $P\{CS\}$ under multiplicativity

The fact that (4.4) factors into a product of two terms, the first (second) being the $P\{CS\}$ associated with 3 observations from a 2-level (3-level) multinomial distribution with $\alpha = (\alpha_1, \alpha_2)$ ($\beta = (\beta_1, \beta_2, \beta_3)$) is not an accident. In fact, this factorization occurs for all (a,b,n) with $a, b \geq 2$. The general result is given as Theorem 4.1 below.

Theorem 4.1: Suppose that the single-stage procedure with sampling rule (3.1) and terminal decision rule (3.2) is used with n independent matrix-observations from a 2-factor multinomial population for which $p_{ij} = \alpha_i \cdot \beta_j$ ($1 \leq i \leq a$, $1 \leq j \leq b$). Then

$$P_{(a,b)}^{(n)}\{CS|\alpha,\beta\} = P_{(a)}^{(n)}\{CS|\alpha\} \times P_{(b)}^{(n)}\{CS|\beta\} \quad (4.7)$$

for all (a,b,n) with $a,b \geq 2, n \geq 1$.

Proof of theorem: Let $\{Y_{ij} = \sum_{r=1}^n X_{ijr} \ (1 \leq i \leq a, 1 \leq j \leq b)\}$ denote the random matrix based on n independent matrix-observations $\{X_{ijr} \ (1 \leq i \leq a, 1 \leq j \leq b)\}$, $r = 1, 2, \dots, n$, and let $Y_{i.} = \sum_{j=1}^b Y_{ij} \ (1 \leq i \leq a)$ and $Y_{.j} = \sum_{i=1}^a Y_{ij} \ (1 \leq j \leq b)$ denote the marginal totals. Let $y_{ij}, y_{i.}$ and $y_{.j}$ be the corresponding observed values. (In the foregoing, for simplicity of notation, we have omitted the subscript n in Y_{ijn} and y_{ijn} .)

We first show for given $\{y_{i.} \ (1 \leq i \leq a), y_{.j} \ (1 \leq j \leq b)\}$ that

$$\begin{aligned} & P\{Y_{i.} = y_{i.} \ (1 \leq i \leq a) \text{ and } Y_{.j} = y_{.j} \ (1 \leq j \leq b)\} \\ &= P\{Y_{i.} = y_{i.} \ (1 \leq i \leq a)\} \times P\{Y_{.j} = y_{.j} \ (1 \leq j \leq b)\}. \end{aligned} \quad (4.8)$$

To see this let n_1 (n_2) denote the number of ways in which we can order the n responses for factor 1 (2) to obtain $Y_{i.} = y_{i.} \ (1 \leq i \leq a) \ (Y_{.j} = y_{.j} \ (1 \leq j \leq b))$. Thus,

$$n_1 = \frac{n!}{\prod_{i=1}^a y_{i.}!} \text{ and } n_2 = \frac{n!}{\prod_{j=1}^b y_{.j}!}.$$

Hence, the number of ways in which we can order the n factor 1 and factor 2 responses to obtain the marginal totals $\{y_{i.} \ (1 \leq i \leq a), y_{.j} \ (1 \leq j \leq b)\}$ is $n_1 \times n_2$. Now

$$\begin{aligned} & P\{Y_{i.} = y_{i.} \ (1 \leq i \leq a) \text{ and } Y_{.j} = y_{.j} \ (1 \leq j \leq b)\} \\ &= \sum_{\substack{n_1 \times n_2 \\ \text{sequences} \\ \text{of trials}}} \prod_i \prod_j (p_{ij})^{y_{ij}} \end{aligned} \quad (4.9)$$

where the sum is over the $n_1 \times n_2$ sequences of n trials leading to matrices with marginal totals $\{y_{i\cdot} \ (1 \leq i \leq a), \ y_{\cdot j} \ (1 \leq j \leq b)\}$. Thus, under multiplicativity, (4.9) becomes

$$\begin{aligned}
& \sum_{\substack{n_1 \times n_2 \\ \text{sequences} \\ \text{of trials}}} \prod_i \prod_j \alpha_i^{y_{ij}} \beta_j^{y_{ij}} &= \sum_{\substack{n_1 \times n_2 \\ \text{sequences} \\ \text{of trials}}} \prod_i \alpha_i^{y_{i\cdot}} \prod_j \beta_j^{y_{\cdot j}} \\
&= n_1 n_2 \prod_i \alpha_i^{y_{i\cdot}} \prod_j \beta_j^{y_{\cdot j}} \\
&\quad (\text{since the probability is the same for each sequence}) \\
&= n_1 \prod_i \alpha_i^{y_{i\cdot}} \times n_2 \prod_j \beta_j^{y_{\cdot j}} \\
&= P\{Y_{i\cdot} = y_{i\cdot} \ (1 \leq i \leq a)\} \times P\{Y_{\cdot j} = y_{\cdot j} \ (1 \leq j \leq b)\}
\end{aligned}$$

as required. This factorization holds for an arbitrary set of marginal totals, not necessarily those that lead to a correct selection.

To complete the proof let $Y_{(i)\cdot}$ ($Y_{\cdot(j)}$) denote the random total associated with $\alpha_{[i]}$ ($1 \leq i \leq a$) ($\beta_{[j]}$ ($1 \leq j \leq b$)). We define the events E_s and F_t where

$$\begin{aligned}
E_s &= \{Y_{i\cdot} \ (1 \leq i \leq a): Y_{(a)\cdot} \text{ is tied for largest with } s-1 \text{ others}\}, \\
F_t &= \{Y_{\cdot j} \ (1 \leq j \leq b): Y_{\cdot(b)} \text{ is tied for largest with } t-1 \text{ others}\}.
\end{aligned}$$

Then

$$\begin{aligned}
P\{\text{CS for factor 1}\} &= \sum_{s=1}^a P\{E_s\} \cdot \frac{1}{s} \\
P\{\text{CS for factor 2}\} &= \sum_{t=1}^b P\{F_t\} \cdot \frac{1}{t}.
\end{aligned}$$

Thus,

$P\{\text{CS for factor 1 and factor 2}\}$

$$= \sum_{s=1}^a \sum_{t=1}^b \sum_{\substack{\{y_{i.}, y_{.j}\} \\ E_s \text{ and } F_t \text{ hold}}} P\{Y_{i.} = y_{i.} \ (1 \leq i \leq a) \text{ and } Y_{.j} = y_{.j} \ (1 \leq j \leq b)\} \times \frac{1}{st}. \quad (4.10)$$

Using the factorization (4.8), the right hand side of (4.10) becomes

$$\begin{aligned} & \sum_{s=1}^a \sum_{t=1}^b \sum_{\substack{\{y_{i.}, y_{.j}\} \\ E_s \text{ and } F_t \text{ hold}}} P\{Y_{i.} = y_{i.} \ (1 \leq i \leq a)\} \\ & \quad \times P\{Y_{.j} = y_{.j} \ (1 \leq j \leq b)\} \times \frac{1}{st} \\ &= \sum_{s=1}^a \sum_{t=1}^b \sum_{\substack{\{y_{i.}\} \\ E_s \text{ holds}}} \sum_{\substack{\{y_{.j}\} \\ F_t \text{ holds}}} P\{Y_{i.} = y_{i.} \ (1 \leq i \leq a)\} \\ & \quad \times P\{Y_{.j} = y_{.j} \ (1 \leq j \leq b)\} \times \frac{1}{st} \\ &= \sum_{s=1}^a \sum_{\substack{\{y_{i.}\} \\ E_s \text{ holds}}} P\{Y_{i.} = y_{i.} \ (1 \leq i \leq a)\} \cdot \frac{1}{s} \\ & \quad \times \sum_{t=1}^b \sum_{\substack{\{y_{.j}\} \\ E_t \text{ holds}}} P\{Y_{.j} = y_{.j} \ (1 \leq j \leq b)\} \cdot \frac{1}{t} \\ &= P\{\text{CS for factor 1}\} \times P\{\text{CS for factor 2}\}. \quad \text{QED} \end{aligned}$$

4.4 Least favorable configuration

In order to implement the single-stage procedure we seek the smallest value of n that will guarantee (2.1). The first step in the search is the determination of those probability vectors α, β which, for given (a, b, n) , will minimize $P_{(a, b)}^{(n)}\{\text{CS} | \alpha, \beta\}$ subject to the constraints specified in (2.1). The elements of these minimizing vectors are called the least favorable (LF) configuration of the α_i and β_j . An important consequence of Theorem 4.1 is that, because of the factorization, the determination of these elements reduces

the problem to one that had been solved earlier by Kesten and Morse (1959) for single-factor experiments. Using their result (K-M, p. 120), it follows that this LF-configuration, which is independent of n , is given for our two-factor experiment by

$$\begin{aligned}\alpha_{[a]} &= \theta_{\alpha}^* \alpha_{[i]} \quad (1 \leq i \leq a-1) \\ \beta_{[b]} &= \theta_{\beta}^* \beta_{[j]} \quad (1 \leq j \leq b-1)\end{aligned}\tag{4.11}$$

which, since $\sum_{i=1}^a \alpha_{[i]} = \sum_{j=1}^b \beta_{[j]} = 1$, implies

$$\begin{aligned}\alpha_{[1]} &= \dots = \alpha_{[a-1]} = 1/(\theta_{\alpha}^* + a - 1), \quad \alpha_{[a]} = \theta_{\alpha}^*/(\theta_{\alpha}^* + a - 1) \\ \beta_{[1]} &= \dots = \beta_{[b-1]} = 1/(\theta_{\beta}^* + b - 1), \quad \beta_{[b]} = \theta_{\beta}^*/(\theta_{\beta}^* + b - 1).\end{aligned}\tag{4.12}$$

Thus, e.g., when (4.11) holds we have for Example 4.1 that

$$P_{(2,3)}^{(3)}\{\text{CS} | (\theta_{\alpha}^*, \theta_{\beta}^*), \text{LF}\} = P_{(2)}^{(3)}\{\text{CS} | \theta_{\alpha}^*, \text{LF}\} \times P_{(3)}^{(3)}\{\text{CS} | \theta_{\beta}^*, \text{LF}\}$$

where

$$P_{(2)}^{(3)}\{\text{CS} | \theta_{\alpha}^*, \text{LF}\} = \frac{(\theta_{\alpha}^*)^2}{(\theta_{\alpha}^* + 1)^3} \{\theta_{\alpha}^* + 3\}\tag{4.13}$$

and

$$P_{(3)}^{(3)}\{\text{CS} | \theta_{\beta}^*, \text{LF}\} = \frac{\theta_{\beta}^*}{(\theta_{\beta}^* + 2)^3} \{(\theta_{\beta}^*)^2 + 6\theta_{\beta}^* + 2\}.\tag{4.14}$$

In the same way it is possible to find exact expressions for $P_{(a)}^{(n)}\{\text{CS} | \theta^*, \text{LF}\}$ for given a , n and θ^* . This $P\{\text{CS}\}$ increases and approaches unity with n . Using tables of these probabilities for the present independence case, $p_{ij} = \alpha_i \cdot \beta_j$ (all i, j), we can determine the smallest n necessary to guarantee (2.1) for given (a, b) and specified $(\theta_{\alpha}^*, \theta_{\beta}^*, P^*)$ provided that the available tables extend to n -values which are sufficiently large. In Section 5 we provide such tables for $a = 2(1)5$, selected θ^* and $n \leq 200$.

5. TABLES OF $P_{(a)}^{(n)}\{CS|\theta^*, LF\}$ FOR SINGLE-FACTOR EXPERIMENTS

Tables of the type just described for single-factor experiments are contained in B-E-M (1959). In that article Tables A-2, A-3 and A-4 give the exact $P_{(a)}^{(n)}\{CS|\theta^*, LF\}$ for $a = 2, 3, 4$, respectively, $n = 1(1)30$ and $\theta^* = 1.02(0.02)1.10(0.10)2.00(0.20)3.00, 10.00$. The tables were calculated using polynomials like (4.13) and (4.14) for the (a, n, θ^*) of interest. Two large-sample normal approximations to the exact probabilities are also given and their use illustrated.

Tables II, III and IV of the present paper extend Tables A-2, A-3 and A-4, respectively, of B-E-M (1959) to $n = 1(1)70(2)140(4)200$ for $\theta^* = 1.1, 1.2(0.2)2.2, 3.0$. Our present Table V for $a = 5$ covers this same range of arguments. These tables were calculated in double precision using an iterative procedure on an IBM 4381 computer at Georgia Institute of Technology. The entries should be correct to the five decimal places reported.

Gibbons, Olkin and Sobel (G-O-S) (1977), pp. 445-447, provides extensive tables of the smallest n necessary to guarantee $P_{(a)}^{(n)}\{CS|\theta^*, LF\} \geq P^*$ for all combinations of (a, θ^*, P^*) with $a = 2(1)10$, $\theta^* = 1.2(0.2)3.0$ and $P^* = 0.75, 0.90, 0.95, 0.975, 0.999$. As will be seen in Section 7, these single-factor tables are not in a form that is appropriate for our purposes. (See, however, Section 8.)

6. SAVINGS ARISING FROM MULTI-FACTOR EXPERIMENTATION UNDER MULTIPLICATIVITY

The motivation for using 2-factor multinomial experimentation rather than independent single-factor multinomial experimentation when multiplicativity is present follows from Theorem 6.1. An analogous result holds for f -factor experiments ($f > 2$).

TABLE II

 $P_{(a)}^{(n)}\{\text{CS}|\theta^*, \text{LF}\}$ as a function of n for $a = 2$ and selected θ^*

n	θ^*							
	1.1	1.2	1.4	1.6	1.8	2.0	2.2	3.0
1	.52381	.54545	.58333	.61538	.64286	.66667	.68750	.75000
2	.52381	.54545	.58333	.61538	.64286	.66667	.68750	.75000
3	.53569	.56799	.62384	.67000	.70845	.74074	.76807	.84375
4	.53569	.56799	.62384	.67000	.70845	.74074	.76807	.84375
5	.54457	.58476	.65338	.70879	.75364	.79012	.81999	.89648
6	.54457	.58476	.65338	.70879	.75364	.79012	.81999	.89648
7	.55196	.59861	.67731	.73939	.78821	.82670	.85718	.92944
8	.55196	.59861	.67731	.73939	.78821	.82670	.85718	.92944
9	.55842	.61064	.69767	.76473	.81600	.85515	.88514	.95107
10	.55842	.61064	.69767	.76473	.81600	.85515	.88514	.95107
11	.56421	.62137	.71548	.78633	.83897	.87791	.90677	.96567
12	.56421	.62137	.71548	.78633	.83897	.87791	.90677	.96567
13	.56951	.63112	.73136	.80508	.85830	.89646	.92381	.97571
14	.56951	.63112	.73136	.80508	.85830	.89646	.92381	.97571
15	.57442	.64011	.74569	.82155	.87479	.91177	.93741	.98270
16	.57442	.64011	.74569	.82155	.87479	.91177	.93741	.98270
17	.57901	.64846	.75875	.83618	.88898	.92452	.94836	.98761
18	.57901	.64846	.75875	.83618	.88898	.92452	.94836	.98761
19	.58334	.65628	.77075	.84926	.90129	.93523	.95725	.99110
20	.58334	.65628	.77075	.84926	.90129	.93523	.95725	.99110
21	.58744	.66366	.78183	.86102	.91204	.94428	.96451	.99358
22	.58744	.66366	.78183	.86102	.91204	.94428	.96451	.99358
23	.59134	.67063	.79211	.87165	.92145	.95195	.97046	.99535
24	.59134	.67063	.79211	.87165	.92145	.95195	.97046	.99535
25	.59508	.67727	.80169	.88129	.92974	.95849	.97537	.99663
26	.59508	.67727	.80169	.88129	.92974	.95849	.97537	.99663
27	.59866	.68359	.81064	.89007	.93706	.96407	.97942	.99755
28	.59866	.68359	.81064	.89007	.93706	.96407	.97942	.99755
29	.60211	.68964	.81904	.89809	.94354	.96886	.98278	.99822
30	.60211	.68964	.81904	.89809	.94354	.96886	.98278	.99822
31	.60543	.69544	.82693	.90542	.94929	.97298	.98556	.99870
32	.60543	.69544	.82693	.90542	.94929	.97298	.98556	.99870
33	.60864	.70101	.83436	.91215	.95441	.97652	.98789	.99905
34	.60864	.70101	.83436	.91215	.95441	.97652	.98789	.99905
35	.61176	.70637	.84137	.91833	.95897	.97958	.98982	.99930
36	.61176	.70637	.84137	.91833	.95897	.97958	.98982	.99930
37	.61477	.71154	.84800	.92402	.96305	.98222	.99144	.99949
38	.61477	.71154	.84800	.92402	.96305	.98222	.99144	.99949
39	.61771	.71653	.85427	.92927	.96669	.98450	.99280	.99963
40	.61771	.71653	.85427	.92927	.96669	.98450	.99280	.99963

TABLE II (continued)

 $P_{\{a\}}^{(n)}\{\text{CS}|\theta^*, \text{LF}\}$ as a function of n for $a = 2$ and selected θ^*

n	θ^*							
	1.1	1.2	1.4	1.6	1.8	2.0	2.2	3.0
41	.62056	.72136	.86022	.93411	.96995	.98648	.99393	.99973
42	.62056	.72136	.86022	.93411	.96995	.98648	.99393	.99973
43	.62334	.72603	.86586	.93858	.97287	.98820	.99488	.99980
44	.62334	.72603	.86586	.93858	.97287	.98820	.99488	.99980
45	.62605	.73056	.87123	.94272	.97550	.98970	.99568	.99985
46	.62605	.73056	.87123	.94272	.97550	.98970	.99568	.99985
47	.62869	.73495	.87633	.94656	.97786	.99100	.99635	.99989
48	.62869	.73495	.87633	.94656	.97786	.99100	.99635	.99989
49	.63127	.73922	.88118	.95011	.97998	.99213	.99692	.99992
50	.63127	.73922	.88118	.95011	.97998	.99213	.99692	.99992
51	.63380	.74337	.88581	.95341	.98188	.99311	.99740	.99994
52	.63380	.74337	.88581	.95341	.98188	.99311	.99740	.99994
53	.63627	.74740	.89022	.95647	.98360	.99397	.99780	.99996
54	.63627	.74740	.89022	.95647	.98360	.99397	.99780	.99996
55	.63869	.75133	.89443	.95932	.98515	.99472	.99814	.99997
56	.63869	.75133	.89443	.95932	.98515	.99472	.99814	.99997
57	.64106	.75515	.89845	.96196	.98655	.99537	.99842	.99998
58	.64106	.75515	.89845	.96196	.98655	.99537	.99842	.99998
59	.64339	.75888	.90229	.96443	.98781	.99594	.99866	.99998
60	.64339	.75888	.90229	.96443	.98781	.99594	.99866	.99998
61	.64567	.76252	.90596	.96672	.98895	.99644	.99887	.99999
62	.64567	.76252	.90596	.96672	.98895	.99644	.99887	.99999
63	.64791	.76606	.90947	.96885	.98998	.99688	.99904	.99999
64	.64791	.76606	.90947	.96885	.98998	.99688	.99904	.99999
65	.65011	.76953	.91283	.97084	.99091	.99726	.99919	.99999
66	.65011	.76953	.91283	.97084	.99091	.99726	.99919	.99999
67	.65227	.77291	.91605	.97270	.99175	.99760	.99931	.99999
68	.65227	.77291	.91605	.97270	.99175	.99760	.99931	.99999
69	.65439	.77621	.91913	.97443	.99251	.99789	.99941	.99999+
70	.65439	.77621	.91913	.97443	.99251	.99789	.99941	.99999+
72	.65648	.77944	.92209	.97604	.99320	.99815	.99950	
74	.65854	.78260	.92492	.97755	.99382	.99837	.99958	
76	.66056	.78569	.92764	.97896	.99439	.99857	.99964	
78	.66256	.78872	.93024	.98027	.99490	.99874	.99969	
80	.66452	.79168	.93274	.98150	.99537	.99889	.99974	
82	.66645	.79458	.93515	.98265	.99579	.99903	.99978	
84	.66836	.79742	.93745	.98373	.99617	.99914	.99981	
86	.67024	.80020	.93967	.98473	.99652	.99925	.99984	
88	.67209	.80293	.94180	.98568	.99683	.99934	.99986	
90	.67392	.80561	.94384	.98656	.99712	.99942	.99988	

TABLE II (continued)
 $P_{(a)}^{(n)}\{\text{CS}|\theta^*, \text{LF}\}$ as a function of n for $a = 2$ and selected θ^*

n	θ^*							
	1.1	1.2	1.4	1.6	1.8	2.0	2.2	3.0
92	.67572	.80823	.94581	.98738	.99738	.99949	.99990	
94	.67750	.81081	.94770	.98815	.99761	.99955	.99992	
96	.67926	.81333	.94952	.98888	.99783	.99960	.99993	
98	.68099	.81581	.95127	.98955	.99802	.99965	.99994	
100	.68270	.81825	.95296	.99019	.99820	.99969	.99995	
102	.68439	.82064	.95458	.99078	.99836	.99973	.99996	
104	.68607	.82298	.95614	.99134	.99851	.99976	.99996	
106	.68772	.82529	.95764	.99187	.99864	.99979	.99997	
108	.68935	.82755	.95909	.99236	.99876	.99981	.99997	
110	.69096	.82977	.96049	.99282	.99887	.99984	.99998	
112	.69256	.83196	.96183	.99325	.99897	.99985	.99998	
114	.69413	.83411	.96312	.99365	.99906	.99987	.99998	
116	.69569	.83622	.96437	.99403	.99914	.99989	.99999	
118	.69724	.83830	.96557	.99439	.99922	.99990	.99999	
120	.69876	.84034	.96673	.99473	.99929	.99991	.99999	
122	.70027	.84235	.96785	.99504	.99935	.99992	.99999	
124	.70177	.84432	.96893	.99533	.99941	.99993	.99999	
126	.70325	.84626	.96997	.99561	.99946	.99994	.99999	
128	.70471	.84818	.97097	.99587	.99951	.99995	.99999	
130	.70616	.85006	.97193	.99612	.99955	.99995	.99999	
132	.70760	.85191	.97287	.99635	.99959	.99996	.99999+	
134	.70902	.85373	.97377	.99656	.99963	.99996		
136	.71042	.85552	.97464	.99677	.99966	.99997		
138	.71182	.85729	.97547	.99696	.99969	.99997		
140	.71320	.85903	.97628	.99714	.99972	.99997		
144	.71592	.86242	.97782	.99746	.99976	.99998		
148	.71859	.86572	.97925	.99775	.99980	.99998		
152	.72121	.86892	.98058	.99801	.99984	.99999		
156	.72379	.87202	.98182	.99823	.99986	.99999		
160	.72633	.87503	.98299	.99843	.99989	.99999		
164	.72882	.87796	.98407	.99861	.99990	.99999		
168	.73127	.88080	.98508	.99877	.99992	.99999+		
172	.73367	.88357	.98603	.99891	.99993			
176	.73604	.88625	.98691	.99903	.99994			
180	.73838	.88886	.98773	.99914	.99995			
184	.74067	.89140	.98851	.99923	.99996			
188	.74294	.89388	.98923	.99932	.99997			
192	.74516	.89628	.98990	.99940	.99997			
196	.74736	.89862	.99053	.99946	.99998			
200	.74952	.90090	.99112	.99952	.99998			

TABLE III

 $P_{\{a\}}^{(n)}\{\text{CS}|\theta^*, \text{LF}\}$ as a function of n for $a = 3$ and selected θ^*

n	θ^*							
	1.1	1.2	1.4	1.6	1.8	2.0	2.2	3.0
1	.35484	.37500	.41176	.44444	.47368	.50000	.52381	.60000
2	.35484	.37500	.41176	.44444	.47368	.50000	.52381	.60000
3	.36222	.38965	.44026	.48560	.52617	.56250	.59508	.69600
4	.36699	.39880	.45702	.50846	.55379	.59375	.62901	.73440
5	.36952	.40395	.46738	.52370	.57342	.61719	.65568	.76896
6	.37361	.41200	.48260	.54487	.59925	.64648	.68742	.80352
7	.37722	.41904	.49574	.56290	.62100	.67090	.71362	.83117
8	.37940	.42344	.50434	.57510	.63602	.68799	.73210	.85052
9	.38261	.42977	.51624	.59136	.65539	.70935	.75453	.87181
10	.38556	.43553	.52688	.60566	.67214	.72751	.77328	.88845
11	.38753	.43950	.53455	.61628	.68483	.74145	.78779	.90141
12	.39024	.44485	.54448	.62954	.70015	.75775	.80426	.91460
13	.39279	.44984	.55361	.64155	.71381	.77207	.81851	.92539
14	.39461	.45349	.56054	.65090	.72463	.78352	.82997	.93399
15	.39700	.45820	.56918	.66216	.73723	.79646	.84255	.94252
16	.39928	.46266	.57724	.67251	.74865	.80802	.85362	.94962
17	.40097	.46605	.58358	.68085	.75798	.81755	.86278	.95542
18	.40314	.47031	.59127	.69062	.76858	.82803	.87258	.96103
19	.40522	.47438	.59851	.69969	.77828	.83751	.88131	.96577
20	.40680	.47755	.60437	.70719	.78640	.84549	.88868	.96970
21	.40880	.48146	.61132	.71580	.79544	.85411	.89642	.97344
22	.41072	.48522	.61791	.72385	.80378	.86196	.90338	.97663
23	.41223	.48822	.62335	.73064	.81089	.86869	.90934	.97931
24	.41408	.49186	.62971	.73830	.81867	.87584	.91551	.98182
25	.41588	.49536	.63576	.74551	.82590	.88240	.92110	.98399
26	.41731	.49821	.64085	.75168	.83214	.88809	.92594	.98582
27	.41906	.50162	.64670	.75855	.83890	.89407	.93090	.98752
28	.42076	.50491	.65231	.76505	.84520	.89958	.93541	.98899
29	.42213	.50763	.65707	.77068	.85071	.90441	.93936	.99025
30	.42378	.51084	.66250	.77688	.85660	.90944	.94336	.99141
31	.42538	.51396	.66772	.78277	.86212	.91409	.94702	.99242
32	.42670	.51657	.67220	.78792	.86699	.91820	.95025	.99328
33	.42827	.51961	.67726	.79355	.87216	.92244	.95349	.99408
34	.42981	.52257	.68213	.79890	.87702	.92638	.95647	.99477
35	.43108	.52508	.68637	.80362	.88134	.92989	.95911	.99536
36	.43258	.52798	.69110	.80874	.88588	.93348	.96175	.99591
37	.43404	.53081	.69567	.81363	.89017	.93683	.96418	.99638
38	.43527	.53322	.69967	.81797	.89400	.93983	.96635	.99679
39	.43671	.53599	.70412	.82265	.89802	.94288	.96850	.99717
40	.43812	.53870	.70842	.82712	.90181	.94573	.97048	.99749

TABLE III (continued)

 $P_{(a)}^{(n)}\{\text{CS}|\theta^*, \text{LF}\}$ as a function of n for $a = 3$ and selected θ^*

n	θ^*							
	1.1	1.2	1.4	1.6	1.8	2.0	2.2	3.0
41	.43931	.54102	.71221	.83112	.90522	.94830	.97227	.99778
42	.44069	.54368	.71640	.83540	.90877	.95090	.97403	.99803
43	.44205	.54628	.72045	.83950	.91213	.95333	.97565	.99826
44	.44321	.54853	.72405	.84319	.91517	.95553	.97712	.99846
45	.44454	.55109	.72800	.84712	.91832	.95775	.97856	.99864
46	.44586	.55359	.73183	.85089	.92130	.95983	.97990	.99879
47	.44698	.55577	.73525	.85430	.92402	.96172	.98111	.99893
48	.44827	.55823	.73899	.85791	.92681	.96362	.98229	.99905
49	.44954	.56064	.74262	.86138	.92947	.96540	.98339	.99916
50	.45063	.56275	.74587	.86453	.93189	.96702	.98439	.99925
51	.45189	.56513	.74941	.86786	.93438	.96864	.98536	.99934
52	.45312	.56746	.75285	.87106	.93674	.97017	.98626	.99942
53	.45419	.56952	.75596	.87398	.93891	.97157	.98709	.99948
54	.45540	.57181	.75932	.87704	.94112	.97296	.98789	.99954
55	.45660	.57407	.76258	.88000	.94323	.97427	.98863	.99959
56	.45764	.57607	.76554	.88270	.94517	.97547	.98931	.99964
57	.45882	.57829	.76874	.88553	.94715	.97666	.98997	.99968
58	.45999	.58048	.77184	.88826	.94903	.97779	.99059	.99972
59	.46101	.58242	.77467	.89077	.95076	.97882	.99115	.99975
60	.46216	.58458	.77770	.89339	.95253	.97985	.99169	.99978
61	.46329	.58670	.78066	.89591	.95421	.98082	.99220	.99980
62	.46429	.58859	.78336	.89824	.95577	.98171	.99266	.99982
63	.46541	.59069	.78626	.90066	.95734	.98259	.99311	.99984
64	.46652	.59275	.78908	.90300	.95885	.98343	.99353	.99986
65	.46750	.59460	.79166	.90516	.96024	.98420	.99392	.99988
66	.46859	.59663	.79441	.90741	.96165	.98496	.99429	.99989
67	.46968	.59864	.79710	.90957	.96300	.98567	.99464	.99990
68	.47063	.60044	.79957	.91158	.96425	.98634	.99496	.99991
69	.47170	.60242	.80220	.91366	.96551	.98699	.99526	.99992
70	.47276	.60437	.80477	.91567	.96672	.98761	.99555	.99993
72	.47475	.60806	.80965	.91947	.96897	.98875	.99607	.99995
74	.47670	.61168	.81437	.92307	.97106	.98978	.99652	.99996
76	.47874	.61542	.81911	.92659	.97305	.99072	.99693	.99997
78	.48065	.61893	.82358	.92987	.97486	.99157	.99729	.99997
80	.48254	.62239	.82791	.93299	.97655	.99234	.99760	.99998
82	.48450	.62594	.83225	.93603	.97814	.99304	.99788	.99998
84	.48634	.62930	.83635	.93887	.97961	.99368	.99813	.99999
86	.48816	.63260	.84033	.94157	.98097	.99425	.99834	.99999
88	.49005	.63600	.84431	.94420	.98226	.99478	.99853	.99999
90	.49183	.63921	.84808	.94667	.98344	.99525	.99870	.99999

TABLE III (continued)

 $P_{(a)}^{(n)}\{\text{CS}|\theta^*, \text{LF}\}$ as a function of n for $a = 3$ and selected θ^*

n	θ^*							
	1.1	1.2	1.4	1.6	1.8	2.0	2.2	3.0
92	.49359	.64238	.85174	.94901	.98454	.99568	.99885	.99999
94	.49541	.64563	.85541	.95129	.98558	.99608	.99899	.99999+
96	.49714	.64871	.85888	.95343	.98654	.99643	.99910	
98	.49884	.65175	.86225	.95546	.98743	.99675	.99920	
100	.50061	.65486	.86562	.95744	.98828	.99705	.99930	
102	.50228	.65782	.86883	.95930	.98905	.99732	.99938	
104	.50394	.66074	.87194	.96107	.98978	.99756	.99945	
106	.50565	.66373	.87505	.96279	.99046	.99778	.99951	
108	.50728	.66658	.87801	.96441	.99109	.99798	.99957	
110	.50889	.66939	.88089	.96595	.99168	.99816	.99962	
112	.51055	.67227	.88376	.96745	.99223	.99833	.99966	
114	.51213	.67501	.88649	.96886	.99274	.99848	.99970	
116	.51370	.67772	.88915	.97021	.99322	.99861	.99973	
118	.51532	.68049	.89180	.97151	.99367	.99874	.99976	
120	.51686	.68313	.89434	.97274	.99408	.99885	.99979	
122	.51839	.68575	.89680	.97391	.99447	.99895	.99981	
124	.51997	.68842	.89925	.97505	.99484	.99905	.99984	
126	.52147	.69097	.90160	.97612	.99517	.99913	.99985	
128	.52296	.69349	.90388	.97715	.99549	.99921	.99987	
130	.52450	.69607	.90615	.97814	.99579	.99928	.99989	
132	.52596	.69854	.90833	.97908	.99606	.99934	.99990	
134	.52742	.70097	.91044	.97997	.99632	.99940	.99991	
136	.52892	.70346	.91255	.98084	.99656	.99946	.99992	
138	.53036	.70585	.91456	.98166	.99678	.99950	.99993	
140	.53178	.70821	.91653	.98244	.99699	.99955	.99994	
144	.53465	.71292	.92035	.98391	.99737	.99963	.99995	
148	.53747	.71753	.92399	.98526	.99770	.99969	.99996	
152	.54021	.72198	.92742	.98649	.99799	.99974	.99997	
156	.54296	.72640	.93072	.98761	.99825	.99979	.99998	
160	.54567	.73072	.93385	.98865	.99847	.99982	.99998	
164	.54829	.73491	.93682	.98959	.99866	.99985	.99998	
168	.55093	.73906	.93967	.99045	.99883	.99988	.99999	
172	.55354	.74313	.94238	.99124	.99897	.99990	.99999	
176	.55607	.74707	.94495	.99197	.99910	.99991	.99999	
180	.55861	.75098	.94742	.99263	.99921	.99993	.99999	
184	.56112	.75481	.94977	.99324	.99931	.99994	.99999+	
188	.56356	.75853	.95199	.99380	.99940	.99995		
192	.56601	.76222	.95413	.99431	.99947	.99996		
196	.56843	.76583	.95617	.99477	.99954	.99997		
200	.57079	.76935	.95810	.99520	.99960	.99997		

TABLE IV

 $P_{(a)}^{(n)}\{\text{CS}|\theta^*, \text{LF}\}$ as a function of n for $a = 4$ and selected θ^*

n	θ^*							
	1.1	1.2	1.4	1.6	1.8	2.0	2.2	3.0
1	.26829	.28571	.31818	.34783	.37500	.40000	.42308	.50000
2	.26829	.28571	.31818	.34783	.37500	.40000	.42308	.50000
3	.27308	.29543	.33790	.37741	.41406	.44800	.47940	.58333
4	.27775	.30469	.35583	.40314	.44661	.48640	.52273	.63889
5	.28040	.30998	.36622	.41824	.46594	.50944	.54898	.67361
6	.28269	.31470	.37595	.43284	.48502	.53248	.57542	.70833
7	.28569	.32077	.38803	.45039	.50727	.55859	.60457	.74306
8	.28830	.32602	.39834	.46519	.52581	.58010	.62830	.77006
9	.29041	.33032	.40697	.47774	.54167	.59859	.64875	.79313
10	.29245	.33452	.41549	.49020	.55740	.61682	.66876	.81470
11	.29469	.33907	.42454	.50312	.57337	.63496	.68828	.83427
12	.29674	.34324	.43277	.51483	.58775	.65119	.70560	.85103
13	.29858	.34701	.44032	.52563	.60103	.66614	.72150	.86592
14	.30035	.35067	.44768	.53614	.61389	.68050	.73660	.87941
15	.30223	.35450	.45525	.54677	.62667	.69454	.75114	.89162
16	.30399	.35811	.46238	.55675	.63860	.70755	.76451	.90241
17	.30563	.36149	.46909	.56615	.64980	.71972	.77691	.91203
18	.30722	.36477	.47563	.57527	.66062	.73135	.78865	.92071
19	.30887	.36816	.48229	.58443	.67131	.74268	.79992	.92859
20	.31046	.37141	.48866	.59316	.68144	.75332	.81041	.93560
21	.31195	.37449	.49472	.60146	.69103	.76335	.82020	.94189
22	.31340	.37750	.50065	.60954	.70030	.77294	.82948	.94756
23	.31490	.38059	.50665	.61761	.70943	.78226	.83837	.95270
24	.31635	.38357	.51244	.62536	.71814	.79108	.84670	.95731
25	.31773	.38642	.51799	.63279	.72645	.79942	.85451	.96144
26	.31908	.38922	.52344	.64003	.73448	.80742	.86192	.96518
27	.32046	.39207	.52893	.64724	.74238	.81517	.86900	.96856
28	.32181	.39484	.53425	.65420	.74994	.82253	.87566	.97160
29	.32310	.39752	.53939	.66090	.75718	.82953	.88193	.97434
30	.32437	.40014	.54443	.66744	.76420	.83624	.88788	.97681
31	.32566	.40281	.54950	.67394	.77108	.84273	.89357	.97905
32	.32692	.40541	.55444	.68023	.77769	.84892	.89892	.98106
33	.32814	.40793	.55923	.68631	.78404	.85481	.90398	.98288
34	.32934	.41042	.56394	.69225	.79020	.86047	.90878	.98452
35	.33056	.41293	.56866	.69815	.79623	.86594	.91336	.98601
36	.33175	.41540	.57327	.70387	.80204	.87116	.91769	.98735
37	.33291	.41779	.57775	.70941	.80763	.87615	.92178	.98855
38	.33405	.42015	.58216	.71484	.81306	.88093	.92566	.98965
39	.33521	.42254	.58658	.72021	.81837	.88556	.92937	.99064
40	.33635	.42489	.59091	.72544	.82349	.88998	.93287	.99153

TABLE IV (continued)

 $P_{(a)}^{(n)}\{\text{CS}|\theta^*, \text{LF}\}$ as a function of n for $a = 4$ and selected θ^*

n	θ^*							
	1.1	1.2	1.4	1.6	1.8	2.0	2.2	3.0
41	.33746	.42717	.59512	.73051	.82844	.89421	.93619	.99234
42	.33855	.42943	.59928	.73548	.83323	.89827	.93934	.99307
43	.33965	.43171	.60343	.74040	.83792	.90219	.94235	.99373
44	.34074	.43395	.60750	.74519	.84245	.90594	.94519	.99433
45	.34180	.43614	.61148	.74985	.84683	.90953	.94789	.99486
46	.34285	.43831	.61540	.75441	.85108	.91299	.95045	.99535
47	.34391	.44049	.61932	.75892	.85523	.91632	.95290	.99579
48	.34496	.44264	.62317	.76332	.85925	.91951	.95521	.99619
49	.34598	.44474	.62693	.76761	.86313	.92257	.95741	.99655
50	.34699	.44683	.63065	.77181	.86691	.92551	.95949	.99688
51	.34801	.44893	.63435	.77596	.87059	.92835	.96148	.99718
52	.34902	.45099	.63799	.78001	.87416	.93107	.96337	.99744
53	.35000	.45302	.64157	.78396	.87761	.93368	.96516	.99768
54	.35098	.45503	.64509	.78783	.88097	.93619	.96686	.99790
55	.35197	.45706	.64860	.79166	.88424	.93861	.96848	.99810
56	.35294	.45905	.65206	.79539	.88742	.94093	.97002	.99828
57	.35389	.46101	.65545	.79904	.89049	.94316	.97148	.99844
58	.35484	.46296	.65880	.80262	.89348	.94530	.97286	.99859
59	.35579	.46491	.66214	.80615	.89639	.94737	.97419	.99872
60	.35674	.46684	.66543	.80960	.89922	.94935	.97544	.99884
61	.35766	.46874	.66866	.81298	.90196	.95125	.97664	.99895
62	.35858	.47062	.67185	.81629	.90462	.95309	.97777	.99905
63	.35951	.47251	.67502	.81955	.90722	.95485	.97885	.99914
64	.36042	.47438	.67815	.82274	.90974	.95655	.97988	.99922
65	.36132	.47622	.68123	.82587	.91219	.95818	.98085	.99929
66	.36222	.47805	.68427	.82893	.91456	.95974	.98178	.99936
67	.36312	.47988	.68730	.83195	.91688	.96125	.98266	.99942
68	.36401	.48170	.69028	.83491	.91913	.96270	.98350	.99948
69	.36489	.48349	.69322	.83780	.92131	.96410	.98430	.99952
70	.36576	.48526	.69612	.84064	.92343	.96544	.98506	.99957
72	.36750	.48881	.70186	.84618	.92751	.96798	.98647	.99965
74	.36920	.49227	.70744	.85150	.93136	.97032	.98774	.99971
76	.37091	.49572	.71292	.85664	.93500	.97249	.98890	.99976
78	.37257	.49910	.71826	.86157	.93844	.97450	.98994	.99980
80	.37423	.50245	.72350	.86634	.94170	.97636	.99089	.99984
82	.37586	.50575	.72860	.87092	.94478	.97809	.99174	.99987
84	.37749	.50902	.73362	.87535	.94769	.97968	.99252	.99989
86	.37908	.51223	.73851	.87961	.95045	.98116	.99322	.99991
88	.38067	.51543	.74332	.88373	.95306	.98253	.99385	.99993
90	.38223	.51856	.74800	.88769	.95553	.98380	.99443	.99994

TABLE IV (continued)

 $P_{(a)}^{(n)}\{CS|\theta^*, LF\}$ as a function of n for $a = 4$ and selected θ^*

n	θ^*							
	1.1	1.2	1.4	1.6	1.8	2.0	2.2	3.0
92	.38379	.52168	.75261	.89152	.95786	.98498	.99495	.99995
94	.38532	.52475	.75710	.89521	.96007	.98607	.99542	.99996
96	.38685	.52780	.76152	.89877	.96217	.98708	.99585	.99997
98	.38835	.53080	.76583	.90220	.96415	.98801	.99624	.99997
100	.38985	.53378	.77007	.90551	.96602	.98888	.99659	.99998
102	.39133	.53671	.77421	.90870	.96780	.98969	.99691	.99998
104	.39280	.53963	.77828	.91179	.96948	.99043	.99720	.99998
106	.39425	.54251	.78225	.91476	.97108	.99112	.99746	.99999
108	.39570	.54536	.78616	.91764	.97258	.99176	.99769	.99999
110	.39713	.54818	.78998	.92041	.97401	.99236	.99791	.99999
112	.39855	.55098	.79373	.92309	.97537	.99291	.99810	.99999
114	.39996	.55374	.79741	.92567	.97665	.99342	.99828	.99999
116	.40136	.55649	.80101	.92817	.97786	.99389	.99844	.99999
118	.40274	.55919	.80454	.93057	.97901	.99433	.99858	.99999+
120	.40412	.56189	.80801	.93290	.98010	.99474	.99871	
122	.40548	.56454	.81141	.93514	.98114	.99512	.99883	
124	.40684	.56719	.81474	.93731	.98212	.99547	.99894	
126	.40818	.56979	.81801	.93941	.98304	.99580	.99904	
128	.40952	.57239	.82122	.94143	.98392	.99610	.99913	
130	.41084	.57495	.82436	.94338	.98475	.99638	.99921	
132	.41217	.57749	.82745	.94527	.98554	.99664	.99928	
134	.41347	.58001	.83048	.94709	.98629	.99688	.99935	
136	.41477	.58251	.83345	.94885	.98700	.99710	.99941	
138	.41606	.58498	.83636	.95055	.98767	.99731	.99946	
140	.41734	.58744	.83922	.95220	.98831	.99750	.99951	
144	.41988	.59228	.84479	.95532	.98948	.99785	.99960	
148	.42239	.59705	.85014	.95823	.99054	.99814	.99967	
152	.42487	.60173	.85530	.96095	.99149	.99840	.99973	
156	.42731	.60634	.86028	.96349	.99234	.99862	.99978	
160	.42973	.61087	.86507	.96586	.99311	.99881	.99981	
164	.43212	.61533	.86969	.96808	.99380	.99897	.99985	
168	.43449	.61972	.87414	.97015	.99442	.99911	.99987	
172	.43682	.62405	.87843	.97208	.99498	.99924	.99990	
176	.43914	.62831	.88257	.97389	.99548	.99934	.99991	
180	.44143	.63250	.88657	.97558	.99593	.99943	.99993	
184	.44369	.63663	.89042	.97716	.99634	.99951	.99994	
188	.44593	.64070	.89413	.97863	.99670	.99958	.99995	
192	.44815	.64471	.89772	.98001	.99703	.99963	.99996	
196	.45035	.64866	.90117	.98130	.99733	.99968	.99997	
200	.45253	.65256	.90451	.98250	.99759	.99973	.99997	

TABLE V

 $P_{\{a\}}^{(n)}\{\text{CS}|\theta^*, \text{LF}\}$ as a function of n for $a = 5$ and selected θ^*

n	θ^*							
	1.1	1.2	1.4	1.6	1.8	2.0	2.2	3.0
1	.21569	.23077	.25926	.28571	.31034	.33333	.35484	.42857
2	.21569	.23077	.25926	.28571	.31034	.33333	.35484	.42857
3	.21900	.23760	.27348	.30758	.33987	.37037	.39915	.49854
4	.22291	.24547	.28929	.33101	.37041	.40741	.44203	.55852
5	.22562	.25093	.30012	.34690	.39094	.43210	.47038	.59707
6	.22746	.25472	.30798	.35886	.40683	.45165	.49325	.62951
7	.22946	.25888	.31674	.37223	.42458	.47336	.51845	.66360
8	.23170	.26350	.32621	.38637	.44294	.49537	.54348	.69508
9	.23372	.26765	.33467	.39891	.45910	.51460	.56517	.72149
10	.23544	.27120	.34202	.40993	.47339	.53166	.58445	.74470
11	.23705	.27457	.34909	.42058	.48725	.54817	.60302	.76635
12	.23870	.27803	.35632	.43139	.50115	.56454	.62122	.78649
13	.24037	.28149	.36348	.44198	.51462	.58023	.63845	.80471
14	.24191	.28472	.37016	.45185	.52715	.59475	.65432	.82099
15	.24335	.28774	.37646	.46120	.53901	.60847	.66923	.83583
16	.24475	.29069	.38265	.47036	.55058	.62176	.68356	.84951
17	.24616	.29365	.38882	.47944	.56194	.63467	.69734	.86208
18	.24755	.29658	.39488	.48828	.57292	.64705	.71042	.87353
19	.24888	.29938	.40068	.49675	.58339	.65879	.72275	.88395
20	.25015	.30207	.40630	.50493	.59347	.67004	.73447	.89349
21	.25140	.30472	.41182	.51295	.60331	.68091	.74570	.90226
22	.25264	.30735	.41729	.52085	.61291	.69144	.75647	.91030
23	.25387	.30996	.42268	.52857	.62222	.70156	.76673	.91767
24	.25506	.31248	.42790	.53605	.63120	.71126	.77648	.92440
25	.25622	.31494	.43300	.54333	.63990	.72059	.78579	.93058
26	.25735	.31736	.43802	.55047	.64838	.72962	.79471	.93625
27	.25848	.31977	.44299	.55750	.65666	.73835	.80325	.94146
28	.25960	.32215	.44789	.56438	.66471	.74677	.81142	.94623
29	.26069	.32447	.45267	.57108	.67251	.75487	.81921	.95061
30	.26176	.32675	.45736	.57764	.68009	.76268	.82666	.95463
31	.26281	.32900	.46199	.58407	.68748	.77024	.83380	.95832
32	.26386	.33124	.46657	.59040	.69470	.77755	.84065	.96171
33	.26490	.33345	.47109	.59661	.70173	.78461	.84720	.96482
34	.26591	.33562	.47552	.60268	.70856	.79142	.85346	.96767
35	.26691	.33776	.47988	.60862	.71521	.79800	.85945	.97029
36	.26789	.33987	.48419	.61446	.72169	.80437	.86520	.97270
37	.26887	.34197	.48845	.62021	.72803	.81053	.87070	.97491
38	.26984	.34405	.49265	.62586	.73421	.81649	.87598	.97695
39	.27080	.34610	.49680	.63139	.74022	.82225	.88103	.97881
40	.27174	.34812	.50087	.63681	.74608	.82781	.88587	.98053

TABLE V (continued)

 $P_{(a)}^{(n)}\{\text{CS}|\theta^*, \text{LF}\}$ as a function of n for $a = 5$ and selected θ^*

n	θ^*							
	1.1	1.2	1.4	1.6	1.8	2.0	2.2	3.0
41	.27267	.35012	.50491	.64215	.75180	.83320	.89051	.98210
42	.27360	.35211	.50890	.64740	.75739	.83841	.89496	.98355
43	.27452	.35409	.51284	.65256	.76285	.84346	.89922	.98488
44	.27542	.35603	.51673	.65763	.76816	.84834	.90331	.98610
45	.27632	.35795	.52057	.66260	.77335	.85306	.90722	.98722
46	.27720	.35986	.52436	.66749	.77841	.85763	.91098	.98825
47	.27808	.36175	.52812	.67231	.78336	.86205	.91458	.98920
48	.27896	.36364	.53184	.67705	.78819	.86634	.91804	.99007
49	.27982	.36549	.53551	.68170	.79291	.87048	.92135	.99087
50	.28068	.36733	.53913	.68628	.79751	.87449	.92452	.99161
51	.28153	.36916	.54271	.69078	.80200	.87838	.92757	.99229
52	.28237	.37097	.54627	.69521	.80639	.88214	.93049	.99291
53	.28321	.37278	.54978	.69958	.81068	.88579	.93329	.99348
54	.28403	.37456	.55326	.70387	.81487	.88931	.93598	.99401
55	.28486	.37632	.55669	.70808	.81896	.89273	.93856	.99449
56	.28567	.37808	.56009	.71224	.82296	.89604	.94103	.99493
57	.28648	.37982	.56346	.71633	.82686	.89924	.94340	.99534
58	.28728	.38155	.56679	.72035	.83068	.90235	.94568	.99571
59	.28808	.38327	.57009	.72431	.83441	.90535	.94786	.99606
60	.28887	.38497	.57335	.72821	.83805	.90826	.94996	.99638
61	.28966	.38666	.57658	.73205	.84160	.91109	.95197	.99667
62	.29044	.38834	.57978	.73583	.84508	.91382	.95390	.99694
63	.29121	.39001	.58295	.73955	.84848	.91647	.95575	.99718
64	.29198	.39167	.58609	.74322	.85180	.91903	.95752	.99741
65	.29275	.39332	.58920	.74682	.85505	.92152	.95923	.99762
66	.29351	.39495	.59227	.75038	.85822	.92392	.96086	.99781
67	.29426	.39658	.59532	.75388	.86132	.92626	.96243	.99798
68	.29501	.39819	.59834	.75732	.86436	.92852	.96393	.99815
69	.29576	.39980	.60134	.76072	.86732	.93071	.96538	.99829
70	.29650	.40139	.60430	.76406	.87022	.93283	.96677	.99843
72	.29797	.40455	.61014	.77060	.87582	.93688	.96937	.99867
74	.29942	.40767	.61588	.77694	.88117	.94068	.97177	.99888
76	.30085	.41075	.62151	.78310	.88629	.94425	.97399	.99905
78	.30227	.41380	.62705	.78908	.89118	.94761	.97602	.99920
80	.30367	.41681	.63248	.79488	.89585	.95075	.97790	.99932
82	.30506	.41979	.63783	.80052	.90033	.95371	.97963	.99942
84	.30644	.42274	.64308	.80599	.90460	.95649	.98123	.99951
86	.30780	.42566	.64823	.81130	.90869	.95911	.98269	.99959
88	.30915	.42855	.65331	.81647	.91260	.96156	.98405	.99965
90	.31048	.43141	.65829	.82148	.91634	.96386	.98529	.99970

TABLE V (continued)

 $P_{\{a\}}^{(n)}\{\text{CS}|\theta^*, \text{LF}\}$ as a function of n for $a = 5$ and selected θ^*

n	θ^*							
	1.1	1.2	1.4	1.6	1.8	2.0	2.2	3.0
92	.31180	.43424	.66320	.82635	.91992	.96603	.98644	.99975
94	.31311	.43704	.66802	.83109	.92335	.96807	.98750	.99979
96	.31441	.43981	.67276	.83568	.92662	.96998	.98848	.99982
98	.31570	.44257	.67743	.84015	.92975	.97178	.98938	.99985
100	.31698	.44529	.68201	.84450	.93275	.97346	.99021	.99987
102	.31824	.44799	.68653	.84872	.93562	.97505	.99097	.99989
104	.31950	.45066	.69097	.85282	.93837	.97654	.99167	.99991
106	.32074	.45331	.69534	.85681	.94099	.97795	.99232	.99992
108	.32198	.45594	.69964	.86069	.94350	.97926	.99292	.99993
110	.32320	.45854	.70388	.86446	.94591	.98050	.99347	.99994
112	.32442	.46113	.70805	.86812	.94821	.98167	.99398	.99995
114	.32563	.46369	.71215	.87168	.95041	.98276	.99445	.99996
116	.32683	.46623	.71619	.87515	.95252	.98379	.99488	.99997
118	.32802	.46875	.72016	.87852	.95454	.98476	.99528	.99997
120	.32920	.47125	.72408	.88179	.95647	.98567	.99565	.99998
122	.33037	.47372	.72793	.88497	.95832	.98652	.99599	.99998
124	.33154	.47619	.73173	.88807	.96009	.98733	.99630	.99998
126	.33269	.47862	.73547	.89108	.96178	.98808	.99659	.99998
128	.33385	.48105	.73915	.89401	.96341	.98879	.99685	.99999
130	.33499	.48345	.74278	.89686	.96496	.98946	.99710	.99999
132	.33612	.48583	.74635	.89963	.96644	.99009	.99732	.99999
134	.33725	.48820	.74986	.90233	.96787	.99068	.99753	.99999
136	.33837	.49055	.75333	.90495	.96923	.99123	.99772	.99999
138	.33949	.49288	.75674	.90750	.97053	.99176	.99790	.99999
140	.34059	.49520	.76011	.90998	.97178	.99225	.99806	.99999
144	.34279	.49978	.76668	.91474	.97412	.99314	.99835	.99999+
148	.34496	.50430	.77307	.91924	.97626	.99393	.99860	
152	.34711	.50876	.77927	.92351	.97823	.99463	.99881	
156	.34923	.51315	.78528	.92754	.98003	.99525	.99898	
160	.35133	.51749	.79113	.93136	.98169	.99580	.99914	
164	.35340	.52177	.79681	.93498	.98320	.99628	.99926	
168	.35546	.52600	.80232	.93840	.98459	.99671	.99937	
172	.35750	.53017	.80768	.94164	.98586	.99709	.99947	
176	.35951	.53429	.81289	.94471	.98703	.99743	.99955	
180	.36151	.53836	.81795	.94762	.98810	.99772	.99961	
184	.36349	.54237	.82286	.95037	.98908	.99798	.99967	
188	.36545	.54634	.82764	.95297	.98999	.99822	.99972	
192	.36739	.55027	.83229	.95544	.99081	.99842	.99976	
196	.36932	.55414	.83680	.95778	.99157	.99860	.99980	
200	.37123	.55798	.84119	.95999	.99227	.99876	.99983	

Theorem 6.1: Under multiplicativity, and for the symmetric case ($a = b$, $\theta_\alpha^* = \theta_\beta^* = \theta^*$) in the LF-configuration, the same number of matrix-observations is necessary to guarantee

$$P_{(a,a)}^{(n)}\{\text{CS} | (\theta^*, \theta^*), \text{LF}\} \geq P^* \quad (6.1)$$

as the number of vector-observations necessary to guarantee

$$P_a^{(n)}\{\text{CS} | \theta^*, \text{LF}\} \geq \sqrt{P^*}. \quad (6.2)$$

Proof of theorem: We know from Theorem 4.1 that

$$\begin{aligned} P_{(a,b)}^{(n)}\{\text{CS} | (\theta_\alpha^*, \theta_\beta^*), \text{LF}\} = \\ P_a^{(n)}\{\text{CS} | \theta_\alpha^*, \text{LF}\} \times P_b^{(n)}\{\text{CS} | \theta_\beta^*, \text{LF}\}. \end{aligned} \quad (6.3)$$

Thus, in the symmetric case ($a = b$, $\theta_\alpha^* = \theta_\beta^* = \theta^*$) we have

$$P_{(a,b)}^{(n)}\{\text{CS} | (\theta^*, \theta^*), \text{LF}\} = [P_a^{(n)}\{\text{CS} | \theta^*, \text{LF}\}]^2$$

We seek the smallest number (n) of matrix-observations such that

$$P_{(a,a)}^{(n)}\{\text{CS} | (\theta^*, \theta^*), \text{LF}\} \geq P^*, \quad (6.4)$$

i.e., such that

$$P_a^{(n)}\{\text{CS} | \theta^*, \text{LF}\} \geq \sqrt{P^*}. \quad \text{QED} \quad (6.5)$$

Hence from (6.3) we see that $2n$ vector-observations (n for each of the two factors) are necessary to guarantee P^* whereas only n matrix-observations are necessary to guarantee P^* . More generally, for an f -factor ($f > 2$) multinomial experiment with multiplicativity and complete symmetry, fn vector-observations (n for each of the f factors) are necessary to guarantee P^* whereas only n "matrix"-observations are necessary to guarantee P^* .

Remark 6.1: In the non-symmetric case ($a \neq b$ and/or $\theta_\alpha^* \neq \theta_\beta^*$) the problem is more complicated. A 2-factor experiment with n matrix-observations and marginal probabilities in the least-favorable configurations has probabilities of a correct selection for the two factors given by

$$P_1 = P_{(a)}^{(n)} \{CS | \theta_\alpha^*, LF\}$$

and

$$P_2 = P_{(b)}^{(n)} \{CS | \theta_\beta^*, LF\}.$$

The optimal choice of sample size is the minimum value of n for which $P_1 \cdot P_2 \geq P^*$. This is illustrated in the computations given in Table VI.

If a separate single-factor experiment is used for each factor, different sample sizes can be employed in each case. With sample sizes n_1 and n_2 we have

$$P_1 = P_{(a)}^{(n_1)} \{CS | \theta_\alpha^*, LF\}$$

and

$$P_2 = P_{(b)}^{(n_2)} \{CS | \theta_\beta^*, LF\},$$

and the optimal sample sizes are given by the pair (n_1, n_2) which minimizes $n_1 + n_2$ subject to $P_1 \cdot P_2 \geq P^*$.

It is easily shown that $n \leq n_1 + n_2 \leq 2n$ where n is the minimum number of matrix observations and $n_1 + n_2$ the minimum total number of vector observations needed to guarantee an overall $P\{CS\} \geq P^*$ in the least favorable configuration. Clearly the greatest savings arise in the symmetric case where $n_1 + n_2 = 2n$.

7. AN APPLICATION OF THE PROCEDURE

The following is an example of how the procedure might be applied in a real-life situation:

A market research organization is planning a phone survey for one of its clients. To increase efficiency, two characteristics of a product are to be surveyed simultaneously. Every person who is phoned will be asked to state which combination of levels of each characteristic is most preferred. Because

of the nature of the characteristics, it is believed that the responses to the two questions will be independent. It is desired to determine the number of phone calls to be made in order that the client will have sufficient assurance that a correct selection will be made for both characteristics, and that the probability of being correct simultaneously for both characteristics is at least a

prespecified P^* . Suppose that the formulation of Section 2 is adopted, and that there are 3 levels of the first characteristic and 4 of the second. Suppose further that $\theta_\alpha^* = 1.4$ and $\theta_\beta^* = 2.0$ are specified. If $P^* = 0.90$ is specified then we see from Table VI that approximately $n = 128$ phone calls will guarantee (2.1). (Note that $P^* = 0.95$ will require approximately $n = 186$ phone calls, and it is clear that the number of required phone calls will become very large for θ_α^* and/or $\theta_\beta^* \rightarrow 1$ and/or $P^* \rightarrow 1$.)

A less efficient way of conducting the survey in the independence case is to ask each person one question. Then the overall problem can be treated as two separate problems (each as in B-E-M (1959)), one with $(a, \theta_\alpha^*, P_1^*)$ and the other with $(b, \theta_\beta^*, P_2^*)$ where $P_1^* \times P_2^* = P^* = 0.9$. Let n_1 and n_2 , respectively, denote the number of phone calls required for each problem. Of interest is $n_1 + n_2$ which is to be compared with $n = 128$. For convenience

one might take $P_1^* = P_2^* = \sqrt{P^*}$; in this case for which $\sqrt{P^*} = \sqrt{0.90} = 0.94868$, it is necessary to phone $n_1 = 183$ people to find their preferences for the first characteristic and a further $n_2 = 60$ people to find their preferences for the second characteristic. This total of 243 responses can be reduced by a more judicious choice of (P_1^*, P_2^*) subject to their satisfying $P_1^* \times P_2^* = P^*$.

By enumeration it can be shown that the optimal pair $(P_1^*, P_2^*) = (0.9239, 0.9742)$ to minimize $n_1 + n_2$ yields $n_1 = 148$ and $n_2 = 78$. (Several

neighboring pairs with $n_1 + n_2 = 226$ also guarantee $P^* = 0.90$ although their achieved $P\{\text{CS}\}$ is lower than that associated with the optimal pair.) Comparison of this total number of 226 phone calls with the original number of 128 illustrates the advantage of asking each respondent both questions.

TABLE VI

Table of $P_{(a)}^{(n)}\{\text{CS}|\theta^*, \text{LF}\}$ for
 $(a, \theta^*) = (3, 1.4)$ and $(4, 2.0)$
and selected n

n	$P_{(3)}^{(n)}\{\text{CS} 1.4, \text{LF}\}^\dagger$	$P_{(4)}^{(n)}\{\text{CS} 2.0, \text{LF}\}^{\dagger\dagger}$	$P_{(3)}^{(n)}\{\text{CS} 1.4, \text{LF}\} \times P_{(4)}^{(n)}\{\text{CS} 2.0, \text{LF}\}$
126	0.90160	0.99580	0.89781
128	0.90388	0.99610	0.90035
130	0.90615	0.99638	0.90287
184	0.94977	0.99951	0.94930
186	0.95089	0.99955	0.95046
188	0.95199	0.99958	0.95159

† Probabilities obtained from TABLE III.

†† Probabilities obtained from TABLE IV.

8. EXPERIMENTATION WHEN MULTIPLICATIVITY DOES NOT HOLD

If the assumption of independence is not plausible, then the procedure described in the present paper is not applicable since now interaction is present. However, the problem can be regarded as that of sampling from a k -level ($k = a \cdot b$) multinomial population with probability vector $(p_{11}, \dots, p_{1b}, p_{21}, \dots, p_{2b}, \dots, p_{a1}, \dots, p_{ab}) = (\gamma_1, \gamma_2, \dots, \gamma_k)$ (say) with the ordered probabilities given by $\gamma_{[1]} \leq \dots \leq \gamma_{[k]}$. Then the problem reduces to the one considered in B-E-M (1959) with the following.

GOAL: To select the event associated with $\gamma_{[k]}$. (8.1)

A correct selection for goal (8.1) is equivalent to a correct selection for the goal of Section 2 under independence but need not be equivalent if dependence is present. This can be seen by the following example.

Example 8.1: Suppose that $\underline{\alpha} = (3/5, 2/5)$ and $\underline{\beta} = (4/7, 3/7)$ are the marginal success probabilities. Then under independence we have $p_{11} = 12/35$, $p_{12} = 9/35$, $p_{21} = 8/35$, $p_{22} = 6/35$, and the event associated with (α_1, β_1) has the largest probability. However, with $\underline{\alpha}, \underline{\beta}$ as above but with dependence we might have $p_{11} = 9/35$, $p_{12} = 12/35$, $p_{21} = 11/35$, $p_{22} = 3/35$. Then the event associated with (α_1, β_2) has the largest probability. The experimenter thus must be aware of the possibility that the event associated with the largest probability can change if he adopts (8.1) when dependence is present.

If the experimenter is prepared to adopt the goal (8.1) in the dependence situation and also adopts the associated probability requirement

$$P\{CS\} \geq P^* \text{ whenever } \gamma_{[k]} \geq \theta^* \gamma_{[k-1]}, \quad (8.2)$$

then tables such as those given in G—O—S (1977), pp. 445–497, are applicable. If the experimenter adopts (8.1) and (8.2) under independence, then $\gamma_{[k]} = \alpha_{[a]} \beta_{[b]}$ and $\gamma_{[k-1]} = \max\{\alpha_{[a]} \beta_{[b-1]}, \alpha_{[a-1]} \beta_{[b]}\}$. The resulting n -value required to guarantee (8.2) will usually be quite large since $k = a \cdot b$ is now large.

9. CONCLUDING REMARKS

We have demonstrated that single-stage factorial multinomial experiments in which observations are taken in matrices are more efficient than independent single-stage single-factor multinomial experiments in which observations are taken in vectors provided that the factor-level responses of each category are independent of the factor-level responses of all other categories. If this independence condition holds then the number of matrix-observations to guarantee P^* will be less (often substantially so) than the total number of vector-observations to guarantee that same P^* when both methods of experimentation seek to select the event associated with the best factor-level combination; the savings are greatest under symmetry (i.e., $(a = b, \theta_{\alpha}^* = \theta_{\beta}^*)$ for 2-factor experiments). However, if independence does not exist, then reducing the selection problem to one involving a $(k = a \cdot b)$ -level single-factor multinomial with a goal and probability

requirement as in B-E-M (1959) may result in a different event being associated with the best factor-level combination, and a much larger number of vector-observations being required.

Even if independence and symmetry are present, the number of matrix-observations necessary to guarantee P^* if single-stage experimentation is used will be very large for θ^* and/or P^* close to unity. However, if sequential experimentation is feasible for the application being studied, it may be possible to guarantee (2.1) with a smaller expected number of matrix-observations than the corresponding single-stage sample size n by truncating the generalized B-K-S open sequential procedure described in Section 4.4 of B-G (1988a). This was done successfully for the normal means selection procedure in B-G (1988b). We plan to undertake such a study for the multinomial.

ACKNOWLEDGMENTS

This research was partially supported by the U.S. Army Research Office through the Mathematical Sciences Institute of Cornell University and by U.S. Army Research Office Contract DAAL03-86-K-0046 at Cornell University. The writers are indebted to Professor Santner for his constructive comments. We also wish to thank Ms. Kathy King for her expertise in typing this manuscript.

BIBLIOGRAPHY

- Alam, K. (1971). On selecting the most probable category. Technometrics 13, 843-850.
- Bechhofer, R.E., Elmaghraby, S. and Morse, N. (1959). A single-sample multiple decision procedure for selecting the multinomial event which has the highest probability. Ann. Math. Statist. 30, 102-119.
- Bechhofer, R.E. and Goldsman, D.M. (1985a). On the Ramey-Alam sequential procedure for selecting the multinomial event which has the largest probability. Comm. Statist.—Simula. Computa. B14(2), 263-282.
- Bechhofer, R.E. and Goldsman, D.M. (1985b). Truncation of the Bechhofer-Kiefer-Sobel sequential procedure for selecting the multinomial event which has the largest probability. Comm. Statist.—Simula. Computa., B14(2), 283-315.