



Rhubarb Curculio

Lixus concavus Say; Family: Curculionidae

Table of Contents

- [Injury](#)
- [Description](#)
- [Life History](#)
- [Management](#)



Adult rhubarb curculio, with yellowish coating present.
Photos by M. Malkin.



Darker-colored rhubarb curculio. Size: about 1/2 inch long.

Injury

Rhubarb is a hardy perennial that is not usually seriously affected by insect pests. However, attacks of the rhubarb curculio occasionally do occur. The leaf stalks of the rhubarb may show exuding sap and partial decay from late-May through early summer, due to the feeding and egg laying punctures of the rhubarb curculio. Feeding injury appears as notches in the stem and on the leaf edges. Sap exudes from wounds of either type and collects as glistening drops of gum when fresh. Fortunately, the eggs of this insect do not hatch when deposited in rhubarb.

Description

The rhubarb curculio (or rhubarb weevil) is a large snout beetle, about 1/2 inch long. It is dark colored, with a yellow powdery material dusted on its back. The yellowish covering easily rubs off when the insect is handled. The head has a downwardly curved snout, at the end of which are the mandibles (the chewing mouth parts). The eggs are oblong and yellow- white in color. The mature larva is a legless grub about 3/4 inch in length, with a brown head.

Life History

The curculio overwinters as an adult, in piles of debris or in other protected places near the rhubarb planting. In about mid-May the adults appear, and are seen resting on the stalks and leaves of rhubarb, dock, thistle or sunflower. They soon begin laying eggs. Eggs are deposited singly in cavities about 1/8 inch deep in the stalks of host plants, and hatching occurs in a week to ten days, in all plants but the rhubarb. The rhubarb curculio survives in weeds in or near the garden. Eggs deposited in rhubarb do not hatch, but are killed by the actively growing plant tissue, which crushes them. In other hosts the newly hatched larva begins burrowing its way down through the stalk, so that when it reaches maturity in eight to nine weeks, it has reached the bottom of

