

**Establishing A Baseline Fetal Ovine Plasma ACTH Value Does Not Require
Frequent Sampling: Absence Of Significant Peaks Of Fetal ACTH
Concentration In Plasma In Late Gestation**

by

Sadowsky, D.W., McCulloch, C.E. and Nathanielsz, P.W.

BU-1222-M

September 1993

1 ESTABLISHING A BASELINE FETAL OVINE PLASMA ACTH VALUE
2 DOES NOT REQUIRE FREQUENT SAMPLING: ABSENCE OF
3 SIGNIFICANT PEAKS OF FETAL ACTH CONCENTRATION IN PLASMA
4 IN LATE GESTATION

5 Drew W. Sadowsky, Charles E. McCulloch*, and Peter W. Nathanielsz

6 Dept. Physiology, College of Veterinary Medicine and *Biometrics Unit, Dept. Plant
7 Breeding and Biometry, Cornell University, Ithaca, NY

8 Please address all correspondence to: Peter W. Nathanielsz, MD, PhD, ScD
9 Laboratory for Pregnancy and Newborn
10 Research
11 College of Veterinary Medicine
12 Dept. Physiology
13 Cornell University
14 Ithaca, NY 14853-6401

1 CONDENSATION

2 Baseline fetal ovine plasma ACTH varies during the 24h day but this variability does
3 not change with gestational age and is not composed of measurable discrete pulses.

4 Key Words: fetus, sheep, adrenocorticotropin, pulsatility

1 **ABSTRACT**

2 **Objective:** Fetal plasma ACTH may show short-term temporal variability. Our objective
3 was to determine whether one fetal ovine plasma sample provides adequate information to
4 establish a baseline fetal plasma ACTH concentration. If true, the ability to establish
5 baselines with a single sample would greatly reduce the amount of plasma removed from
6 a fetus to determine the baseline for an experiment.

7 **Study design:** We sampled plasma from four non-stressed fetuses at four different
8 sampling rates. Four different gestational age windows were used to investigate age-related
9 changes. Ewes were instrumented at 118 days gestational age (dGA). Fetuses were
10 sampled at 125-129, 130-134, 135-139, 140-146 dGA. Twelve consecutive fetal carotid
11 samples were withdrawn at each age at four sampling rates; every two, five, ten, and thirty
12 minutes.

13 Mean, standard deviation, and coefficient of variation (CV) were determined for each
14 12 sample series. Gestational age increases in mean plasma ACTH were analyzed by linear
15 regression of the four age groups. The CV's of the sampling rates, combined from all age
16 groups, were compared by two way ANOVA, with the individual sheep as the second factor.
17 Bonferroni's test of multiple comparisons was performed where necessary. The CV's of the
18 four gestational age windows, combined from all sampling rates, were similarly compared.
19 The presence of pulses were examined using the PULSAR program.

20 **Results:** Fetal blood gases were within normal limits for all time points. The mean
21 regression equation was $ACTH = 1.55 (dGA) - 178 \text{ pg.ml}^{-1}$, and was significant at the $p =$
22 0.05 level. The CV's were as follows: 125 dGA (48.8,25.1,8.1,8.5); 130 dGA (16.4,8.6,13.8,
23 34.1); 135 dGA (13.0,16.3,18.6,17.9); and 140 dGA (10.3,9.4,23.6, and 25.7); for the 2, 5, 10,
24 and 30 min sampling rates, respectively. None of the comparisons were significant. The few
25 pulses of ACTH detected were primarily detected in the 30 min sampling rate series.

26 **Conclusions:** These results support previous studies that show that fetal plasma ACTH
27 does rise gradually in late gestation. Baseline fetal ovine plasma ACTH varies over time,
28 but this variability does not change with gestational age and is not composed of measurable
29 discrete pulses of ACTH.

1 **BACKGROUND**

2 Fetal ovine corticotropin (ACTH) may be released in a pulsatile manner. Some
3 evidence exists which suggests that the adult pituitary secretes ACTH as discrete pulses (1).
4 Plasma ACTH concentrations have been shown to vary over a one hour period in late
5 gestation fetal sheep (2). Fetal plasma cortisol levels also show considerable variation over
6 a 90 minute period (3). These cortisol fluctuations in undisturbed sheep may reflect an
7 underlying ACTH variability.

8 Norman *et al.* (1985) demonstrated that basal ACTH levels in the fetal sheep vary
9 with gestational age, and that this basal level increases at a rate of 1 pg.ml⁻¹.day⁻¹ over the
10 last three weeks of gestation (2). We have described a similar rate of increase in the fetal
11 sheep (4). This slow and steady rise in fetal plasma ACTH could reflect a change in pulse
12 amplitude or frequency of ACTH release.

13 The existence of pulses of ACTH in fetal sheep plasma would complicate the
14 assessment of fetal plasma ACTH concentration under basal conditions made following a
15 single measurement of ACTH. The detection of basal ACTH pulses would be complicated
16 by the low magnitude of the reported variability (2). While the pituitary response to a
17 stressor may involve the release of a single, large, "pulse" of ACTH, the basal ACTH plasma
18 level would be more likely to be maintained by regularly occurring pulses of low amplitude.
19 Sampling plasma ACTH at a rate faster than the pulse frequency should more readily reveal
20 the pulses than would less frequent sampling. In addition, sampling at several different
21 frequencies should eliminate any variability inherent in the process of repeated measures
22 and avoid false pulse detection.

23 Various methods have been used to detect pulses using time-series hormone data (1,
24 5,6,7). The PULSAR program (7) uses a running mean of the hormone series being
25 investigated as the baseline, and the size of the template used to tag hormone peaks is
26 based on the variability of the radioimmunoassay used for the hormone being investigated.
27 This method was employed in the present study to detect pulses of fetal plasma
28 immunoreactive-ACTH under the four different sampling rates.

29 The four following hypotheses were investigated in this study: the fetal sheep in late

1 gestation secretes basal ACTH in a pulsatile manner; the variability of fetal plasma ACTH
2 increases with increasing gestational age, reflecting the greater maturity of the pituitary; the
3 variability remains sufficiently low that a single ACTH sample provides a useful measure
4 of the basal ACTH level; and the mean basal fetal plasma ACTH concentration increases
5 in late gestation.

6 MATERIALS AND METHODS

7 **Care of Animals:** Four Rambouillet and Dorset cross ewes of known gestational
8 age were used in these studies. Each ewe carried one fetus. Animals were housed in
9 metabolic cages and given free access to alfalfa cubes and water *ad libitum*. Lights were
10 on for 16 hours each day beginning 0600. New York State College of Veterinary Medicine
11 guidelines for the care and use of the sheep were followed. Experiments were conducted
12 under an approved Cornell Institutional Animal Care and Use Committee protocol.

13 Surgery was conducted under halothane general anesthesia at 118 ± 2 days
14 gestational age (dGA) (mean \pm SD) using techniques described in detail previously (8).
15 Ewes were instrumented with polyvinyl catheters placed in the jugular vein and carotid
16 artery and with multistrand bipolar stainless steel wires (AS632; Cooner Sales Co. Inc.,
17 Chatsworth, CA) sewn 5 mm apart into the myometrium of the pregnant horn to record the
18 myometrial electromyogram. Fetuses were instrumented with polyvinyl catheters inserted
19 into the jugular vein and carotid artery (8). The myometrial electromyogram was recorded
20 as previously described (8), and was used only to indicate established labor, at which time
21 the sampling protocol was terminated and the fetus allowed to be born.

22 **Blood Sampling:** Maternal and fetal blood samples were drawn into sterile ice-
23 water chilled syringes, centrifuged at 4°C and the plasma was immediately flash frozen in
24 liquid nitrogen. Fetal and maternal blood gases were determined on 0.5 ml of blood in an
25 ABL2 Acid Base Laboratory (Radiometer A/S, Copenhagen, Denmark) at 39°C. Fetal
26 blood was handled aseptically and the erythrocytes were returned at the end of each
27 sampling series.

1 **Sampling Regimen:** Fetuses were sampled in four age windows: 125-129, 130-
2 134, 135-139, and 140-146 dGA. Twelve consecutive fetal carotid blood samples (0.7 ml)
3 were withdrawn over 15 seconds at each of four sampling rates: one sample every two (2)
4 minutes, one every five (5) minutes, one every ten (10) minutes, and one every thirty (30)
5 minutes. At least two hours elapsed between sampling series. Two sets of sampling rates
6 were performed on one day, and the other two were performed the following day. The
7 order in which the four sampling protocols was carried out was randomized. Morning
8 sampling began between 0900 and 1100h and afternoon sampling began between 1400 and
9 1700h.

10 **Immunoreactive Corticotropin (ACTH) Radioimmunoassay:** Plasma ACTH
11 was measured by radioimmunoassay in unextracted plasma using a miniaturized version of
12 the INC Star ACTH kit (Stillwater, MN, USA) validated specifically for use in fetal sheep
13 plasma and previously described in detail (9). Plasma was thawed on ice and 50 μ l plasma
14 was incubated with rabbit anti-human ACTH(1-24) antiserum for 24 hours at 4 ° C, and then
15 incubated with iodinated ACTH(1-24) for another 24 hours at 4 ° C. The bound fraction was
16 precipitated with a goat anti-rabbit second antibody (30 minutes at room temperature, 20-
17 22° C).

18 The standard curve was generated from human plasma standards of concentrations
19 5, 10, 20, 50, and 100 pg.ml^{-1} . Interassay coefficient of variation (CV) for a fetal sheep
20 plasma pool (mean concentration 28.8 pg.ml^{-1} , n=9) was 12.5%, and for maternal pool
21 (mean concentration 42.8 pg.ml^{-1} , n=8) was 8.2%. Intraassay CV were 6.1 and 3.0% respec-
22 tively. The assay sensitivity, calculated as the 90% bound fraction over the free fraction
23 (B/B0) limit of detection, was 5.0 pg.ml^{-1} of sample.

24 **PULSAR Program Parameters:** One ACTH assay was devoted to various fetal
25 and maternal ovine plasma pools which covered the range of the assay. Each pool was run
26 in a series of ten samples and used to generate an assay variability curve. This curve
27 determined the "Noise Units" portion of the PULSAR program; the determination of how
28 much variability would be due to the assay variability, rather than to a physiological ACTH
29 pulse.

1 The PULSAR program used amplitude and duration as the criteria for peak
2 determination. A smaller amplitude was required for a pulse containing two points than for
3 a peak containing only one point. This diminishing amplitude-criterion was continued out
4 to five points. The amplitude of each of these amplitude-criteria was determined empiri-
5 cally, using two data sets. The first data set consisted of the series of ten replicates of the
6 pools used above, and was used as a negative control where no pulses should be detected.
7 The first pass through each series had the ten replicates in the original assay order. The
8 second pass through each series had the individual replicates ordered in such a way as to
9 simulate one pulse. The second data set served as the positive controls and consisted of
10 samples from three fetuses used in another experimental protocol. Twelve fetal plasma
11 samples were obtained at a sampling rate of five minutes, and were taken at several
12 different gestational ages from each sheep.

13 Up to six iterations were allowed in the determination of the running baseline. The
14 iteration method weights points composing a pulse so as to minimize their effect on the
15 determination of the trend line composing the baseline. A pulse was split into two when
16 the intervening points fell by one standard deviation unit (SDU) below that of the peak
17 concentration, even though the intervening point may not reach the baseline concentration.

18 **Statistics:** Gestational age increases in mean plasma ACTH were analyzed by
19 linear regression. The mean ACTH values for each 12-sample series, fifty-one points in
20 total, was used to generate the mean regression equation. Individual linear regression for
21 each sheep were also conducted, using the mean ACTH value from each sampling rate
22 versus the actual dGA.

23 The coefficient of determination, R^2 , was used as the measure of the variability in
24 mean plasma ACTH due to the gestational age. Analysis of variance was conducted on the
25 regression results, where the F-test ($\alpha=0.05$) was used to determine whether the calculated
26 slope was significantly different from zero.

27 The CV from each sampling rate within an age group was compared by two-way
28 analysis of variance using the individual sheep as the second factor. Significance at the
29 $p<0.05$ level was then further analyzed by Bonferroni's test of multiple comparisons. The

1 analysis was conducted using all sampling rates combined. The CV's from each sampling
2 rate were similarly compared across the combined gestational age groups.

3 The number of pulses per twelve sample series was analyzed similarly to that of the
4 coefficients of variation. Two-way analysis of variation was conducted using both the
5 gestational age and individual sheep analysis, and the sampling rate and individual sheep
6 analysis. Data throughout are provided as mean \pm S.D. and n = 4 unless otherwise noted.

7 RESULTS

8 **Fetal Condition:** Fetal blood gases were within normal limits for all time points.
9 Fetal PO₂ was 18.8 \pm 1.2 mm Hg at the start of the experiment, and had not changed at the
10 time of its conclusion.

11 The twelve-sample series of samples did not show a rise in trend in the ACTH levels
12 from beginning to end of the series. This observation indicates that the sampling procedure
13 did not disturb the fetus significantly. All ewes underwent spontaneous delivery. All fetuses
14 were born alive and in good condition.

15 **Linear Regressions:** The mean linear regression equation was ACTH = 1.55
16 (dGA) -178, p < 0.05. This indicated an increase of 1.55 pg.ml⁻¹ in fetal plasma ACTH per
17 increase of one day of gestational age during the period studied, 125-146 days gestation.

18 Table 1 gives the results of the individual regression equations. The variables pre-
19 sented are the slope, coefficient of determination R² (adjusted for the degrees of freedom),
20 and the F-statistic. All individual fetal ACTH regressions were significant and fit a linear
21 model. However, fetus 3006 demonstrated an increase in the slope between 130 and 135
22 dGA windows, suggesting that a more complex, nonlinear, model may be more appropriate
23 for this fetus.

24 **Coefficients of Variation of Fetal Plasma ACTH:** Table 2 gives the mean
25 plasma ACTH values of each time series for each animal. These values were used to
26 generate the mean regression equation. As seen here, the mean fetal plasma ACTH values
27 obtained from the four different sampling rates for one sheep over one gestational age
28 window were not always identical. Figure 1 gives one representative example of the twelve

1 fetal plasma ACTH samples obtained from fetus 3006 over the 125-129 dGA window at
2 each of the four sampling rates. This age window was selected to illustrate the basal
3 variability present in fetal plasma ACTH as no pulses were present.

4 Table 3 gives the coefficients of variation for each of the four dGA windows at each
5 of the four sampling rates. Missing values seen in Tables 2 and 3 arose from early problems
6 with catheter blockages and the presence of active labor. Due to the incomplete block
7 structure of the results, the interaction term by two-way analysis of variance could only be
8 determined for the sampling rate and individual fetal factors, but not for the gestational age
9 window and individual fetal factors. The interaction term was not significant at the $\alpha=0.05$
10 level. None of the analysis of variance tests for the sampling rate, the gestational age
11 window, and the individual fetal factors were significant.

12 **PULSAR pulse detection results:** For the determination of the "Noise Units"
13 in the PULSAR program, a binomial regression curve was fit to a fetal ovine ACTH
14 radioimmunoassay variability curve. To generate this curve, ACTH values of multiple
15 replicates of seven different plasma pools and their serial dilutions, covering the range of
16 9 to 120 pg.ml⁻¹, were quantitated by the RIA and plotted with the mean of the pools on
17 the abscissa and the standard deviation *s* of the pool replicates on the ordinate. The
18 binomial regression curve was expressed as:

$$19 \quad y = 1.21 + 0.0119x + 0.000818x^2.$$

20 The PULSAR program's amplitude-criteria, determined by using the two test data
21 sets, for the fetal plasma ACTH model were as follows: for a one-point pulse, the
22 amplitude-criterion was 2.8 assay standard deviation units (SDU) above the baseline ACTH
23 concentration; for a two-point pulse, 2.3 SDU; for a three-point pulse, 1.8 SDU; for a four-
24 point pulse, 1.5 SDU; and for a five-point pulse, 1.2 SDU. All of the components of the
25 model were now established.

26 Nineteen peaks were detected in fifty-one twelve-sample series representing 122.2
27 hours (Table 4). This gives a basal rate of 1 plasma ACTH pulse per 6.43 hours in a late-
28 gestation ovine fetus. Seven sample series had multiple pulses detected. Of the twenty
29 pulses detected, half were from the thirty-minute sampling rate time series. Half of the

1 pulses were between one and two standard deviations greater than the mean of the sample
2 series, which was calculated including the points composing the pulse.

3 The two-way analysis of variance of pulses detected was not significant in any
4 comparison. The interaction between the sampling rate term and the individual sheep term
5 was also not significant. The interaction between the gestational age window term and the
6 individual sheep term could not be determined due to the incomplete block design.

7 DISCUSSION

8 The lack of increase in plasma ACTH values from the beginning to the end of each
9 sampling series indicated that neither the sampling rate, nor the volume of blood removed,
10 was causing a stress-release of ACTH. Thus, the fetal plasma ACTH results were believed
11 to represent true basal secretion.

12 The rate of increase of 1.6 pg.ml^{-1} in fetal plasma ACTH per increase of one day of
13 gestational age is similar to the rate reported previously by others (2), as well as ourselves
14 (4), and supports the idea that ACTH provides the drive for the fetal plasma cortisol rise
15 which initiates parturition and organ maturation in this species.

16 The low coefficients of variation suggest that the variability in plasma ACTH
17 concentrations is not large enough to warrant the removal of multiple plasma samples from
18 the fetus for the determination of a single time-point value in baseline samples, at rest. The
19 variability does not appear to change with sample frequency or with dGA. However, diffe-
20 rent means for the same gestational age window were obtained with each series of samples;
21 the individual series being no more than three days apart. In contrast, fetal plasma cortisol
22 time series coefficients of variation have been reported to change with gestation (3). Challis
23 et al. (1981) reported that fetal plasma cortisol variability was larger (36%) on days 11 to
24 20 before labor initiation than the variability (15-19%) measured on days 21 to 30 and days
25 5 to 0 before labor (3). The current results would suggest that this observation is due to
26 maturation in the adrenal cortex itself, rather than changes in ACTH secretion.

27 The possibility remains for ACTH to be released in a pulsatile fashion in the fetal
28 sheep in response to a stressor. Of particular interest would be the effect of a chronic

1 stressor, where the repeatability of any ACTH pulses on some timed fashion could be mea-
2 sured.

3 The PULSAR results demonstrate that pulses do occur. However, the frequency was
4 very low suggesting that the pulses detected may have been in response to some change in
5 homeostasis rather than to maintain the basal levels. Reported ACTH half-lives of two to
6 seven minutes in sheep (10) and rats (11) suggest that more frequent pulses would be
7 required to maintain basal levels than the few demonstrated here. However, the assay used
8 may not be sensitive enough to enable PULSAR to detect small repeated pulses. A study
9 which would involve intracellular voltage recording from corticotropin cells in the
10 adenohypophysis might indicate the mechanism of maintenance of basal levels of ACTH
11 release in the ovine fetus, pulsatile or continuous. Such a study has been done in the
12 gonadotropin-releasing hormone pulse generator cells (12). However, the corticotropin cells
13 are more randomly dispersed in the adenohypophysis, which would make such an *in vivo*
14 experiment difficult to perform.

15 In a study measuring both cortisol and ACTH in fetal sheep every five minutes over
16 a period of two hours, pulses (as analyzed by frequency domain methods) of ACTH
17 occurred with a frequency of two per hour at both 133 and 143 dGA (13). The amplitude
18 of the pulses (34 pg.ml^{-1}) was not different at the two ages, while the mean concentration
19 of ACTH rose (13). Recently, Brooks and Challis (1991) have reported fetal ovine plasma
20 ACTH pulses occurring on the average of one every 40 minutes in fetuses sampled for six
21 hours at ten minute intervals at 140 dGA (14). They found much higher variability, but
22 similar means, in their samples (14) than those seen in the current study. Both the rate and
23 the amplitude of the pulses (13, 14) were within the range of detection of PULSAR and the
24 time series. The difference in results may be attributable to the different pulse detection
25 methods.

26 Myometrial contractures produced by the administration of oxytocin to the pregnant
27 ewe result in the release of ACTH by the fetus (15,16). This stimulus to fetal ACTH
28 secretion requires the fall in fetal PO_2 that generally accompanies a contracture (16). In
29 several studies the frequency of contractures during late pregnancy in sheep has been shown

1 to vary from one every 30 to 60 minutes (17,18). Our study made no attempt to specifically
2 sample fetal blood in relation to individual contractures.

3 Adult rats, sampled either for four hours at a rate of two minutes, or for six hours
4 at a rate of fifteen minutes (1), indicated a much more complex pulse pattern than has been
5 shown here. The spectral analysis performed indicated rhythms of plasma ACTH concen-
6 trations with periodicities between four and 220 minutes. This observation further supports
7 the idea that the pulses detected in the ovine fetus were not for the maintenance of basal
8 plasma ACTH levels.

9 CONCLUSION

10 The lack of significance in the age group comparisons suggests that during development of
11 the hypothalamo-pituitary-adrenal cortical axis there are no changes in pulse amplitude or
12 frequency of ACTH. These results do not support the idea that more than one sample
13 needs to be taken to obtain a valid measurement of the unstressed baseline fetal plasma
14 ACTH. However, responses to experimental stimulation may be pulsatile. ACTH
15 concentrations during fetal hypoxia or hypotension should be investigated in a similar
16 manner.

1 REFERENCES

- 2 1. Carnes M, Goodman BM, Lent SJ. High resolution spectral analysis of plasma
3 adrenocorticotropin reveals a multifactorial frequency structure. *Endo* 1991;128:902-
4 10.
- 5 2. Norman LJ, Lye SJ, Wlodek ME, Challis JRG. Changes in pituitary responses to
6 synthetic ovine corticotrophin releasing factor in fetal sheep. *Can J Physiol*
7 *Pharmacol* 1985;63:1398-403.
- 8 3. Challis JRG, Patrick JE, Cross J, Wolkewych J, Manchester E, and Power S. Short-
9 term fluctuations in the concentration of cortisol and progesterone in fetal plasma,
10 maternal plasma, and amniotic fluids from sheep during late pregnancy. *Can J*
11 *Physiol Pharmacol* 1981;59:261-67.
- 12 4. McDonald TJ, Nathanielsz PW. Bilateral destruction of the fetal
13 paraventricular nuclei prolongs gestation in sheep. *Am J Obstet Gynecol*
14 1991;165:764-70.
- 15 5. Merriam GR, Wachter KW. Algorithms for the study of episodic hormone secretion.
16 *Am J Physiol* 1982;243:E310-18.
- 17 6. Clifton DK, Steiner RA. Cycle detection: a technique for estimating the frequency
18 and amplitude of episodic fluctuations in blood hormone and substrate
19 concentrations. *Endo* 1983;112:1057-64.
- 20 7. Veldhuis JD, Johnson ML. Cluster analysis: a simple, versatile, and robust algorithm
21 for endocrine pulse detection. *Am J Physiol* 1986;250:E486-93.
- 22 8. Nathanielsz PW, Bailey A, Poore ER, Thorburn GD, Harding R. The relationship
23 between myometrial activity and sleep state and breathing in fetal sheep throughout
24 the last third of gestation. *Am J Obstet Gynecol* 1980;138:653-59.
- 25 9. McDonald T, Reimers TR, Figueroa JP, Nathanielsz PW. The effect of continuous
26 intracerebroventricular administration of arginine vasopressin (AVP) to the fetal
27 lamb at 125 to 134 days of gestation on fetal plasma ACTH and cortisol concentra-
28 tions. *J Dev Physiol* 1989;12:35-40.
- 29 10. Jones CT, Luther E, Ritchie JWK, Worthington D. The clearance of ACTH from the
30 plasma of adult and fetal sheep. *Endo* 1975;96:231-34.

- 1 11. López FJ, Negro-Vilar A. Estimation of endogenous adrenocorticotropin half-life
2 using pulsatility patterns: a physiological approach to the evaluation of secretory
3 episodes. *Endo* 1988;123:740-46.
- 4 12. Kaufman J-M, Kesner JS, Wilson RC, Knobil E. Electrophysiological manifestation
5 of luteinizing hormone-releasing hormone pulse generator activity in the Rhesus
6 monkey: influence of α -adrenergic and dopaminergic blocking agents. *Endo*
7 1985;116:1327-33.
- 8 13. Apostolakis EM, Longo LD, Veldhuis JD, Yellon SM. Dissociation of pulsatile
9 cortisol and adrenocorticotropin secretion in fetal sheep during late gestation. *Endo*
10 1992;130:2571-78.
- 11 14. Brooks AN, Challis JRG. Effects of naloxone on the preparturient increase in
12 adrenocorticotrophin and cortisol in foetal sheep. *J Neuroendocrinology* 1991;3:419-
13 24.
- 14 15. Lye SJ, Wlodek ME, Challis JRG. Possible role of uterine contractions in the
15 short-term fluctuations of plasma ACTH concentration in fetal sheep. *J Endocrinol*
16 1985;106:R9-R11.
- 17
18 16. Woudstra BR, Kim C, Aarnoudse JG, Nathanielsz PW. Myometrial
19 contracture-related increases in plasma adrenocorticotropin in fetal sheep in the last
20 third of gestation are abolished by maintaining fetal normoxemia. *Endocrinology*
21 1991;129:1709-1713.
- 22
23 17. Harding R, Poore ER, Bailey A, Thorburn GD, Jansen CAM, Nathanielsz PW.
24 Electromyographic activity of the nonpregnant sheep uterus. *Am J Obstet Gynecol*
25 1982;142:448-457.
- 26
27 18. Bailey A., Harding R, Nathanielsz PW, Poore ER, Thorburn GD. Uterine activity
28 and its effect on the sheep fetus. *J Physiol* 1980;308:27P.

1 **Figure 1** Twelve-sample series of Fetal Plasma ACTH values from Fetus 3006 over the
2 125-129 dGA window at the four sampling rates.

1

Figure 1

1

Table 2 Mean plasma ACTH for Individual Time Series

	Fetus	dGA Window	2 minute	5 minute	10 minute	30 minute
2						
3	3006	125	20.43	19.98	19.15	18.30
		130	18.33	22.78	22.69	25.39
		135	46.52	47.56	56.65	54.04
		140	44.82	41.95	30.31	45.97
4	2107	125	--	--	--	--
		130	24.60	24.91	24.56	26.65
		135	30.74	33.46	34.29	30.25
		140	53.99	39.98	41.94	58.67
5	3001	125	--	--	--	--
		130	36.51	35.31	34.42	27.14
		135	50.36	46.22	40.21	46.17
		140	--	--	--	--
6	2057	125	22.77	12.68	16.31	20.37
		130	20.88	26.40	20.76	26.30
		135	15.03	10.37	23.55	19.47
		140	35.12	58.85	--	29.94

1

Table 3 Individual % Coefficients of Variation of ACTH

2	Sheep	dGA Window	2 minute	5 minute	10 minute	30 minute
3	3006	125	7.9	9.2	7.6	9.2
		130	14.4	4.4	9.3	67.9
		135	5.6	10.7	7.6	15.8
		140	8.1	9.5	21.1	28.7
4	2107	125	--	--	--	--
		130	9.8	4.3	21.0	40.9
		135	17.4	35.7	33.5	17.1
		140	7.3	12.0	26.1	28.5
5	3001	125	--	--	--	--
		130	35.2	16.4	18.8	15.5
		135	15.1	10.5	12.0	11.3
		140	--	--	--	--
6	2057	125	89.8	41.1	8.6	7.8
		130	6.3	9.2	6.1	12.2
		135	14.0	8.2	21.1	27.2
		140	15.4	6.8	--	19.8

1 Table 4 Description of 19 fetal plasma ACTH pulses organized by sampling rate:
 2 number of pulses per sample series, number of points composing a pulse, and
 3 pulse amplitude.

	2 min	5 min	10 min	30 min
4 1 pulse ^a	2	3	3	3
5 2 pulse ^b	0	0	1	3
6 2 points ^c	1	2	0	3
7 <1 SD ^d	0	1	0	1
8 >1 SD	1	1	2	6
9 >2 SD	0	0	3	1
10 >3 SD	1	1	0	1

11 ^aNumber of 12-point sample series at a given sampling rate which contained one pulse.

12 ^bNumber of 12-point sample series at a given sampling rate which contained two pulses.

13 ^cNumber of 12-point sample series at a given sampling rate which contained a pulse
 14 composed of two points (all other pulses are composed of only one point).

15 ^dPulse amplitude in units of standard deviation above mean of sample series (including
 16 pulse values).




Figure 5.1