

Culex pipiens Specimen Submission Kit

*** If you are using our full collection kit instead of a protocol that is already established by your agency please use [these guidelines](#) instead for collection and shipping instructions.**

<u>Item</u>	<u>Number included in kit</u>
Falcon Tubes (one to be used per egg raft)	10
White Ladles	1
Mesh Dippers (stick and screen mesh)	2
Whirl-Pak Bags	20
Ice Packs	1
Styrofoam Containers	1
Falcon tube fish food	1

[Larval Identification Key](#)

Specimen Collection Procedure

Culex pipiens larvae and egg rafts are often found in standing water and prefers polluted water with high levels of organic content. This species can be found in a range of containers that hold water, and is often associated with tire piles, catch basins, and storm drains. Another closely related species, *Cx. restuans*, is often found alongside *Cx. pipiens*. These species are difficult to differentiate as adults, but the larvae can be identified to species using the guide posted [here](#).

- 1) Only place a single egg raft into each falcon tube so that once the larvae emerge (~24 hours) each raft can be identified to species individually. Only 2 – 3 larvae per raft need to be examined and they should be chilled in ice water or placed in 70% ethanol prior to examination to immobilize them. If collecting more than one egg raft prepare another falcon tube and keep tubes containing egg rafts upright. The [Larval Identification Key](#) can be used to determine the species of each egg raft. The features used to differentiate 4th instar *Culex* larvae are also present in 1st instar larvae. If you are lacking equipment or training, skip this step and submit the specimens and they will be identified once they arrive.
- 2) Transfer the larvae to a **whirl-pak bag** filled 75% full of water and a small amount of **fish food** (~1/4 a pellet). Make sure there is a small air pocket enclosed in the bag to allow any hatched larvae to breath in transit. Tightly seal the whirl-pak enclose it within a second whirl-pak. Allow the egg rafts to sit at room temperature and hatch into larvae (usually 1 – 2 days).

- 3) Once the larvae have emerged place the bags into the **Styrofoam container** with **ice packs** separated by paper towels to prevent direct contact between the bags and ice packs. Fill out the [Specimen Submission Form](#) and send it to pesticide@cornell.edu to notify us of the impending delivery. Please also print out a copy and include in the package so that the label can be matched once it arrives at Cornell University.

(A)



(B)



Ship the culex (A) place ice packs in the bottom of the Styrofoam container and cover them with paper towels and (B) place whirl pak bags in container

Please send the package overnight* to the address below:

Harrington Lab
Cornell University
129 Garden Ave
3131 Comstock Hall
Ithaca, NY 14853

** If you require a fedex number to send the package overnight, one can be obtained by contacting pesticide@cornell.edu. Please note, we cannot support shipping for all agencies, so please only request a fedex number if necessary.*