

SCHOOL OF OPERATIONS RESEARCH
AND INDUSTRIAL ENGINEERING
COLLEGE OF ENGINEERING
CORNELL UNIVERSITY
ITHACA, NEW YORK 14853-3801

TECHNICAL REPORT NO. 1010

May 1992

**ON THE HEIGHT OF THE
MINIMAL HILBERT BASIS**

by

**J. Liu, L.E. Trotter, Jr.
and G.M. Ziegler**

This research was partially supported by NSF Grant DMS89-20550.

On the Height of the Minimal Hilbert Basis

JIYONG LIU, LESLIE E. TROTTER, JR. & GÜNTER M. ZIEGLER

Abstract. We present a simple geometric proof of a result due to Ewald and Wessels: in a pointed, polyhedral cone of dimension $n \geq 3$ with integer-valued generators, any linearly independent generator representation for a minimal Hilbert basis element has coefficient sum less than $n - 1$.

We consider here subsets of \mathbb{R}^n of the form $K = \{\sum_{i=1}^m \lambda_i a_i : \lambda_i \geq 0, 1 \leq i \leq m\}$, with $a_1, \dots, a_m \in \mathbb{Q}^n$, i.e., finitely generated, rational (*convex*) cones. We denote $K = \text{cone}\{a_1, \dots, a_m\}$. K is *nontrivial* if it properly contains $\{0\}$ and *pointed* if it contains no linear subspace properly containing $\{0\}$. When K is nontrivial and pointed, its *extreme rays* provide a unique (up to positive scaling) minimal set of generators.

When K is finitely generated, there also exists a finite set of vectors, a *Hilbert basis*, which generates $K \cap \mathbb{Z}^n$ under nonnegative integral combinations. For instance, when $K = \text{cone}\{a_1, \dots, a_m\}$, with a_1, \dots, a_m integral, the integral elements in the (half-open) *zonotope* $\{\sum_{i=1}^m \lambda_i a_i : 0 \leq \lambda_i < 1, 1 \leq i \leq m\}$ constitute a Hilbert basis for K . When K is nontrivial and pointed, then its integral elements have a unique *minimal* Hilbert basis consisting of those members of $K \cap \mathbb{Z}^n$ which are not sums of other members of $K \cap \mathbb{Z}^n$. Our interest here is in properties of minimal Hilbert bases and we thus assume henceforth that $K = \text{cone}\{a_1, \dots, a_m\}$ is nontrivial and pointed. We also assume, for $1 \leq i \leq m$, that a_i is the *shortest integral* vector on the extreme ray $\{\lambda a_i : \lambda \geq 0\}$, i.e., that $a_i \in \mathbb{Z}^n$ and $\gcd\{a_{ij} : 1 \leq j \leq n\} = 1$, where a_{ij} denotes the j -th component of the vector a_i . Finally, we assume with no loss in generality that K is *full-dimensional*; i.e., $\text{rank}\{a_1, \dots, a_m\} = n$. For background material on cones and Hilbert bases, the reader is referred to Schrijver [5]. Now let x be an element of the minimal Hilbert basis for K . By Carathéodory's Theorem, there are linearly independent generators a_{i_1}, \dots, a_{i_n} so that $x \in \text{cone}\{a_{i_1}, \dots, a_{i_n}\}$; i.e., x is in the *simplicial* subcone of K generated by a_{i_1}, \dots, a_{i_n} . Of course, x may be contained

in many such simplicial subcones, and we define the *height* of x as

$$h(x) = \max\left\{\sum_{i=1}^n \lambda_i : x = \lambda_1 a_{i_1} + \dots + \lambda_n a_{i_n}; \lambda_i \geq 0, 1 \leq i \leq n; \text{rank}\{a_{i_1}, \dots, a_{i_n}\} = n\right\}.$$

Motivated by a question of algebraic geometry, G. Ewald [1] posed the problem of determining an upper bound on $h(x)$. Subsequently, Ewald and Wessels [2] established that $h(x) \leq n - 1$ and demonstrated, moreover, that this bound is sharp. Here we indicate a simple geometric proof of this result, discovered independently and motivated by results in Liu [3]. We also see that equality $h(x) = n - 1$ cannot occur for $n > 2$.

Theorem. *Let K be an n -dimensional pointed polyhedral rational cone, $n \geq 3$, and let x be a minimal Hilbert basis element for K . Then $h(x) < n - 1$, and this bound is best-possible.*

Proof. Suppose $x \in C'$, where $C' = \text{cone}\{a_1, \dots, a_n\}$ and the generators a_1, \dots, a_n define a simplicial subcone of K . We consider the polyhedron defined by the nonzero, integral elements of C' ; i.e.,

$$P' := \text{conv}\{\mathbb{Z}^n \cap (C' \setminus \{0\})\}.$$

The facets of P' separating P' from the origin induce a partition of C' into subcones, each generated by elements of the minimal Hilbert basis for C' . If any of these facets has an integral element in its (relative) interior, this element can be used to further refine the partition. Moreover, by triangulating, we may assume that each cone in the partition is simplicial (see [4, Theorem 3.4] for details). We will thus assume that C' is partitioned into simplicial subcones, say $C' = \bigcup_{i=1}^p C_i$, with each C_i generated by elements of the minimal Hilbert basis of C' . Consider $x \in C_1 = \text{cone}\{b_1, \dots, b_n\}$, where b_1, \dots, b_n define a facet of P' , where b_i is in the minimal Hilbert basis for C' , with $b_i = \sum_{j=1}^n \lambda_{ij} a_j$, $\sum_{j=1}^n \lambda_{ij} \leq 1$ and $\lambda_{ij} \geq 0$, for $1 \leq i, j \leq n$.

Now x is in the minimal Hilbert basis for K , hence also for the subcones C' and C_1 . Thus x is in the parallelotope defined by b_1, \dots, b_n and we can write $x = \sum_{i=1}^n \alpha_i b_i$, where $0 \leq \alpha_i \leq 1$, for $1 \leq i \leq n$. If $\sum_{i=1}^n \alpha_i > n - 1$, then the element x' obtained by reflecting x from the apex, $\sum_{i=1}^n b_i$, of the parallelotope, i.e., $x' = (\sum_{i=1}^n b_i) - x = \sum_{i=1}^n (1 - \alpha_i) b_i$, would be an integral vector of C_1 , and hence of C' , not in P' . Thus we must have $\sum_{i=1}^n \alpha_i \leq n - 1$. Finally, since $x = \sum_{i=1}^n \alpha_i b_i = \sum_{i=1}^n \alpha_i (\sum_{j=1}^n \lambda_{ij} a_j) = \sum_{j=1}^n (\sum_{i=1}^n \alpha_i \lambda_{ij}) a_j$, with $\sum_{i=1}^n \sum_{j=1}^n \alpha_i \lambda_{ij} \leq n - 1$, it follows that the height of x is bounded above by $n - 1$.

Furthermore, if $\sum_{i=1}^n \alpha_i = n - 1$, then the reflected point x' has to coincide with one of the vectors b_j , by construction. But this implies that $x = \sum_{i \neq j} b_i$, which is not an element of the minimal Hilbert basis for $n \geq 3$.

To see that the bound is sharp, let $d > 1$ be an integer and suppose $K \subseteq \mathbb{Q}^n$ is the simplicial cone generated by $(1, \dots, 1, d)$ and the $n - 1$ unit vectors e_1, \dots, e_{n-1} . It is not difficult to see that the minimal Hilbert basis for K consists of the d vectors $\frac{t}{d}(1, \dots, 1, d) + \sum_{i=1}^{n-1} \frac{d-t}{d} e_i$, for $0 \leq t \leq d - 1$, along with the n generators of K . The vector of maximum

height is achieved for $t = 1$, i.e., for $x = \frac{1}{d}(1, \dots, 1, d) + \sum_{i=1}^{n-1} \frac{d-1}{d} e_i$. Thus $h(x) = \frac{1}{d} + \frac{(n-1)(d-1)}{d} = (n-1) - \frac{n-2}{d}$, so that as d increases, $h(x)$ is arbitrarily close to $n-1$. \square

The class of examples used in the proof to demonstrate that the height bound is as tight as possible was also discovered by Ewald and Wessels [2]. Note that for these examples, the nonzero vectors in the parallelotope for K coincide with the minimal Hilbert basis. Such *parallelotope cones* are studied further in Liu [3].

Finally, we remark that the proof here still applies for any minimal Hilbert basis (not necessarily unique) when K is not pointed; in this situation the height of any element of any minimal Hilbert basis is less than $n-1$.

References

- [1] GÜNTER EWALD: personal communication, 1990.
- [2] GÜNTER EWALD & UWE WESSELS: *On the ampleness of invertible sheaves in complete projective toric varieties*, *Results in Mathematics* **19** (1991), 275-278.
- [3] JIYONG LIU: Hilbert Bases with the Carathéodory Property, Ph. D. Thesis, Cornell University, Ithaca NY, 1991.
- [4] JIYONG LIU & LESLIE E. TROTTER, JR.: *Hilbert covers and partitions of rational, polyhedral cones*, preprint, Cornell University 1991.
- [5] ALEXANDER SCHRIJVER: *Theory of Linear and Integer Programming*, John Wiley and Sons, Chichester, 1986.

JIYONG LIU, LESLIE E. TROTTER, JR.
 School of OR&IE
 Cornell University
 Ithaca, New York 14853, USA

GÜNTER M. ZIEGLER
 ZIB-Berlin
 Heilbronner Str. 10
 W-1000 Berlin 31, Germany