May 1983



A.E. Res. 83-20

THE ECONOMIC POTENTIAL OF CROP ROTATIONS IN LONG ISLAND POTATO PRODUCTION

S.S. Lazarus

G.B. White

Department of Agricultural Economics Cornell University Agricultural Experiment Station New York State College of Agriculture and Life Sciences A Statutory College of the State University Cornell University, Ithaca, New York, 14853

It is the policy of Cornell University actively to support equality of educational and employment opportunity. No person shall be denied admission to any educational program or activity or be denied employment on the basis of any legally prohibited discrimination involving, but not limited to, such factors as race, color, creed, religion, national or ethnic origin, sex, age or handicap. The University is committed to the maintenance of affirmative action programs which will assure the continuation of such equality of opportunity.

Ŀ

TABLE OF CONTENTS

Page
INTRODUCTION
METHODOLOGY
RESULTS
SUMMARY AND CONCLUSIONS 17
REFERENCES
APPENDIX A 23
Appendix Table A-1: Potato Budget 24
Appendix Table A-2: Rye Budget 25
Appendix Table A-3: Corn Budget 26
Appendix Table A-4: Winter Wheat Budget
Appendix Table A-5: Soybean Budget 28
Appendix Table A-6: Oats Budget 29
Appendix Table A-7: Sunflower Budget 30
Appendix Table A-8: Dry Bean (Red Kidney) Budget
Appendix Table A-9: Cauliflower Budget
Appendix Table A-10: Cabbage Budget
Appendix Table A-11: Onion Budget 35
Appendix Table A-12: Spinach Budget
Appendix Table A-13: Machinery Complement for Potato Production
Appendix Table A-14: Distribution of Annual Labor Needs by Semi-Monthly Time Periods

LIST OF TABLES

	<u>ra</u>
TABLE 1 - Optimal rotations with various limitations on maximum potate acreage (field crop rotations - North Fork Model.	» 6
TABLE 2 - Optimal rotations with various limitations on maximum potato acreage (field crop and cole crop rotations) - North Fork Model.	8
TABLE 3 - Optimal rotations with various limitations on maximum potato acreage (field crop rotations) - South Fork Model.	9
TABLE 4 - Optimal rotations with various limitations on maximum potato acreage (field crop and cole crop rotations) - South Fork Model.	11
TABLE 5 - Reduced income if an acre of various crop rotations was raised, potato acreage not constrained.	13
TABLE 6 - Optimal rotations if chemical costs for potatoes increase by various percentages.	15
TABLE 7 - Optimal rotations if potato yields decrease by various percentages.	16
TABLE 8 - Optimal rotations if potato yields in rotations increase by various percentages.	18
TABLE 9 - Optimal rotations for the North and South Fork models (all rotations).	19

LIST OF FIGURES

FIGURE 1 - Returns above variable costs for various potato acreage 12 constraints.

Page

THE ECONOMIC POTENTIAL OF CROP ROTATIONS IN LONG ISLAND POTATO PRODUCTION S.S. Lazarus and G.B. White*

INTRODUCTION

Suffolk County, the easternmost county on Long Island, has the highest value of farm receipts of any county in New York State. One of the major agricultural commodities raised there is potatoes. In 1981, 18,500 acres of potatoes were produced. However, in recent years potato production has declined. (For example, in 1970 potato acreage was 31,000.) There are several reasons for this decrease. Urban encroachment is a major problem.

Potato pests (disease, insects, and weeds) also cause many problems for Long Island growers. Since potato production is very intense on the island, pest populations tend to build up. It is believed that more pesticides are used per acre on Long Island potato fields than on potatoes in any other region of the United States. (For one estimate of per acre costs in various potato producing regions, see Putnam, 1981.) The insects on Long Island have become resistant to some pesticides and new effective chemicals must constantly be sought.

Aldicarb (Temik), a systemic insecticide, was widely used to control the Colorado potato beetle. In the late 1970's it was discovered that the ground water had become contaminated with aldicarb. In 1980 the use of aldicarb was banned on Long Island. Heavy use of other pesticides may also cause ground water contamination.

The withdrawal of aldicarb has caused an awareness of some of the problems of intense pesticide use. Continuous potato production has, in the past, been an economical practice for the fertile Long Island land; it may not be economical in the future given the pest management options now available to growers.

Integrated Pest Management (IPM) is a potential solution to some of the potato production problems on Long Island. IPM is the use of chemical, cultural, genetic, and biological pest control methods. These techniques are used in such a way as to have a minimum effect on nontarget organisms and the environment (Apple et al., 1979).

One IPM strategy that reduces pesticide use and incorporates other pest management tactics is crop rotation. Crop rotation can help reduce the population of potato pests. Crop rotation on Long Island potato farms will not become a major IPM technique until several economic questions are answered: 1) How will crop rotation affect growers' net income? 2) Are sufficient labor and other needed inputs available for other rotations? and 3) Are there markets for all crops raised in the rotations?

*Research Support Specialist and Assistant Professor, Department of Agricultural Economics, Cornell University, Ithaca, NY 14853-0398. This research was supported by Cooperative Agreement No. 58-32U4-2-389 between the Agricultural Research Service, USDA, and Cornell University. This paper focuses on the first of these three questions. A model was developed to examine the profitability of various crop rotations that are agronomically feasible for Long Island. It is important to examine the broad question of the economic feasibility of rotation as an IPM strategy on Long Island. Insufficient labor to raise certain vegetables can then be examined within the perspective of the relative profitability of these crops. Likewise solutions to various marketing problems may be found if certain crops are found to have economic potential. These questions may be addressed in the future depending upon the results of this research to evaluate the effects of crop rotation on growers' incomes.

METHODOLOGY

A linear programming model was used to determine the best crop production plan. Linear programming is a mathematical programming technique in which an optimal mix of production methods and crops is derived by allocating limited resources (such as land, labor, and capital) among various production activities to achieve the highest net return over variable costs. Additional analyses of a linear program solution can indicate when an opticides), a change in the market price of a crop, or a change in the quantity of resources used.

There are two major areas that produce potatoes on Long Island. Both are on the eastern portion of the Island. One area is commonly called the South Fork, the other the North Fork. Linear programming models were developed for representative 150 acre farms on each Fork. This was necessary because the Forks have different soil types, and growers use different cultural practices on the two Forks. The soil on the North Fork is very light and irrigation is required to raise potatoes. Many growers on the North Fork have traditionally raised continuous potatoes. Rye, which is planted as a cover crop to prevent wind erosion during the winter, is plowed down in the spring. Although the land on the North Fork is well suited for the production of various vegetable crops, many potato growers prefer not to raise vegetables due to the problems of hiring seasonal labor.

Compared to the North Fork, the soil on the South Fork is heavier. Irrigation is not widely used to grow potatoes, due to the greater waterholding capacity of the soil. Even though many South Fork growers do not irrigate, potato yields are estimated to be approximately 5-10 percent higher than on the North Fork. Thus, South Fork growers have lower costs because they do not need irrigation equipment. At the same time they receive higher gross returns due to the higher yields. Growers on the South Fork have traditionally raised two years of potatoes followed by a year of rye. Like the growers on the North Fork, they plant a rye cover crop, but allow it to mature every third year. Few South Fork potato growers raise vegetables. In addition to the labor problems, irrigation equipment may need to be purchased to grow vegetables economically.

There are major difficulties hiring seasonal labor to weed and harvest vegetable crops. Migrant labor is typically used to harvest vegetable crops, and many New York State and federal laws regulate the employment of migrants. Many growers who have traditionally specialized in potato production lack the managerial expertise or are not inclined to handle a seasonal labor crew. In the past, these farms have relied largely upon family labor or used full-time hired employees. Potato growers have a strong preference to continue operating in a similar manner. There are some farmers on Long Island who raise large quantities of vegetables in spite of the labor difficulties. This implies that it would be possible for potato farmers to surmount the labor difficulties and raise vegetable crops.

Another problem with some rotations is that potatoes are raised on soil with a low pH. The pH is kept low to reduce the incidence of potato scab. Rye, cauliflower and cabbage are three crops that, like potatoes, can be grown on a low pH soil. However, fields planted to cauliflower or cabbage are generally limed prior to planting. These three crops are relatively common on Long Island. But many other crops require a higher pH to produce a high yield. It is possible to raise the soil pH slightly to allow the production of these crops, yet not so much that potato scab would be a major problem the following year.

The following rotations using various potato and field crop combinations were considered in the model:

- 1) Year 1 Potatoes; Year 2 Rye
- 2) Year 1 Potatoes; Year 2 Corn
- 3) Year 1 Potatoes; Year 2 Double Crop of Winter Wheat and Soybeans
- 4) Year 1 Potatoes: Year 2 Double Crop of Winter Wheat and Soybeans;
- Year 3 Corn
- 5) Year 1 Potatoes; Year 2 Oats
- 6) Year 1 Potatoes; Year 2 Sunflowers

7) Year 1 - Potatoes; Year 2 - Dry Beans (Red Kidney Beans)

In addition to these rotations, continuous potatoes (the production practice currently used by many growers) was considered for the North Fork. Two years of potatoes followed by a year of rye was considered to be the traditional rotation in the South Fork model.

Some growers on Long Island raise large quantities of cabbage and cauliflower in spite of the labor difficulties. Thus, in some versions of the model two vegetable crop rotations were considered in addition to the field crop rotations:

- 1) Year 1 Potatoes; Year 2 Cauliflower
- 2) Year 1 Potatoes; Year 2 Cabbage

The models were also run to include two other rotations which contain crops not commonly grown on Long Island, but which are possibilities in the future:

1) Year 1 - Potatoes; Year 2 - Onions

2) Year 1 - Potatoes; Year 2 - Double Crop of Spinach and Soybeans

These two rotations may have some agronomic problems due to the higher pH soils required by the onions and spinach. The utilization of scab resistant varieties will help make these rotations possible. However, presently available scab resistant varieties can develop scab under severe conditions. Spinach is currently raised by some growers on Long Island. It takes somewhat less labor than many other vegetable crops. A rotation of spinach followed by soybeans, a less labor intensive field crop, might provide a compromise between the high labor needs of vegetable crops and the relatively low per acre return of some field crops.

Budgets were constructed for each of the crops considered in the models (Appendix Tables Al through Al2). Information for these budgets was gathered from a variety of sources. Three Long Island potato growers were interviewed to discover their current crop raising practices, as well as their costs and returns for potato production. Average yields and prices over the past five years for various Long Island crops were obtained from New York Agricultural Statistics. If the information was not available for Long Island, average New York State data were used. Cost data for field crops was obtained from Knoblauch (1981). Revenue and cost data for sunflowers was obtained from W. Lazarus (1982). Pesticide usage for crops other than potatoes were estimated from Cornell Recommends for Field Crops and Cornell Recommendations for Commercial Vegetable Production. The potato pesticide usage was obtained from 1981 surveys of Long Island potato growers participating in a Cornell-sponsored IPM program. Additional information about production practices for vegetable crops was obtained from Dhillon (1979), Phelps and How (1981), and Snyder (1981). Prices were obtained from several Long Island input suppliers.

Labor and machinery costs were estimated for each crop. An economic engineering approach was used. Using this approach, machinery costs and labor inputs were calculated based on such factors as machine width, operating speed, and machine efficiency (Benson 1974; Knoblauch, et al., 1980).

The representative farm was assumed to have the machinery complement presented in Appendix Table A13. The farm was assumed to have sufficient machinery to plant the entire farm in potatoes since this crop was currently grown. The farm was also assumed to have sufficient machinery to raise the various vegetable crops considered in some of the model variations. Since raised some vegetables in the past and thus would have the necessary

Custom corn planting, custom combining, and custom grain drying were assumed for the rotations requiring these operations. The use of custom machinery is a way to avoid the problem of having too few acres of a particular crop to justify (economically) the purchase of a specialized machine. A grower trying a new rotation with just a few acres of a field crop is not likely to purchase an expensive machine to produce that crop. The South Fork model farm, however, was assumed to purchase an irrigation system to irrigate any vegetable crops produced. It is handled as a continuous input rather than using an assumed equipment capacity.

The labor requirements for various crops are presented in Appendix Table Al4. It was assumed that the grower and his family could provide 217 hours of labor during each semi-monthly period. This is the equivalent of two people each working a 50 hour week. Additional labor could be hired for \$5.50 per hour (wages, taxes, and benefits).

The farm could borrow operating capital at a 12 percent annual rate for nine months. The various crops were assumed to be sold at harvest. The modeled farms could either raise the rye that is used as seed for the cover crop or buy the seed. It would cost \$5.00 per bushel to buy rye seed. Excess rye could be sold for \$2.80 per bushel. In the models which included vegetable crop rotations, the acreage of any one crop was limited to 25 acres. This was done because of the price risk of having a large acreage in any one vegetable crop. The yield of potatoes was assumed to be five percent higher on the South Fork than on the North due to the better soil. The vegetable crops were assumed to have the same yield on both Forks. Field crops which were assumed to be irrigated on both Forks had the same yields, but yields of nonirrigated field crops were assumed to be 10 percent higher on the South Fork than on the North Fork.

The models were also run to examine what affect various acreage limitations on potato production would have on returns over variable costs. The model first allowed all acreage (150 acres) on the North Fork and 100 acres on the South Fork to be planted to potatoes. Maximum potato acreage was then reduced by increments of 25 acres in successive runs. This procedure examined possible reductions in farm returns over variable costs of using crop rotation as an IPM tactic.

Potato yields in the future might stay at current levels at the cost of relatively higher pesticide costs for potato crops. So a model variation was run that examined higher potato pesticide usage while the quantities of chemicals used for other crops was held constant. Another model variation examined the effect of potato yield reductions if chemicals no longer effectively controlled the Colorado potato beetle. It is also possible that potato yields in the past have been somewhat suppressed due to the high intensity of potato production. Another variation of the model examined the effect of yield increases for potatoes grown in a rotation.

RESULTS

If only field crop rotations were considered, continuous potato production was the most profitable cropping practice on the North Fork (Table 1). If all 150 acres of the farm were planted to potatoes, the highest return over variable costs would be attained. This result is not surprising since growers can be expected to have found a profitable cropping pattern by trial and error, and since existing machinery is geared to potato production. Relatively little seasonal labor must be hired. Large amounts of pesticides must be used to produce continuous potatoes.

Growers might be willing to raise other field crops if they could be subsidized for the income loss of not raising continuous potatoes. Pesticide use could be reduced if only two-thirds or half of each farm's acreage

Table 1.	Optimal	market and a
	operman	rotations with various limitations and set
	acrease	(field anon potato
	-crca5c	(iteld crop rotations) - North Fork Model

		Maxir	um Potato	Acreage	
	150	125	100	75	50
Rotation:					
Continuous Potatoes	1.5/) 1 00			
(1)Potatoes (2)Rye		- 10	50		
(1)Potatoes (2)Corn		- 12	12	12	8
(1)Potatoes (2)Winter Wheat/Soubon					
(1)Potatoes (2)Winter Wheat/Soybea	ns	- 38	88	138	
(1) Potatoes (2) Octo		·		· · · · · · · · · · · · · · · · · · ·	130
(1) Potatoos (2) San Si		•			
(1) Potatoos (2) Due n					
(1) Iotaloes (2) Dry Beans					
Actual Number of Acres in Potatoes	150	125	100	75	50
Return over Variable Costs	\$101,08	8 \$88,618	\$75,471	\$61,972	\$44 92
Family Labor Activity Land W				, ,	· • • • • • • • • • • • • • • • • • • •
March (second half)					
April (first half)	147	123	98	74	50
April (second helf)	147	123	98	74	50
May (first half)	165	138	110	83	50
May (nonord half)	15	13	10	8	26
June (finat half)	15	13	10	8	14
June (nerest 1 1 1 s)	63	.48	32	17	19
July (Second nalf)	199	173	148	124	12
July (first half)	167	160	159	150	83
July (second half)	167	140	112	100	149
August (first half)	217	217	202	157	98
August (second half)	217	217	202	157	14/
September (first half)	144	120	05	10/	147
September (second half)	144	124	104	/1	47
October (first half)	144	130	110	85	47
October (second half)	144	126	100	107	94
		140	109	93	84
red Labor Activity Level, Hours					
ugust (first half)	76	30			
ugust (second half)	76	30			
her Major Activities					
orrow operating canital	27 600	41.07 0.00		÷	
uy rye seed (bu.)	225	\$107,966	\$90,390	\$73,140	\$58,268
sticide Activo Inconti					
ungicide (lba A T)					
$\frac{1}{100} = \frac{1}{100} = \frac{1}$	1,907	1,589	1,271	953	630
asecutide (IDS. A.I.)	, 047	3,314	2,580	1.8/7	1 /1/
colcide (lbs. A.T.)	000	1 - 2 - 2	-,	+ ; 047	1,410

-

was planted to potatoes resulting in less risk of ground water contamination. The Colorado potato beetle is less likely to develop resistance to various chemicals that are used less intensively.

If a grower on the North Fork did not raise potatoes on more than half of his acreage in any given year, the optimal cropping pattern would contain two rotations (Table 1). A potato/rye rotation would be raised on 12 acres (i.e. six acres of potatoes and six acres of rye). A rotation of potatoes followed the next year by a double crop of winter wheat and soybeans would be planted on the remaining 138 acres of the farm.¹ The return over variable costs was reduced by \$39,116 if potato acreage was restricted by 50 percent. Pesticide use would be cut in half.

The potato/rye rotation was raised on just enough acreage to provide seed for the cover crop in all cases where potato acreage was restricted (Table 1). This was due to the dual pricing system used for rye in the model. Rye seed could be purchased for \$5.00 per bushel, but the grower could sell excess rye which he raised for only \$2.80 per bushel.

If both field and cole crop rotations were considered in the North Fork model, continuous potatoes would be planted on 100 acres in the optimal solution (Table 2). A two year rotation of potatoes and cauliflower would be planted on the remaining 25 acres. The return over variable costs would be \$107,515. The 25 acres of cauliflower would provide the farm with a higher return than if continuous potatoes were raised on the entire farm. The cauliflower acreage required considerable hired labor. This modeled farm was representative of the current situation on some North Fork potato farms where some high income vegetable crops are grown.

If the maximum potato acreage was limited to half of the farm in any given year, three rotations would be raised (Table 2). Twelve acres of the two year potatoes and rye rotation would be raised. Eighty-eight acres of the two year rotation of the potatoes and double crop (winter wheat, soybeans) would be raised. The remainder of the farm (50 acres) would be planted in the potatoes and cauliflower rotation. The return over variable costs was \$24,600 less than if no restrictions were placed on potato acreage. Pesticide use was reduced by 23 percent from the optimal plan. A constraint limited the acreage of any one vegetable crop to 25 acres, due to the price risk of raising vegetables. If the cauliflower acreage had not been limited, even more acres of the potato/cauliflower rotation would have been raised in all situations.

The model results on the South Fork were similar to those on the North Fork if only field crop rotations were considered (Table 3). The traditional cropping pattern of two years of potato production followed by a year of rye would be used for the entire 150 acres. The return over variable costs would be \$91,640.

¹A grower would probably set up the rotations so that of the 138 acres, there would be 69 acres of potatoes and 69 acres in double-cropping in any one year. In an average year, potato acreage on the farm would be six acres (from the potato/rye rotation) plus 69 acres from the potato/double crop rotation, or a total of 75 acres of potatoes.

lable 2.	Optimal	rotations with various limitations on mendation
	acreage	(field crop and cole grop notations) in maximum potato
	-	North Fork Model

$\begin{array}{c c c c c c c c c c c c c c c c c c c $		<u></u>	Max	imum Pota	to Acreage	
Retation: 100 100 50		15	0 12	5 100	75	50
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	Rotation:				······	
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	Continuous Potatoes	10) 0 1(۰ ۵ -		
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	(1)Potatoes (2)Rye					
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	(1)Potatoes (2)Corn			1	.2 12	2 8
(1)Potatoes (2)Winter Wheat/Soybeans 38 88 (3)Corn (1)Potatoes (2)Oats	(1)Potatoes (2)Winter Wheat /Sou			· · · · ·		
(1)Potatoes (2)Oats	(1)Potatoes (2)Winter Wheat/Soy (3)Corn	beans beans		3	8 88	3
(1)Potatoes (2)Sunflowers	(1)Potatoes (2)Oats					• 63
(1)Potatoes (2)Dry Beans	(1)Potatoes (2)Sunflowers					•
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	(1)Potatoes (2)Dry Board	**# 				·
C1)Pointoos (2)Callifuser505050505050Actual Number of Acres in Potatoes1251251007550Return over Variable Costs $$107,515$ $$95,505$ $$82,915$ $$64,700$ Family Labor Activity Level, Hours123123987449March (second half)123123987449April (second half)138138110860May (second half)13131089June (first half)4747321611June (second half)172172172172139July (first half)179179172172139July (first half)217217217217217August (first half)217217217217217August (second half)217217217217217Cotober (first half)217217217217217October (first half)217217217217217Obvember (first half)217217217217217Obvember (first half)709709689669637Staber (first half)709709689667641October (second half)224224178133100eptember (first half)693693673651127103overber (first half)693 <td>(1)Potatoes (2)Coult flow</td> <td></td> <td></td> <td></td> <td>······</td> <td></td>	(1)Potatoes (2)Coult flow				······	
Actual Number of Acres in Potatoes1251251007550Return over Variable Costs $\$107,515$ $\$107,515$ $\$95,505$ $\$82,915$ $\$64,700$ Family Labor Activity Level, Hours March (second half)123123987449April (first half)123123987449April (second half)131310814May (first half)131310814June (second half)17217214512180June (second half)17217214512180June (second half)17217214512180June (second half)17217217213913July (second half)171217217217217July (second half)217217217217217September (first half)217217217217217September (first half)217217217217217October (second half)217217217217217Ovember (first half)217217217217217Ovember (first half)224224178133100eptember (second half)224224178133100eptember (first half)709709689669637ctober (second half)224224178133100eptember (firs	(1)Potatoog (2) Call	5	0 5	0 50	0 50	50
Actual Number of Acres in Potatoes1251251007550Return over Variable Costs $\$107,515$ $\$107,515$ $\$95,505$ $\$82,915$ $\$64,700$ Family Labor Activity Level, HoursMarch (second half)123123987449April (first half)123123987449April (second half)1381381108360May (second half)13131089June (first half)4747321611June (second half)17217214512180July (first half)179179172172139August (first half)217217217217217August (second half)217217217217217September (first half)171217217217217September (first half)217217217217217October (first half)217217217217217Ovember (second half)209209209209209Ovember (first half)216176151127103Ovember (second half)217217217217217217Otober (first half)216224224178133100eptember (second half)709709689669637ctober (first half)693693678667641<	(1) ocalles (2) Cabbage	<u>.</u>				
Return over Variable Costs\$107,515 \$107,515 \$95,505 \$82,915 \$64,700Family Labor Activity Level, Hours123 123 98 74 49March (second half)123 123 98 74 49April (first half)123 13 10 83 60May (first half)13 13 10 8 14May (second half)13 13 10 8 9June (first half)172 172 145 121 80June (first half)179 179 172 172 180June (second half)179 179 172 172 180July (first half)177 217 217 217 217 217July (second half)217 217 217 217 217 217 217August (first half)217 217 217 217 217 217August (second half)217 217 217 217 217 217September (first half)217 217 217 217 217 217September (second half)217 217 217 217 217 217October (first half)217 217 217 217 217November (second half)224 224 178 133 100eptember (second half)224 224 178 133 100eptember (second half)709 709 689 669 637ctober (first half)709 709 689 569 637ugust (second half)53 53 53 53 53solut (second half)52 225	Actual Number of Acres in Potato	es 12	5 12	5 100) 75	50
Family Labor Activity Level, Hours March (second half) 123 123 98 74 49 April (first half) 123 123 98 74 49 April (first half) 138 133 10 83 60 May (first half) 13 13 10 8 14 June (first half) 17 172 172 16 11 June (second half) 172 172 145 121 80 July (first half) 179 179 172 172 139 August (first half) 217 217 217 217 217 217 August (second half) 217 217 217 217 217 217 September (first half) 217 217 217 217 217 217 October (second half) 217 217 217 217 217 217 November (first half) 209 209 209 209 209 209 209 209 209 209 209 209	Return over Variable Costs	\$107,5	15 \$107,51	15 \$95,50)5 \$82,91	5 \$64,700
March (second half)123123987449April (first half)123123987449April (second half)1313108360May (first half)131310814May (second half)13131089June (first half)4747321611June (second half)17217214512180July (first half)179179172172139July (second half)209209182154145August (first half)217217217217217August (first half)217217217217217September (second half)217217217217217October (first half)217217217217217Ovember (first half)209209209209209Sovember (second half)217217217217Vovember (first half)209209209209209Sovember (second half)5353535353red Labor Activity Level, Hours828236ugust (first half)176176151127103eptember (first half)683693678667641ctober (second half)693693678667641ctober (second half)608<	Family Labor Activity Level, Hour	s				, , , , , , , , , , , , , , , , , , , ,
April (first half)123123987449April (second half)1381381108360May (first half)131310814June (first half)13131089June (first half)17217214512180July (first half)17217214512180July (first half)179179172172139July (second half)209209182154145August (first half)217217217217217August (second half)217217217217217September (first half)171217217217217Cober (second half)217217217217217October (second half)217217217217217Obvember (first half)217217217217217Obvember (first half)217217217217217Obvember (first half)224224178133100eptember (first half)176176151127103eptember (first half)693693678669637ctober (second half)693693678669637ctober (second half)693693678667641optober (first half)693693678667641ctober	March (second half)	123	} 123	3	, ,	
April (second half)1381381381108360May (first half)131310814June (first half)13131089June (second half)17217214512180July (first half)17917217214512180July (second half)209209182154145August (first half)217217217217217August (second half)217217217217217September (first half)171217217217217Detober (second half)217217217217217Detober (first half)217217217217217Detober (second half)217217217217217Detober (second half)217217217217217Detober (second half)217217217217217Detober (second half)5353535353ugust (first half)176176151127103eptember (first half)176176151127103eptember (first half)693693678667641ctober (second half)709709689669637ctober (second half)608608589573551her Major Activities225225225225 <td< td=""><td>April (first half)</td><td>12</td><td>123</td><td></td><td>2 74</td><td>49</td></td<>	April (first half)	12	123		2 74	49
May (first half)1313108360May (second half)131310814June (first half)1717171216June (second half)17217214512180July (first half)179179172172139July (second half)209209182154145August (first half)217217217217217August (second half)217217217217217September (second half)217217217217217September (second half)217217217217217October (first half)217217217217217October (first half)217217217217217November (second half)209209209209209November (first half)209209209209209November (first half)224224178133100eptember (first half)176176151127103eptember (first half)693693678667641ctober (first half)693693678667641ctober (first half)686689573551her Major Activities51872\$151,872\$131,722\$113,630\$92,846yr yes eed (bu.)225225	April (second half)	138	120	· 90	/4	49
May (second half)131310814June (first half)4747321611June (second half)17217214512180July (first half)1791721721391089July (first half)17917214512180July (second half)209209182154145August (first half)217217217217217August (second half)217217217217217September (first half)171217217217217Detober (second half)217217217217217Detober (second half)217217217217217Detober (second half)217217217217217Detober (second half)209209209209209November (first half)209209209209209November (second half)5353535353red Labor Activity Level, Hours828236ugust (second half)709709689669637ctober (first half)176176151127103eptember (first half)683693678667641ctober (second half)709709689669637ctober (second half)683693678667641	May (first half)	19	່ 1.30	, TIO	83	60
June (first half) 13 10 8 9 June (second half) 172 173 10 8 9 July (first half) 172 172 145 121 80 July (second half) 209 209 182 154 145 August (first half) 217 217 217 208 175 August (second half) 217 217 217 217 217 September (first half) 217 217 217 217 217 September (second half) 217 217 217 217 217 October (first half) 217 217 217 217 217 October (first half) 217 217 217 217 217 October (first half) 209 209 209 209 209 November (first half) 209 209 209 209 209 November (first half) 224 224 178 133 100 eptember (second half) 709 709 689 669 637 ctober (first half) 176 176 151 127 103 eptember (second half) 224 224 236 $$ ugust (second half) 709 709 689 669 637 ctober (first half) 1633 693 678 667 641 eptember (second half) 693 678 667 641 tober (second half) 69	May (second half)	. 13	1 L J	10		14
June (second half) 47 47 47 47 47 32 16 11 July (first half) 172 172 172 145 121 80 July (second half) 209 209 172 172 139 August (first half) 217 217 217 208 154 August (first half) 217 217 217 217 217 September (first half) 217 217 217 217 217 September (second half) 217 217 217 217 217 October (first half) 217 217 217 217 217 October (second half) 217 217 217 217 217 November (first half) 209 209 209 209 209 November (first half) 209 209 209 209 209 November (second half) 53 53 53 53 53 red Labor Activity Level, Hours 82 82 36 $$ ugust (second half) 224 224 178 133 100 eptember (first half) 176 151 127 103 eptember (second half) 693 693 678 667 641 cober (second half) 608 608 589 573 551 her Major Activities 225 225 $$ $$ orrow operating capital tay rye seed (bu.) $1,639$ $1,63$	June (first half)	1	13	10	8	9
July (first half) 172 172 145 121 80July (second half) 209 209 182 154 145 August (first half) 217 217 217 217 217 August (second half) 217 217 217 217 217 September (first half) 171 217 217 217 217 September (second half) 217 217 217 217 217 October (second half) 217 217 217 217 217 October (second half) 217 217 217 217 217 November (first half) 209 209 209 209 209 November (second half) 53 53 53 53 53 red Labor Activity Level, Hours 82 82 36 $$ ugust (first half) 176 176 151 127 103 eptember (first half) 176 176 151 127 103 eptember (second half) 709 709 689 669 637 ctober (second half) 693 693 678 667 641 ctober (second half) 608 608 589 573 551 her Major Activities 225 225 $$ $$ otrow operating capital $\$151, 872$ $\$131, 722$ $\$13, 630$ $\$92, 846$ 225 225 225 $$ $ -$ otrow oper	June (second half)	47	4/	32	16	11
July (second half) $1/9$ $1/79$ 172 172 139 August (first half) 209 209 182 154 145 August (second half) 217 217 217 217 208 175 September (first half) 171 217 217 217 217 217 217 September (second half) 217 217 217 217 217 217 217 October (first half) 217 217 217 217 217 217 217 October (first half) 209 209 209 209 209 209 November (first half) 209 209 209 209 209 November (second half) 53 53 53 53 53 .red Labor Activity Level, Hours 82 82 36 $$ ugust (second half) 224 224 178 133 100 eptember (first half) 176 176 151 127 103 eptember (second half) 709 709 689 669 637 ctober (second half) 693 693 678 667 641 ctober (second half) 608 608 589 573 551 her Major Activities $5151,872$ $$131,722$ $$113,630$ $$92,846$ 225 225 $$ $$ $$ atticide Active Ingredients $1,639$ $1,639$ $1,321$ $1,003$ 687 <	July (first half)	172	1/2	145	121	80
August (first half) 209 209 182 154 145 August (second half) 217 217 217 217 208 175 September (first half) 171 217 217 217 217 217 217 September (second half) 217 217 217 217 217 217 217 October (first half) 217 217 217 217 217 217 217 November (first half) 209 209 209 209 209 209 November (second half) 53 53 53 53 53 red Labor Activity Level, Hours 82 82 36 $$ ugust (second half) 224 224 178 133 100 eptember (first half) 176 176 151 127 103 eptember (second half) 709 689 669 637 ctober (first half) 693 678 667 641 ctober (second half) 693 678 667 641 ctober (second half) 608 608 589 573 551 her Major Activities 225 225 $$ $$ orrow operating capital $$151,872$ $$131,722$ $$113,630$ $$92,846$ uy rye seed (bu.) 225 225 $$ $$ usciticide Active Ingredients $1,639$ $1,639$ $1,321$ $1,003$ 687 mgicide (lbs. A.I.) <td< td=""><td>July (second half)</td><td>1/9</td><td>179</td><td>172</td><td>172</td><td>139</td></td<>	July (second half)	1/9	179	172	172	139
August (linkt half) 217 217 217 217 217 208 175 August (second half) 217 21	August (first half)	209	209	182	154	145
August (second half) 217 <t< td=""><td>August (second half)</td><td>217</td><td>217</td><td>217</td><td>208</td><td>175</td></t<>	August (second half)	217	217	217	208	175
September (first half) 171 217 217 217 217 217 September (second half) 217 217 217 217 217 217 217 October (second half) 217 217 217 217 217 217 217 November (first half) 209 209 209 209 209 209 November (second half) 53 53 53 53 53 .red Labor Activity Level, Hours 82 82 36 $$ ugust (second half) 224 224 178 133 100 eptember (first half) 176 176 151 127 103 eptember (second half) 709 709 689 669 637 ctober (first half) 693 693 678 667 641 ctober (second half) 608 608 589 573 551 her Major Activities 225 225 $$ $$ orrow operating capital $$151,872$ $$131,722$ $$113,630$ $$92,846$ uy rye seed (bu.) 225 225 $$ $$ eticide Active Ingredients $1,639$ $1,639$ $1,321$ $1,003$ 687 upicide (lbs. A.I.) $1,639$ $1,639$ $1,321$ $1,003$ 687 secticide (lbs. A.I.) 775 775 648 522 272	September (first 1-1-1)	217	217	217	217	217
Depression217217217217217217Detober (first half)217217217217217217Detober (second half)217217217217217Detober (second half)209209209209209November (second half)5353535353Stred Labor Activity Level, Hours828236ugust (first half)224224178133100eptember (first half)176176151127103eptember (second half)709709689669637ctober (second half)693693678667641ctober (second half)693693678667641ctober (second half)693693678667641ctober (second half)608608589573551her Major Activities 225 225 225 225 225 eticide Active Ingredients $1,639$ $1,639$ $1,321$ $1,003$ 687uguicide (lbs. A.I.) $3,414$ $3,414$ $2,681$ $1,947$ $1,416$ 775775648522727272	Soptember (III'st half)	171	217	217	217	217
Detober (first half) 217 213 217 213 217 216 217 217	Deplember (second half)	217	217	217	217	217
October (second half) 217 2109 209 209	Detober (first half)	217	217	217	217	217
November (first half) 209 <	Scrober (second half)	217	217	217	217	217
November (second half)5353209209209red Labor Activity Level, Hours ugust (first half) 82 82 36 $$ ugust (second half) 224 224 178 133 100 eptember (first half) 176 176 151 127 103 eptember (second half) 709 709 689 669 637 ctober (first half) 693 693 678 667 641 ber Major Activities 608 608 589 573 551 ber Major Activities $$151,872$ $$151,872$ $$131,722$ $$113,630$ $$92,846$ atticide Active Ingredients $$1,639$ $1,639$ $1,321$ $1,003$ 687 atticide (lbs. A.I.) $1,639$ $1,639$ $1,321$ $1,003$ 687 atticide (lbs. A.I.) $3,414$ $3,414$ $2,681$ $1,947$ $1,416$	November (first half)	209	209	217	217	217
red Labor Activity Level, Hours 82 82 36 ugust (first half) 224 224 178 133 100 eptember (first half) 176 176 151 127 103 eptember (second half) 709 709 689 669 637 ctober (first half) 693 693 678 667 641 ctober (second half) 608 608 589 573 551 her Major Activities 608 608 589 573 551 orrow operating capital \$151,872 \$131,722 \$113,630 \$92,846 ugicide (lbs. A.I.) 1,639 1,639 1,321 1,003 687 sticide (lbs. A.I.) 1,639 1,639 1,321 1,003 687 sticide (lbs. A.I.) 775 775 648 520 252	lovember (second half)	53	53	53	209	209
august (first half) 82 82 36 $$ $$ $ugust$ (second half) 224 224 178 133 100 $eptember$ (first half) 176 176 151 127 103 $eptember$ (second half) 709 709 689 669 637 $ctober$ (first half) 693 693 678 667 641 $ctober$ (second half) 608 608 589 573 551 her Major Activities 608 608 589 573 551 orrow operating capital $$151,872$ $$151,872$ $$131,722$ $$113,630$ $$92,846$ 225 225 225 $$ $$ $$ sticide Active Ingredients $1,639$ $1,639$ $1,321$ $1,003$ 687 $ugust (106 (1bs. A.I.))$ $3,414$ $3,414$ $2,681$ $1,947$ $1,416$ 775 775 648 522 522	red Labor Activity Level, Hours				53	53
ugust (second half) 224 224 178 133 100 eptember (first half) 176 176 151 127 103 eptember (second half) 709 709 689 669 637 ctober (first half) 693 693 678 667 641 ctober (second half) 608 608 589 573 551 her Major Activities 608 608 589 573 551 her Major Activities $$151,872$ $$131,722$ $$113,630$ $$92,846$ orrow operating capital $$151,872$ $$131,722$ $$113,630$ $$92,846$ uy rye seed (bu.) $$225$ 225 $$ $$ sticide Active Ingredients $1,639$ $1,321$ $1,003$ 687 ugicide (lbs. A.I.) $1,639$ $1,3414$ $2,681$ $1,947$ $1,416$ 775 775 648 522 775 775	ugust (first half)	['] 82	82	36	·	
eptember (first half) 176 176 151 127 103 eptember (second half) 709 709 689 669 637 ctober (first half) 693 693 678 667 641 ber Major Activities 608 608 589 573 551 ber Major Activities $5151,872$ $$131,722$ $$113,630$ $$92,846$ ber Major Activities $$151,872$ $$131,722$ $$113,630$ $$92,846$ ber Major Active Ingredients $$1,639$ $1,639$ $1,321$ $1,003$ 687 ber Major Active Ingredients $$1,639$ $1,639$ $1,321$ $1,003$ 687 ber Major (lbs. A.I.) $3,414$ $3,414$ $2,681$ $1,947$ $1,416$ brown of the total (lbs. A.I.) 775 775 648 522	ugust (second half)	224	224	179	1 2 2	
eptember (second half) 709 709 689 669 637 ctober (first half) 693 693 678 667 641 ctober (second half) 608 608 589 573 551 her Major Activities 608 608 589 573 551 orrow operating capital $\$151, 872$ $\$131, 722$ $\$131, 630$ $\$92, 846$ aticide Active Ingredients 225 225 $$ $$ aticide (lbs. A.I.) $1, 639$ $1, 639$ $1, 321$ $1, 003$ 687 asecticide (lbs. A.I.) $3, 414$ $3, 414$ $2, 681$ $1, 947$ $1, 416$ 775 775 648 522 225 225	eptember (first half)	176	176	151	133	100
ctober (first half) 693 693 678 669 637 ctober (second half) 608 608 589 573 551 her Major Activities 608 608 589 573 551 her Major Activities \$151,872 \$131,722 \$113,630 \$92,846 bricide Active Ingredients \$151,872 \$131,722 \$113,630 \$92,846 sticide Active Ingredients 1,639 1,639 1,321 1,003 687 sticide (1bs. A.I.) 1,639 1,639 1,321 1,003 687 sticide (1bs. A.I.) 3,414 3,414 2,681 1,947 1,416 775 775 648 522 225 225 225 225	eptember (second half)	709	700	101	12/	103
ctober (second half) 673 673 667 641 her Major Activities 608 608 589 573 551 her Major Activities \$151,872 \$131,722 \$113,630 \$92,846 bricide Active Ingredients \$151,872 \$121 1,003 687 sticide (lbs. A.I.) 1,639 1,639 1,321 1,003 687 sticide (lbs. A.I.) 3,414 3,414 2,681 1,947 1,416 775 775 648 522 523 523	ctober (first half)	693	603	009	669	637
her Major Activities 500 589 573 551 her Major Activities \$151,872 \$131,722 \$113,630 \$92,846 by rye seed (bu.) 225 225 \$121,003 \$92,846 sticide Active Ingredients 1,639 1,639 1,321 1,003 687 secticide (1bs. A.I.) 3,414 3,414 2,681 1,947 1,416 rbicide (1bs. A.I.) 775 775 648 522 521	ctober (second half)	602 803	- 400	0/8	667	641
inter major Activities perrow operating capital iny rye seed (bu.) sticide Active Ingredients ingicide (lbs. A.I.) issecticide (lbs. A.I.)	har Major Astimiter	000	000	589	573	551
Stick operating capital \$151,872 \$131,722 \$113,630 \$92,846 Sticide Active Ingredients 225 225 sticide (lbs. A.I.) 1,639 1,639 1,321 1,003 687 sticide (lbs. A.I.) 3,414 3,414 2,681 1,947 1,416 rbicide (lbs. A.I.) 775 775 648 522	arrow apprendices					
sticide Active Ingredients ungicide (lbs. A.I.) 1,639 1,639 1,321 1,003 687 usecticide (lbs. A.I.) 3,414 3,414 2,681 1,947 1,416 urbicide (lbs. A.I.) 775 775 648 522	y rye seed (bu.)	\$151,872	\$151,872	\$131,722	\$113,630	\$92,846
IngledientsIngicide (lbs. A.I.)1,6391,3211,003687Issecticide (lbs. A.I.)3,4143,4142,6811,9471,416Irbicide (lbs. A.I.)775775648522775	sticide Active Increase		440	يبغر فيدغيه		
Institute (1bs. A.I.)1,6391,6391,3211,003687issecticide (1bs. A.I.)3,4143,4142,6811,9471,416rbicide (1bs. A.I.)775775648522	indicide (lba A T)	_				
rbicide (1bs. A.I.) 3,414 3,414 2,681 1,947 1,416 775 775 648 522	marciae (IDS. A.L.)	1,639	1,639	1,321	1.003	697
775 775 648 522 1,747 1,416	productice (10s. A.I.)	3,414	3,414	2,681	1 947	1 /12
	······································	775	775	648	500	1,410 960

		Maximum	Potato Ac	reage	
	150	125	100	75	50
Rotation:	· · · · · · · · · · · · · · · · · · ·				
(1) Patatoon (3) Rye	150	150	150		
(1)Potatoes (2)Potatoes (3)Mye				11	3
(1)Potatoes (2)Corn	·-				
(1)Potatoes (2)Uinter Wheat/Sovbeau	ns			139	
(1)Potatoes (2)Winter Wheat/Sovbea	as				717
(1)Potatoes (2)Winter wheat, co, co					147
(3) LOIN		·		<u> </u>	
(1) Potatoes (2) Sunflowers					
(1) Potatoes (2) Dry Beans					
(1) Polatoes (2) bij bedan					50
Astronal Number of Acres in Potatoes	100	100	100	15	50
Actual Number of Theres 211				+77 070	A55 044
Return over Variable Costs	\$91,6 40	\$ 91, 640	\$91,640	Ş/ /, 3/8	ŞJJ, 944
Return over variable totte					•
Remily Labor Activity Level, Hours			00	7/.	50
March (second half)	98	98	98	74	50
March (Second half)	98	.98	98	/4 02	66
April (record half)	110	110	110	ده د	25
April (second half) (f_{ij})	11	11	11	8	15
May (first half)	11	11	11	8	10
May (second nall)	32	32	32	1/	12
June (first nall)	42	42	42	55	3/
June (second half)	29	29	29	91	. 62
July (first half)	29	29	29	16	11
July (second half)	104	104	104	88	59
August (first half)	104	104	104	88	59
August (second half)	104	98	98	71	47
September (first half)	90	08	98	85	57
September (second half)	98	90	98	105	97
October (first half)	98	90	90	92	87
October (second half)	98	90	20		
Other Major Activities	+74 001	676 001	\$76 001	\$67.115	\$52,902
Borrow operating capital	\$76,001	\$70,001	1 900		
Sell rve seed (bu.)	1,900	1,900	1,900		91
Buy rye seed (bu.)					
· · · · · · · · · · · · · · · · · · ·					
Pesticide Active Ingredients	1 584	1.584	1,584	1,188	796
Fungicide (1bs. A.I.)	· · · · · · · · · · · · · · · · · · ·	2,903	2,903	2,147	1,628
Insecticide (lbs. A.I.)	2,70J 60¢	625	625	522	374
Herbicide (1bs. A.I.)	620	. 020	029		

Table 3. Optimal rotations with various limitations on maximum potato acreage (field crop rotations) - South Fork Model. Potatoes have traditionally been raised less intensively on the South Fork. The use of two years of potatoes followed by a year of rye as the traditional cropping pattern automatically limited potato production in any given year to two-thirds of the farm's acreage. If potato acreage was limited to half of the farm, 11 acres would be planted in the potato/rye rotation, providing enough seed for the cover crop. The remaining 139 acres would be planted in the potato/double crop (winter wheat, soybean) rotation. As noted before, growers would probably set up rotations so that approximately half the farm would be planted to potatoes each year. The return over variable costs would be \$77,378, which was \$14,262 less than if no restrictions were placed on potato acreage.

If both field and cole crop rotations were considered in the South Fork model, 100 acres would be planted in two years of potatoes, followed by a year of rye (Table 4). The remaining 50 acres would be planted in a potato and cauliflower rotation. The return over variable costs was \$106,833. The use of pesticides was relatively high.

Pesticide use was reduced by 17 percent if the maximum potato acreage was restricted to half of the farm (Table 4). In this situation, 11 acres would be planted in the potato/rye rotation. The rotation of potatoes followed the next year by a double crop of winter wheat and soybeans would be planted on 89 acres. Potatoes followed by a year of cauliflower would be planted on 50 acres (an annual average of 25 acres of cauliflower, or the maximum acreage permitted by the model's constraints). The return over variable costs was \$97,479, which was \$9,354 less than it would be if no restrictions were placed on potato acreage.

The results shown in Tables 1 through 4 were summarized in Figure 1. The returns were higher on both Forks if field and cole crop rotations were considered rather than just field crop rotations. The flat portions of the two curves for the South Fork in Figure 1 reflect the current practice of growing two years of potatoes followed by a year of rye, resulting in no more than 100 acres of potatoes in a given year. The curves show the tradeoff in returns above variable costs as potato acreage is reduced.

Another way to study the economic feasibility of various potential crop rotations is to examine the amount that the optimal return over variable costs would be reduced if the modeled farm was forced to plant an acre of a nonoptimal rotation. For example, Table 5 shows that in the North Fork field crop rotation model, income would be reduced by \$268.28 if an acre of the potato/dry beans rotation was forced into the solution. The greatest reduction in income would occur if an acre of the potato/oats rotation (\$310.56) or the potato/double crop (winter wheat, soybean)/corn rotation (\$372.15) was forced into the North Fork field crop model (Table 5).

With a corn price of \$2.75 (see Appendix Table A3) and assuming that variable costs were not increased as yields increased, it is possible to calculate the corn yield required for an acre of the corn rotations to come into the model. For an acre of the two year rotation of potatoes and corn to come into the optimal solution, a minimum corn yield of 210 bushels of dry, shelled corn for the North Fork field crop model and 149 bushels for the South Fork field crop model would be required. In both these scenarios, the corn yield required to make corn a profitable alternative is probably

					· · · · · · · · · · · · · · · · · · ·
		Maximum	Potato Ad	reage	
	150	125	100	/5	
Rotation:					
(1)Potatoes (2)Potatoes (3)Rye	100	100	100		
(1)Potatoes (2)Rye				11	
(1)Potatoes (2)Corn					
(1)Potatoes (2)Winter Wheat/Soybea	ns			89	
(1)Potatoes (2)Winter Wheat/Soybea	ns	· .			75
(3)Corn					75
(1)Potatoes (2)Oats		· · · · · · · · · · · · · · · · · · ·			
(1)Potatoes (2)Sunflowers					
(1)Potatoes (2)Dry Beans				 EO	50
(1)Potatoes (2)Cauliflower	50	50	. 50	50	
(1)Potatoes (2)Cabbage					
Actual Number of Acres in Potatoes	92	92	92	75	50
Return over Variable Costs	\$106,833	\$106,833	\$106,833	\$97,479	\$74,448
Family Labor Activity Level, Hours					50
March (second half)	90	90	90	/4	50
April (first half)	90	. 90	90	74	50
April (second half)	101	101	101	83	01
May (first half)	10	10	10	.8	10
May (second half)	10	10	10	8	10
June (first half)	27	27	27	. 1/	12
June (second half)	44	44	44	52	
July (first half)	64	64	. 64	104	77
July (second half)	84	84	84	/6	116
August (first half)	155	155	155	145	217
August (second half)	217	217	217	217	217
September (first half)	217	217	217	217	217
September (second half)	217	217	217	217	217
October (first half)	217	217	217	217	217
October (second half)	217	217	217	217	217
November (first half)	125	125	125	209	209
November (second half)	32	32	32		
Hand Labor Activity Level Hours	ана се				
August (assord balf)	78	78	78	67	39
August (second half)	148	148	148	130	106
September (IIISt hall)	685	685	685	676	649
September (second half)	660	660	660	664	644
October (second half)	575	575	575	570	553
Other Major Activities			411/ 010	4107 210	200 / LA
Borrow operating capital	\$114,318	\$114,318	5 5114,318	\$107,329	¢¢0,404
Sell rye seed (bu.) Buy rye seed (bu.)	1,192	1,192	1,192	العديدة. يتر 44- عد ج ند	151
Pesticide Active Ingredients				1 000	01.1
Fungicide (1bs. A.I.)	1,492	1,492	1,492	1,238	1 (20
Insecticide (1bs. A.I.)	2,732	2,732	2,732	2,24/	1,632
Herbicide (1bs. A.I.)	588	588	588	523	303

Table 4. Optimal rotations with various limitations on maximum potato acreage (field crop and cole crop rotations) - South Fork Model.



FIGURE I RETURNS ABOVE VARIABLE COSTS FOR VARIOUS POTATO ACREAGE CONSTRAINTS

	N7	h Fork	Sout	h Fork
	Field Crop Rotations	Field & Cole Crop Rotations	Field Crop Rotations	Field & Cole Crop Rotations
Continous Potatoes			NA	NA
(1)Potatoes (2)Potatoes (3)Rye	NA	NA	- -	
(1)Potatoes (2)Rye	\$218.59	\$207.42	\$112.16	\$107.35
(1)Potatoes (2)Corn	295.74	287.31	128.63	126.55
(1)Potatoes (2)Winter Wheat/Soybeans	259.30	250.74	94.73	91.83
(1)Potatoes (2)Winter Wheat/Soybeans (3)Corr	372.15	360.33	238.88	233.06
(1)Potatoes (2)Oats	310.56	299.16	155.39	150.35
(1)Potatoes (2)Sunflow	er 294.15	285.24	135.65	133.10
(1)Potatoes (2)Dry Beau	ns 268.28	259.37	108.73	106.18
(1)Potatoes (2)Caulifi	ower NA		NA	
(1)Potatoes (2)Cabbage	NA	330.74	NA	94.78

Table 5. Reduced income if an acre of various crop rotations was raised, potato acreage not constrained.

unrealistically high.

It is possible that the chemical costs required to produce potatoes will increase. If the Colorado potato beetle becomes more resistant to currently used pesticides, more applications of possibly more expensive chemicals will be needed to maintain potato yields. Table 6 shows the result of increased chemical costs. There are two columns for each group of possible rotations for the North Fork. The left column in each grouping gives the percentage change of chemical costs required for the optimal solution to change. For example, if only field crop rotations were considered as possibilities on the North Fork, the optimal rotation of 150 acres of potatoes (as shown in Table 1) would continue to be optimal until chemical costs increased by 98.8 percent (but returns over variable costs would decrease). If chemical costs increased by 98.8 percent, 138 acres of continuous potatoes should be raised and 12 acres of the potato/rye rotation. Pesticide cost increases of 100 percent would not cause any additional changes in the

The model was more sensitive to pesticide cost changes if both field and cole crop rotations were considered. The optimal solution would change in this situation if chemical costs increased by 93.8 percent on the North Fork. Pesticide cost increases of 100 percent would not change the optimal rotations on the South Fork.

In the past, potato producers have been able to maintain potato yields by using new chemicals or by increasing rates. In the future, however, the Colorado potato beetle may develop high levels of resistance to all registered chemicals. In this situation potato yields might decrease. Table 7 shows the changes in the optimal solution if potato yields decreased when all other factors were held constant. Potato yields must decrease by 27.5 percent on the North Fork and by 33.6 percent on the South Fork before the optimal solution changes if only field crop rotations were considered. A 26.1 percent yield decrease on the North Fork (32.9 percent on the South) was required for the optimal solution to change if both field and cole crop rotations were considered as cropping alternatives.

Table 7 also shows the percentage yield decrease required for the second change in the optimal solution. For example, on the North Fork, if only field crop rotations are considered, the optimal solution changed after a yield decrease of 27.5 percent. At this point 138 acres of continuous potatoes should be raised and 12 acres of the potato/rye rotation. If potato yields decreased by 32.6 percent, the optimal solution would again change. At this point, 67 acres of continuous potatoes, 12 acres of potatoes/rye, and 71 acres of potatoes/double crop (winter wheat, soybeans)

Potato yields may, instead of decreasing in the future, increase in rotations. In this scenario, potato yields were held at current levels for continuous potatoes on the North Fork. They also remained constant for the second year of potatoes in the three year rotation of potatoes/potatoes/rye on the South Fork. The yields may increase in the rotations due to less pest problems if potatoes do not follow potatoes. Table 8 shows the percentages that potato yields in rotations would have to increase for the optimal solution to change. If only field crop rotations were considered

		North	Fork		South	Fork
	7 F L V F A	2	Fiel & Cole	d Crons	Field Crops	Field & Cole Crops
	98.8%	100%	93.8%	100%	100%	100%
		-		acres	1	
Continuous potatoes	138	138	88	88	NA	NA
<pre>(1)Potatoes (2)Potatoes (3)Rye</pre>	NA	NA	NA	NA	150	100
<pre>(1)Potatoes (2)Rye</pre>	12	12	12	12	-	-
<pre>(1)Potatoes (2)Corn</pre>			2		-	*:
<pre>(1)Potatoes (2)Winter Wheat/Soybeans</pre>						
<pre>(1)Potatoes (2)Winter Wheat/Soybeans (3)Corn</pre>	1 1		1	-		-
(1)Potatoes (2)Oats	1			.]		
<pre>(1)Potatoes (2)Sunflowers</pre>			- - - - - - 	2	440 m	
(1)Potatoes (2)Dry Beans	1					
<pre>(1)Potatoes (2)Cauliflower</pre>	NA	NA	20	50	NA	20
<pre>(1)Potatoes (2)Cabbage</pre>	NA	NA			NA	
Return over Variable Costs	\$38 , 998	\$38,342	\$58,968	\$55 , 956	\$55,865	\$74,658

Table 6. Optimal rotations if chemical costs for potatoes increase by various percentages.

15

ı

the product of the second s

Optimal rotations if potato yields decrease by various percentages. Table 7.

1.1

36.8% \$51,247 & Cole Crops 32.9% 36.8% NA Ξ 89 50 i Field \$56,252 NA 16 84 50 South Fork ļ Field Crops 3.6% 38.1% \$35,325 \$29,381 NA 139 Ц NA NA 33.6% 16 NA 134 NA Μ ļ 1 --acres--\$45**,**356 <u>& Cole Crops</u> 6.1% 31.5% 10 NA 12 78 50 ł Field 26.1% \$55**,**659 88 NA 12 North Fork 50 2 Field Crops \$23,794 67 NA 12 71 NA NA ļ 27.5% \$35,511 138 NA 12 NA NA (1)Potatoes (2)Winter Wheat/Soybeans (1)Potatoes (2)Winter Wheat/Soybeans (1)Potatoes (2)Potatoes (3)Rye (1)Potatoes (2)Cauliflower Return over Variable Costs (1)Potatoes (2)Sunflowers (1)Potatoes (2)Dry Beans (1)Potatoes (2)Cabbage Continuous potatoes (1)Potatoes (2)Corn (1)Potatoes (2)Oats (1)Potatoes (2)Rye (3)Corn

the yield would have to increase by 27.5 percent and 33.5 percent on the North and South Forks respectively. (The next optimal solution changes would be at 32.6 percent for the North Fork and 38.0 percent for the South Fork.) It is probably unrealistic to expect potato yields to increase this much due to the benefits of rotation.

The optimal solution was less sensitive to potato yield increases if both field and cole crop rotations were considered in the model (Table 8). In this situation, if potato yields increased by 26.1 percent on the North Fork and 32.8 percent on the South Fork, the optimal solution would change. Potato yield increases of about 32 percent would be associated with large increases in potatoes grown in rotations.

So far the results of the model have been discussed for field crop rotations and for field and cole crop rotations. A third model variation considered field and cole crops as well as two additional rotations. These rotations were as follows: (1) potatoes followed by a year of onions, and (2) potatoes followed by a double crop of spinach and soybeans in the second year. Both of these rotations have agronomic problems due to the higher pH soils required for the onion and spinach production. If potato varieties could be developed that were resistant to scab in higher pH soil, these rotations may have potential.

Table 9 shows that the maximum acreage of 50 acres of both the potato/ onion and the potato/double crop (spinach, soybean) rotations would be raised on each Fork. (No more than 25 acres of any one vegetable crop was permitted in the model due to large possible price fluctuations.) The return over variable costs was higher than the returns have been with the traditional rotations on both Forks. The yield on vegetable crops was assumed to be the same on both Forks, but irrigation equipment had to be purchased for production of vegetables on the South Fork. Although returns above variable costs were approximately the same for vegetable crops on both Forks, total returns above variable costs were higher on the South Fork due to higher relative returns for potatoes on the 75 acres grown in the optimal solution on both Forks.

Two major problems must be solved before these two rotations will have potential on Long Island potato farms. First, an acceptable scab-resistant potato variety for relatively high pH soils must be developed. Second, growers must be able to handle large amounts of seasonal labor to successfully raise rotations with onions or spinach.

SUMMARY AND CONCLUSIONS

The current potato production practices on Long Island are the most profitable of the field crop rotations considered. On the North Fork, continuous potato production gave the highest return over variable costs. On the South Fork, two years of potatoes followed by a year of rye (a common current practice) gave the highest returns. There are many problems with intensive potato production. Researchers are constantly investigating new pesticides to stay ahead of insect resistance build-up. Heavy use of some alternative chemicals may result in ground water contamination similar to the problems caused by aldicarb. Optimal rotations if potato yields in rotations increase by various percentages. Table 8.

36.6% \$101,089 \$101,583 \$117,887 \$120,580 \$119,759 \$125,288 \$138,974 \$143,629 & Cole Crops NA Ц 89 ļ ļ 50 Field 32.8% NA 16 84 South Fork ļ 50 ļ Field Crops NA 139 Π NA NA 33.5% MA 16 134 NA MA ł --acres--& Cole Crops 74 1% 31.6% 10 NA 12 78 30 Field 26.1% 88 MA 12 North Fork 20. ł Field Crops 67 MA 27 7 NA M 27.5% 138 MA 2 NA NA (1)Potatoes (2)Winter Wheat/Soybeans (1)Potatoes (2)Winter Wheat/Soybeans (1)Potatoes (2)Potatoes (3)Rye (1)Potatoes (2)Cauliflower Return over Variable Costs (1)Potatoes (2)Sunflowers (1)Potatoes (2)Dry Beans (1)Potatoes (2)Cabbage Continuous potatoes (1)Potatoes (2)Corn (1) Potatoes (2) Oats (1)Potatoes (2)Rye (3)Corn

		North Fork	 South Fork
Rotations, acres			
Continous potatoes			NA
(1)Potatoes (2)Rye		NA	
(1)Potatoes (2)Corn			
(1)Potatoes (2)Winter Wheat/Soybeans			
(1)Potatoes (2)Winter Wheat/Soybeans	(3)Corn		
(1)Potatoes (2)Oats			
(1)Potatoes (2)Sunflowers			· · · · · · · · · · · · · · · · · · ·
(1)Potatoes (2)Dry Beans			
(1)Potatoes (2)Cauliflower		50	50
(1)Potatoes (2)Cabbage			
(1)Potatoes (2)Onions		50	50
(1)Potatoes (2)Spinach/Soybeans		50	50
Actual Number of Acres in Potatoes		75	75
Return over Variable Costs		\$120,287	\$132,683
Family Labor Activity Leven, Hours			150
March (second half)	i.	85	150
April (first half)		217	84
April (second half)		217	217
May (first half)		217	217
May (second half)		217	217
June (first half)		217	217
June (second half)		217	154
July (first half)		- 217	200
July (second half)		217	217
August (first half)		217	217
August (second half)		217	217
Sentember (first half)		142	217
September (second half)		217	217
October (first half)		217	217
October (in an indif)		217	217
Nevember (first balf)		209	209
November (record balf)	1	53	53
November (second nativ			
Hired Labor Activity Level, nours			
April (first half)		81	· /1
April (second half)		69	F1
May (first half)		494	502
May (second half)		562	202
June (first half)		568	515
June (second half)		4	
July (first half)		46	
July (second half)		674	681
August (first half)		704	125
August (second half)		846	862
September (first half)		134	134
September (second half)		667	671
October (first half)		655	656
October (second half)		570	571
Other Major Activities			
Pernow Operating Capital		\$151,455	\$132,683
Buy Rye Seed, bushels		225	225
Pesticide Active Ingredients		4 0.07	1 /50
Fungicide, Ibs. A.I.		1,222	. 1,400
Insecticide, Ibs. A.I.		2,001	 2,001
Herbicide, Ibs. A.I.		625	020

Table 9. Optimal rotations for the North and South Fork models (all rotations).

A variety of field crop rotations could be raised on Long Island, but all would result in lower returns over variable costs than traditional cropping practices. If potato acreage was limited, the two most economically feasible field crop rotations on both Forks are: (1) A year of potatoes followed by a year of rye, and (2) a year of potatoes followed the next year by a double crop of winter wheat and soybeans. The potato/rye rotation would be raised on only a few acres of land to provide seed for the rye cover crop. The potato/double crop (winter wheat, soybean) rotation is the most feasible replacement for large amounts of potato acreage. But, potato growers are unlikely to raise these rotations unless there is legislation forcing them to raise less potatoes or they are subsidized for the resulting income loss.

When cole crop rotations were also considered in the linear programming model, returns were increased. Returns over variable costs were higher (six percent for the North Fork, 16 percent for the South Fork) for the optimal plan with nonrestricted potato acreage if 25 acres of the farm was planted in a potato/cauliflower rotation. Cauliflower tolerates low soil pH. Relatively large quantities of seasonal labor must be hired if more than a few acres of cauliflower are raised. If a substantial number of growers grew 25 acres of cauliflower, however, the price of cauliflower may be significantly reduced, an event that cannot be handled by the farm-level models constructed for this research.

The development of insect resistance to pesticides on Long Island potato fields has caused many problems. In the past, growers have been able to cope by using new and/or heavier applications of insecticides. In the future potato production costs or potato yields (and thus returns) might change due to insect resistance to available chemicals. It might become more expensive to control the Colorado potato beetle. A second scenario is that potato yields might decrease due to the insect problems, or conversely, potato yields in rotations might increase since there would be less pressure from potato pests if potatoes were raised less intensively. The optimal solution to the linear programming model is not very sensitive to these changes. The traditional rotations of continuous potatoes on the North Fork and two years of potatoes followed by a year of rye on the South remain the most economically feasible unless extreme changes would occur in potato returns or production costs.

The highest returns in the linear programming model resulted when some vegetable crop rotations with agronomic problems were considered. Potatoes/onions and potatoes/double crop (spinach, soybeans) had significantly higher returns over variable costs than the traditional potato cropping pattern. Spinach and onions require higher pH soils than potatoes. If potato varieties could be developed which are resistant to scab, these rotations become possible alternatives.

Markets for some of the crop alternatives discussed in this report would need to be developed. (Cauliflower has the advantage of having an already well developed market.) However, the marketing problems could perhaps be solved if substantial acreage of these crops were raised on Long Island.

Labor is a problem in the vegetable crop rotations. Many of the potato growers have little experience managing seasonal labor and do not want to

have to deal with the extra management required to utilize migrant labor. But, some vegetable growers on Long Island do use migrant labor, so perhaps potato farmers should consider vegetable crop rotations as a possible alternative.

The results of this research suggest other rotations or markets which will be analyzed in the future. Some of these are as follows:

1) A potato-potato-corn rotation with a rye cover crop as an alternative to the traditional rotation of potato-potato-rye on the South Fork.

na da antiga da na mangazi da na su

 Markets for oats and straw to horse owners. Some relatively high prices for oats on Long Island have been reported, but the market is perhaps limited.

These analyses may show a greater potential for field crops in mitigating losses from rotations.

Crop rotation has the potential of being used on Long Island, along with other IPM practices, to help solve some of the potato pest problems. The rotations must be carefully chosen to avoid significant losses in farm income.

REFERENCES

- Apple, J.L., P.S. Benepal, R. Berger, G.W. Bird, F. Maxwell, W.G. Ruesink, P. Santelmann, and G.B. White. Integrated Pest Management: A Program of Research for the State Agricultural Experiment Stations and the <u>College of 1890.</u> Committee Report, The Intersociety Consortium for Plant Protection for the Experiment Station Committee on Organization and Policy. Raleigh, N.C. 1979.
- Benson, F.J. "1974 Maine Farm Planning Guide", Bulletin 584, Cooperative Extension Service, University of Maine, May 1974.
- Dhillon, P.S. "Cost of Producing Selected Fresh Market Vegetables in South Jersey", Bulletin B-853, Department of Agricultural Economics and Marketing, Rutgers University, August 1979.
- Knoblauch, W.A. "Field Crop Enterprise Budgets: Cost and Returns for 1981", mimeo, Department of Agricultural Economics, Cornell University, 1981.
- Knoblauch, W.A., R.A. Milligan, R.J. Haslem, and M.M. vanLieshout "An Economic Analysis of New York Field Crop Enterprises", A.E. Res. 80-6, Department of Agricultural Economics, Cornell University, April 1980.

Lazarus, W.F. Personal Communication, October 1982.

- New York Crop Reporting Service. <u>New York Agricultural Statistics</u>. Albany, NY: New York State Department of Agriculture and Markets, various issues.
- New York State College of Agriculture and Life Sciences. <u>Cornell Recommen-</u> <u>dations for Commercial Vegetable Production - 1982.</u> Ithaca, NY:

New York State College of Agriculture and Life Sciences. <u>1982 Cornell</u> <u>Recommends for Field Crops.</u> Ithaca, NY: Cornell University, 1981.

- Phelps, J.B. and R.B. How. "Planning Data for Small Scale Commercial Vegetable and Strawberry Production in New York", A.E. Res. 81-20, Department of Agricultural Economics, Cornell University, October 1981.
- Putnam, J.N. <u>Aroostook County, Maine Potato Industry Study</u>. Springfield, MA: Farm Credit Banks, 1981.

Snyder, D.P. "Cost of Production Update for 1980", A.E. Res. 81-11, Department of Agricultural Economics, Cornell University, July 1981.

APPENDIX A

Budget
Potato
A-1:
Table

. 1				North Continuous	Fork ^a	South F	ork
	Unit	Price	Outstite	Potatoes	Rotation	becond Year of Potatoes	Potato Rotation
Receipts:			Auanlilly		Total		
North Fork -							
90% size A, U.S. No. 1 10% culls and size B	cwt. cwt.	\$5.30 2.50	286 31	\$1,515.80 77.50	\$1,515.80 77 50		·
South Fork -				2 3			
90% size A, U.S. No. 1 10% culls and size B	cwt. cwt.	5.30 2.50	301 33			\$1,595.30 82 50	\$1,595.30
Total Receipts				\$1.593.30	¢1 503 30	00.20	82.50
Expenses:						ş1,6//.80	\$ 1,6 77.80
Seed Fertilizer - Nitrogen Phospherous Potassium	1b. 1b. 1b.	.073 .32 .28	2,130 N.F175;S.F175 N.F300;S.F350 N.F175;S.F175	\$ 155.49 56.00 84.00 24.50	\$ 155.49 56.00 84.00 24.50	\$ 155.49 56.00 98.00	\$ 155.49 56.00 98.00
Chemicals - Fungicide Insecticide				38.59 356.00	38.59 33.50	46.19 91.04	24.50 46.19
Herbicide Machinery Variable Cost				23.98 96.95	23.98 94.49	312.75 23.98 61.00	262.88 23.98 58 54
Selected Variable Cost				\$ 835.51	\$ 809.55	16.777 \$	\$ 725.58
Return over Selected Variable Costs				\$ 757.79	\$ 783.75	\$ 899.89	\$ 952.22

a irrigated

i

Table A-2: Rye Budget

				North Fork	South Fork	Cover Crop Only
	Unit_	Price	Quantity		Total	
Receipts:						
North Fork	bu.	\$2.80	34	\$95.20		
South Fork	bu.	2.80	38		\$106.40	
Cover Crop	bu.	2.80	. 0			\$0.00
Expenses:						
Seed	bu.	(a)	1.5	(a)	(a)	(a)
Fertilizer - Nitrogen	1b.	. 32	20	\$ 6.40	\$ 6.40	\$ 0.00
Chemicals - Herbicide				.70	.70	0.00
Custom Combine				25.00	25.00	0.00
Machinery Variable Cost				5.69	5.69	2.73
Selected Variable Costs				\$37 . 79	\$37.79	\$ 2.73
· · · · · · · · · · · · · · · · · · ·						· ·
Return over Selected Variable Costs	· .	:		\$57.41	\$68.61	\$-2.73

(a)Seed expense was calculated in the linear programming model instead of the budget, since the farmer had the option of buying seed for \$5.00 per bushel or raising his own (valued at \$2.80 per bushel).

Table	A-3:	Corn	Budget
-------	------	-----------------------	--------

	· .				North Fork ^a	South Fork
· · · · · · · · · · · · · · · · · · ·		Unit	Price	Quantity	Tot	al
Receipts:	· .					
North Fork South Fork		bu. bu.	\$2.75 2.75	102^{b} 102^{b}	\$280.50	\$280.50
						I
Expenses:	·					
Seed Fertilizer - Lime Chemicals - : I Custom Machin Machinery Var	Nitrogen Phospherous Potassium Insecticide Herbicide hery - Planting Combining Drying tiable Cost	25,000 seeds 1b. 1b. 1b. ton	17.20 .32 .28 .14 28.00	1 100 50 50 .5		\$ 17.20 32.00 14.00 7.00 14.00 .46 12.65 5.00 40.00 30.00 11.70
Selected V	Variable Costs				\$207.77	\$184.01
Return ove Variable C	r Selected osts	a status Alta a			\$ 72.73	\$ 96.49

T

^a irrigated

^b dry, shelled corn

	A 1	1.1.1 m to out	Uboat	Budget
Table	A-4:	Winter	wnear	budget

				North Fork	South Fork
	Unit	Price_	Quantity	Tot	tal
Receipts:					
North Fork South Fork	bu. bu.	\$3.25 3.25	46 51	\$148.50	\$165.75
Expenses:					·
Seed Fertilizer - Nitrogen Phospherous Potassium Lime Chemicals - Herbicide Custom Combine Machinery Variable Cost Selected Variable Costs	bu. 1b. 1b. 1b. ton	8.70 .32 .28 .16 28.00	3 30 30 .25	\$ 26.10 9.60 8.40 4.80 7.00 .70 25.00 5.69 \$ 87.29	\$ 26.10 9.60 8.40 4.80 7.00 .70 25.00 5.69 \$ 87.29
Return over Selected Variable Costs				\$ 62.21	\$ 78.46

And an and the second second second

				North Fork	South Fork
	Unit	Price	Quantity	Toi	
Receipts:					
North Fork	bu.	¢ 6 10	àc		
South Fork	5u.	\$ 0.10	26	\$158.60	
	Du.	0.10	29	· · · · · · · · · · · · · · · · · · ·	\$176.90
Expenses:					
Seed	bu	14 40	1. 0		
Fertilizer - Nitrogen	15	14 40	1.2	\$ 17.30	\$ 17.30
Phospherous	15.	• 32	10	3.20	3.20
Potassium	15.	• 40° 16°	40	11.20	11.20
Lime	ton.	28.00	40 0 E	6.40	6.40
Chemicals - Herbicide	con	20,00	• 25	7.00	7.00
Custom Combine				7.95	7.95
Machinery Variable Cost				25.00	25.00
Selected Variable Costs				11.59	11.59
:			·	\$ 89.64	\$ 89.64
Return over Selected					
variable Costs				\$ 68,96	\$ 87.26

Table A-5: Soybean Budget

Table	A-6:	0ats	Budget	
-------	------	------	--------	--

				North Fork	South Fork
	Unit	Price	Quantity	То	tal
Receipts:					
North Fork South Fork	bu. bu.	\$1.70 1.70	65 72	\$110.50	\$122.40
Expenses:					
Seed Fertilizer - Nitrogen Phospherous Potassium Lime Chemicals - Herbicide Custom Combine Machinery Variable Cost Selected Variable Costs	bu. 1b. 1b. 1b. ton	5.50 .32 .28 .16 28.00	3 40 35 35 .25	\$ 16.50 12.80 9.80 5.60 7.00 .70 25.00 9.77 \$ 87.17	\$ 16.50 12.80 9.80 5.60 7.00 .70 25.00 9.77 \$ 87.17
Return over Selected Variable Costs				\$ 23.33	\$ 35.23

				North Fork	South Fork
	Unit	Price	Quantity	То	tal
Receipts:					
North Fork	014	610 40	1.0		
South Fork	CWL +	310.40	10	Ş166.40	1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 -
	CWE.	10.40	18		\$187.20
Expenses:					
					f(x) = f(x) + f(x)
Seed	15.	1.32	6	¢ 7 07	à 7 00
Fertilizer - Nitrogen	1b.	. 32	60	9 / 192 j	\$ 7.92
Phospherous	1b.	.28	20	19.20	19.20
Potassium	15.	.16	20	3 20	5.60
Lime	ton	28.00	.25	7.00	3.20
Chemicals - Herbicide			• 2.5	14 65	
Custom Machinery - Combine	2			25 00	14.65
Drying				12 / 9	- 25.00
Helicor	oter	1		12.40	14.00
sprayi	ng		÷	5.00	5 00
Machinery Variable Cost				9.04	9.06
Selected Variable Costs	•			\$109.09	\$110.61
Potumo orașe din 1					
Recurn over Selected	· .				
variable Costs	· .			\$ 57.31	\$ 76.59
			· .		

Table A-7:	Sunflower	Budget
------------	-----------	--------

				North Fork	South Fork
	Unit	Price	Quantity	To	tal
Receipts:					
North Fork	cwt.	\$21.60	13	\$280.80	
South Fork	cwt.	21.60	14		\$302.40
	-				• ,
Expenses:					
Seed	1b.	.50	90	\$ 45.00	\$ 45.00
Fertilizer - Nitrogen	1b.	.32	25	8.00	8.00
Phospherous	1b.	.28	75	21.00	21.00
Potassium	1b.	.14	50	7.00	7.00
Lime	ton	28.00	.5	14.00	14.00
Chemicals - Fungicide				.46	.46
Insecticide				.23	• 23
Herbicide				28.85	28.85
Custom Combine				30.00	30.00
Machinery Variable Cost				12.90	12.90
Selected Variable Costs				\$167.44	\$167.44
Return over Selected					
Variable Costs				\$113.36	\$134.96

Table A-8: Dry Bean (Red Kidney) Budget

				North Fork ^a	South Fork ^a
	Unit	Price	Quantity	To	otal
Receipts:	cwt.	\$19 .3 0	150	\$2,895.00	\$2,895.00
Expenses:	.÷	• •			
Plants Fortilizer Nu	1,000	26.40	10	\$ 264.00	\$ 264.00
Phospherous	1b. 1b.	• 32 • 28	160 320	51.20 89.60	51,20
Potassium Lime (hydrated)	1b. ton	•14 122	160	22.40	22.40
Chemicals - Insecticide Herbicide			• •	102.96	102.96
Fungicide				10.60	10.60 19.25
Machinery Variable Cost		1,45	429	622.05 73.71	622.05 78-89
Selected Variable Costs				\$1,316.77	\$1,321.95
Return over Selected Variable Costs	· · ·			\$1,578.23	\$1.573.05
		· .	1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -		

Table A-9: Cauliflower Budget

a irrigated

Table A-10: Cabbage Budget

				North Fork ^a	South Fork ^a
	Unit.	Price	Quantity	To	otal
Receipts:	cwt.	\$8.20	257	\$2,107.40	\$2,107.40
Evnenses:					
Plants Fertilizer - Nitrogen Phospherous Potassium Lime (hydrated) Chemicals - Fungicide Insecticide Herbicide	100 1b. 1b. 1b. ton	1.40 .32 .28 .14 122	150 150 100 100 .50	$ \begin{array}{c} $ 210.00 \\ 48.00 \\ 28.00 \\ 14.00 \\ 61.00 \\ 40.92 \\ 75.51 \\ 30.15 \\ (16.00) \\ 40.92 \\ 75.51 \\ 30.15 \\ (16.00) \\ 40.92 \\ 75.51 \\ 30.15 \\ (16.00) \\ 40.92 \\ 75.51 \\ 30.15 \\ (16.00) \\ 40.92 \\ 75.51 \\ 30.15 \\ (16.00) \\ 40.92 \\ 75.51 \\ 30.15 \\ (16.00) \\ 40.92 \\ 75.51 \\ 30.15 \\ (16.00)$	\$ 210.00 48.00 28.00 14.00 61.00 40.92 75.51 30.15 616.80
Crates Machinery Variable Cost Selected Variable Costs	•	1.20	514	616.80 83.88 \$1,208.26	\$1,213.44
Return over Selected Variable Costs				\$ 899.14	\$ 893.96

a irrigated

1			
Table	A-11:	Onion	Budget

		· · · · · · · · · · · · · · · · · · ·		· · · .	
				North Fork ^a	South Fork ^a
	Unit	Price	Quantity	Tc	tal
Receipts:	cwt.	\$ 9.65	175	\$1,688.75	\$1,688.75
Expenses:					
Seed Fertilizer - Nitrogen Phospherous Potassium Lime Chemicals - Fungicide Insecticide Herbicide	1b. 1b. 1b. 1b. ton	15.00 .32 .28 .14 28.00	2.5 100 100 100 1	\$ 37.50 32.00 28.00 14.00 28.00 13.86 2.39	\$ 37.50 32.00 28.00 14.00 28.00 13.86 2.39
Bags Machinery Variable Cost Selected Variable Costs		.35	350	23.20 122.50 53.61 \$ 355.06	23.20 122.50 57.06 \$ 358.51
Return over Selected Variable Costs				\$1,333.69	\$1,330.24

^a irrigated

.

Table A-12: Spinach Budge	ble A	-12:	Spinach	Budget
---------------------------	-------	------	---------	--------

	-			North Fork ^a	South Fork ^a
	Unit	Price	Quantity	To	otal
Receipts:	cwt.	\$22.90	80	\$1,832.00	\$1,832.00
Expenses:			,		
Seed	16.	25.00	5	\$ 125.00	\$ 125.00
Fertilizer - Nitrogen	16.	.32	150	48.00	48.00
Phospherous	16.	.28	100	28.00	28.00
Potassium	16.	.14	100	14.00	14.00
Lime	ton	28.00	.5	14.00	14.00
Chemicals - Fungicide			· · ·	3.06	3.06
Insecticide				12.59	12.59
Packing Boxes		. 85	160	136.00	136.00
Cooling		.30	160	48.00	48.00
Machinery Variable Cost				50.61	54.06
Selected Variable Costs	3			\$ 479.26	\$ 482.71
Return over Selected				· · · ·	
Variable Costs				\$1,352.74	\$1,349.29

a irrigated

Machine	New Cost	Speed (mph)	Field Efficiency
Tractor 60 hp	\$17,900		
Tractor 100 hp	36,300		
Rollover plow with Clodbuster, 4-16" bottoms	9,500	4.0	. 8
Sprayer, 48' boom	13,500	4.5	• 5
Potato cultivator, 4 row	2,400	4.0	. 8
Potato planter, 4 row	15,000	4.0	• 65
2 big gun irrigation sets (80A)	44,000	·	
Disk harrow, 13'	4,950	5.0	•8
Potato harvester, 2 row	31,000	2.0	•6
3 bulk bodies, 18'	13,500		
Seed cutter	4,000		
Grain drill, 18 x 7	5,100	4.0	.7
Precision seeder, 4 row	2,900	2.0	•8
Transplanter, 4 row	2,400	1.0	.7
2 wagons	5,000		

Table A-13: Machinery Complement for Potato Production

	Marc	÷	Anr		Ma	~	- Inc	-	lut		Augu	Ist	Septem	ber	Octof	er	Nover	her
- b								}	 		,		-					
	-	2	•	2	•	2	1	2	-	2	-	2		2	-	2	-	14
										hours								
Potatoes:							•••••											
North Fork - Continuous		86.	.98		-10	•10	.42	1.33	1-11	1.11	1.95	1.95	•84	-84	-84	-84		
Rotation		96.	86.		.10	.10	.22	1.33	1.11	1.11	1.95	1.95	•84	-84	•84	•84		
South Fork - 2nd year		96	-98	1.1	.10	.10	•42	.42	•20	.20	1.03	1.03	-84	-84	.84	.84	_	
Rotation		96	-98		.10	.10	.22	.42	.20	-20	1.03	1.03	-84	•84	.84	-84		
Rye									-17	.17			.14	.14	-14	.14		
Rye Cover Crop		 - - -											.12	.12	.12	.12		
Soybeans									1.07		.14	.14			•17	.17		
Winter Wheat								.34						•00	•29	•20		
Corn - North Fork				.22	.42	.20	•03		-16	16.	-91	-91	_		•26	•26		
South Fork				.22	.42	•20	•03					_			-26	•26		
Oats		.22	.42	.20						11.	.11							
Dry Beans (Red Kldney)					.14	.46	-46	.18	.18	•03			.17	-17				
Sunflowers				.14	-24	•24								17	.17			
Cauliflower - North Fork								.22	1.6	2.29	2.2	7.89	10.9	32.24	31.33	27.94	8.34	2.12
South Fork								.22	1.6	2.4	2.42	8.0	11.01	32.35	31.33	27.94	8.34	2.12
Cabbage - North Fork							1.5	2.5	1.29	7.98	7.98	1.38	38.78	33.53	33.44			
- South Fork							1.5	2•62	1.4	8.08	8.08	1.49	39.89	33.53	33.44			
Onions - North Fork			•22	•22	.37	3.67	4.59	4.59	4.5	32.81	31.9	31.9						
- South Fork			•22	.22	•37	3 67	4.7	4.7	4.61	32.92	31.9	31.9	_					
Splnach - North Fork		.44	8.23	7.89	27.75	27.18	26.18											
South Fork		•44	8.35	0*8.	28.08	27.29	26.18											
									•									

Table A-14: Distribution of Annual Labor Needs by Semi-Monthly Time Periods