# IMPACT OF HYPOXIC NITRATE RESPIRATION ON MYCOBACTERIUM TUBERCULOSIS

A Dissertation Presented to the Faculty of the Graduate School of Cornell University In Partial Fulfillment of the Requirements for the Degree of Doctor of Philosophy

> by Amy Christine Cunningham-Bussel January 2014

© 2014 Amy Christine Cunningham-Bussel

# IMPACT OF HYPOXIC NITRATE RESPIRATION ON MYCOBACTERIUM TUBERCULOSIS

# Amy Christine Cunningham-Bussel, Ph.D. Cornell University 2014

An estimated one third of the world's population is infected with Mycobacterium tuberculosis (Mtb) and suppresses the pathogen's replication without eradicating it, thereby remaining at risk for developing active tuberculosis (TB). It is not clear how Mtb survives, potentially for decades, in a hostile host environment. Respiration of nitrate to nitrite could help Mtb to survive in hypoxic tissues, but was not thought to be significant at physiologic oxygen tensions, nor was the resultant nitrite considered consequential to Mtb's physiology. Here, we found that Mtb infecting human macrophages in vitro produces copious nitrite at physiologic oxygen tensions. The nitrite impacts Mtb in diverse ways, slowing its growth, reducing its consumption of ATP, regulating its survival following treatment with isoniazid and hydrogen peroxide, and remodeling its transcriptome in a manner distinct from that of nitric oxide. Thus, respiration of nitrate and adaptation to nitrite are likely to play a prominent role in Mtb's pathophysiology, whether or not the Mtb resides in hypoxic sites.

## **BIOGRAPHICAL SKETCH**

Amy Christine Cunningham- Bussel graduated *magna cum laude* in biological sciences from Cornell University in 2005 where she majored in neurobiology and behavior. She is a student in the Tri-Institutional MD/PhD program of Weill Cornell Medical College, Rockefeller University and The Sloan-Kettering Institute.

She joined the laboratory of Dr. Carl Nathan in September of 2009 where her research focused on studying the impact of hypoxic nitrate respiration on *Mycobacterium tuberculosis*.

### ACKNOWLEDGMENTS

I first want to thank my mentor Dr. Carl Nathan. I have always appreciated the incredible biologic narrative that unfolds within scientific articles that eventually become integrated into textbooks. My work over the past four years has been my first real opportunity to contribute to that ever-expanding body of knowledge that first captivated me in college. I am thankful to Carl for entertaining my naïve desire to work with primary human, rather than murine, macrophages and for giving me complete freedom to explore a fascinating area of biology, spanning the interaction between the human host and *Mycobacterium tuberculosis*. This particular project required that we travel an incredibly winding road together. Sometimes, even many times, the road was not particularly apparent. Despite this, he never failed to provide encouragement and, when needed, motivate forward motion. Over the years, this project has become exactly the intellectual challenge that I hoped for when I first joined the Nathan laboratory. I am exceedingly grateful to Carl for his mentorship during my graduate training.

I would also particularly like to thank my thesis committee members, Dr. Sabine Ehrt, Dr. Dirk Schnappinger, Dr. Daniel Mucida, Dr. Lionel Ivashkiv and the late Dr. Ralph Steinman, who all graciously sacrificed long hours to meet and provide me with valuable scientific and career advice.

I am privileged and honored to be a member of the Tri-Institutional

MD/PhD program, which I owe in large part to three individuals, Dr. Olaf Andersen, Dr. David Silbersweig and Dr. Emily Stern. I met Olaf nearly 10 years ago when I first told him of my interest in the MD/PhD program. He has contributed to every academic step that I have taken ever since. He is an incredible program director and I am forever grateful to him for providing me with guidance and ceaseless encouragement over the past decade.

Beginning in the summer of 2005, Dr. David Silbersweig and Dr. Emily Stern, kindly offered me a position in the Functional Neuroimaging Laboratory. Over the next two years in addition to being able to participate in an incredible area of scientific research they offered me the opportunity to contribute to a project that became my first, first authored manuscript. They are two of the kindest, most encouraging scientists and clinicians that I know and I am truly thankful to them for supporting my application to the MD/PhD program.

I would like to thank friends in the Nathan, Ehrt, Rhee, Ding and Schnappinger Labs, especially, Poonam Rath, Jennifer Small, Tamu Chidawanyika, Christina Maksymiuk, Kristin Burns, Julia Roberts, Carolina Trujillo, Joeli Marrero, German Rehren, Kathryn O'brien, Jee Kim, Julien Vaubourgeix, Tania Lupoli, Selin Somersan Karakaya, Madhu Nandakumar, Nikki Goodsmith, Michele Fuortes, Heran Darwin, Ben Gold, Ruslana Bryk, Thulasi Warrier, Sumit Chakraborty, Anand Balakrishnan, Gang Lin, Kyu Rhee, Aihao Ding, Xiuju Jiang, Yan Ling. I am thankful for your friendship, encouragement, fascinating scientific discussions, technical advice and assistance, and very especially, for those who worked in the Bronx BSL3, for fun conversations that got me through hours of CFU assays. I am very fortunate to have worked with all of you.

I would also like to acknowledge everyone who donated blood. I am extremely grateful for your contribution.

Through my parents, I was introduced to science and medicine from an early age and was witness to their love of their own careers. It has been a joy to share in their scientific endeavors and to tell them about my own. I am especially thankful to them for all of the opportunities they have unstintingly provided over the years, for their endless love and support through stressful times and for planning many fun vacations and activities that helped me escape the stresses of PhD work.

I first met Tyler Janovitz when I interviewed for the MD/PhD program in 2006 and we were married four years later. I would especially like to thank him for all of the late night discussions and advice that he gave me long after his interest had waned over the course of my graduate training. I also thank him for all of the fun that we have had, which never failed to draw me out of the experiment-has-failed blues.

## **TABLE OF CONTENTS**

BIOGRAPHICAL SKETCHiii
ACKNOWLEDGMENTSiv
<b>CHAPTER 1</b> 1
INTRODUCTION1
The toxic molecules produced by an active immune response1
The host exploits the toxicity and essentiality of transition metals2
The availability of molecular oxygen regulates mycobacterial growth4
Oxygen regulates the growth of Mtb within the host
When oxygen is scarce, Mtb respires the alternative electron acceptor
nitrate7
Mtb respires nitrate within primary human macrophages7
Conclusions
References11
<b>CHAPTER 2</b>
Summary
Introduction
<b>Results</b>
Discussion41
Methods
References
<b>CHAPTER 3</b>
Summary
Introduction
Results
Discussion71
Methods74
References
<b>CHAPTER 4</b>
<i>Future Directions</i>
References
PUBLICATIONS

# LIST OF FIGURES

## CHAPTER 2

Figure 1	
Figure 2	
Figure 3	
Figure 4	
Figure 5	
Figure 6	
Supplementary Figure 1	
Supplementary Figure 2	
Supplementary Figure 3	

# CHAPTER 3

Figure 1	61
Figure 2	
Figure 3	
Figure 4	
Supplementary Figure 1	
Supplementary Figure 2	105

## LIST OF TABLES

## CHAPTER 2

Table 1	
Supplementary Table 1	
Supplementary Table 2	
Supplementary Table 3	

# APPENDICES

## APPENDIX 1

Supplementary Figure 1	
Supplementary Figure 2	
Supplementary Figure 3	
Supplementary Table 1	
Supplementary Table 2	
Supplementary Table 3	

## APPENDIX 2

Supplementary	Figure	<b>1</b> 104
Supplementary	Figure	<b>2</b> 105

## LIST OF ABBREVIATIONS

CFU: Colony forming unit ETH: Ethionamide IFNγ: Interferon gamma INH: Isoniazid MCSF: Macrophage colony-stimulating factor Mtb: Mycobacterium tuberculosis NOS: Nitric oxide synthase OD: Optical density PIMO: Pimonidazole RNS: Reactive nitrogen species ROS: Reactive oxygen species TB: Tuberculosis TNFα: Tumor necrosis factor

# CHAPTER 1 INTRODUCTION

*Mycobacterium tuberculosis* (Mtb) is the second leading cause of death worldwide from a single infectious pathogen (1). Despite the induction of a potent immune response, Mtb is not eradicated and survives to establish lifelong disease. Furthermore, months of multidrug chemotherapy are required to treat patients with tuberculosis (TB) (2). How Mtb survives the arsenal of toxic small molecules that are products of an active immune response, and why months of multidrug chemotherapy are required to treat TB, remain active areas of investigation.

#### The toxic molecules produced by an active immune response.

Humans have co-evolved with Mtb over millennia and are its only known natural host (3, 4). Infection with Mtb begins with the inhalation of droplet nuclei containing Mtb, which arise from infected individuals and become airborne (5, 6). Within the lung alveoli of a newly infected host, Mtb is confined within the phagosomes of resident macrophages. Once activated by interferon gamma (IFN $\gamma$ ), these phagosomal compartments fuse with adjacent intracellular lysosomes and acidify (7). IFN $\gamma$  also triggers the assembly and recruitment of enzymes responsible for the production of reactive nitrogen and oxygen species (RNS and ROS, respectively). Examples of RNS include, nitric oxide ('NO), nitrogen dioxide ('NO<sub>2</sub>) and nitrite (NO<sub>2</sub><sup>-</sup>) and examples of ROS are superoxide (O<sub>2</sub><sup>-</sup>), hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and hydroxyl radical ('OH) (8, 9). Nitrate (NO<sub>3</sub><sup>-</sup>) can give rise to the above RNS within humans, however, this primarily requires enzymatic reduction by nitrate reductase enzymes expressed by oral commensal bacteria (10). This is described in greater detail below.

Despite exposure to these potentially lethal host products, Mtb expresses enzymes that detoxify ROS and RNS directly, as well as enzymes that repair mycobacterial proteins and DNA damaged by these toxic molecules. For example, enzymes which degrade ROS and/or RNS include Mtb's catalase/peroxidase KatG, the superoxide dismutases SodA and SodC, an NADPH dependent peroxidase/peroxynitrite reductase encoded by *ahpC*, *ahpD*, *lpdC* and *dlaT* and the truncated hemoglobins trHbO and trHbN. The majority of these proteins have been shown to contribute to mycobacterial virulence (9, 11, 12). Methionine sulphoxide reductases, MsrA and MsrB, repair the ROS and RNS mediated oxidation of methionine residues and UvrB, a DNA repair protein contributes to resistance against both ROI and RNI *in vivo* (13, 14). Furthermore, mycobacteria synthesize mycothiol rather than glutathione, which serves as a dominant intracellular antioxidant. Mycobacteria unable to synthesize mycothiol are less able to survive treatment with ROS and RNS (15).

### The host exploits the toxicity and essentiality of transition metals

Every living organism depends upon transition metals for growth, DNA replication and metabolic energy production. Moreover, these essential metals are required for the catalytic function of many of the enzymes involved in resistance to ROS and RNS mediated oxidative stress (16). Iron, the most abundant transition metal within human tissue, is actively sequestered away from microbial pathogens by host proteins. Infected human phagocytes stimulated with IFN $\gamma$  repressed the expression of transferrin and the intracellular iron storage protein, ferritin thus reducing the total concentration of intracellular iron and limiting intracellular bacterial replication (17, 18). The host encodes natural resistance- associated macrophage protein 1 (NRAMP1), which transports iron and manganese out of the phagosomal compartment. Though controversial, mutations in NRAMP1 have been demonstrated to confer resistance to tuberculosis in mice and humans (19-22).

In contrast, although transition metals are critical to cellular life, in abundance they are toxic. Copper accumulates within Mtb- associated pulmonary lesions. This is thought to be a strategy by which the immune system restricts mycobacterial growth in vivo (23)9. This is further supported by the observation that, in comparison to other bacteria, Mtb is particularly susceptible to copper in vitro. Moreover, Mtb lacking the copper transporter MctB accumulated 100 fold more intracellular copper *in vitro* and was markedly attenuated within guinea pigs (23). In addition, INFy mediated transportation of copper into the phagolysosomes of activated macrophages was lethal to resident intracellular bacteria (24). This lethality of copper is attributed to the fact that copper disables critical cellular enzymes by dislodging catalytic iron atoms. In addition, copper participates in Fenton chemistry, which results in the production of toxic hydroxyl radicals and may potentiate RNS mediated oxidative stress (25). Thus, the host capitalizes on the dual essentiality and toxicity of transition metals to eradicate invading pathogens (16).

#### The availability of molecular oxygen regulates mycobacterial growth.

In the host, Mtb is thought to reside within oxygen-depleted tissue (26, 27). Individual macrophages, in which Mtb becomes sequestered, contain lower oxygen tensions than the extracellular space (28). These infected cells then aggregate forming avascular structures called granulomas within which oxygen tensions approach 1% (29-32). Although sudden exposure to anaerobiosis is lethal, Mtb can adapt to conditions in which oxygen is gradually depleted (33). In response to the gradual withdrawal of oxygen from the medium, Mtb carries out the orderly shut down of various cellular processes and enters into a nonreplicating persistent (NRP) state beginning once oxygen tensions reach 1%. NRP bacteria no longer divide or synthesize DNA and demonstrate limited RNA and protein synthesis (34). Furthermore, NRP Mtb are no longer sensitive to antibiotics that target these inactive cellular processes (27, 35).

## Oxygen regulates the growth of Mtb within the host.

Before the advent of chemotherapy, surgical intervention to reduce Mtb's access to oxygen was a leading treatment for TB. Intentional lung atelectasis allowed for near-complete pathologic and clinical resolution of disease (36). While this suggests that mycobacterial growth requires access to oxygen *in vivo*, evidence that oxygen regulates the growth of Mtb within infected humans is, necessarily, indirect.

In order to investigate the *in vivo* oxygen dependence of Mtb, oxygen tensions were experimentally manipulated within the lung tissue of Mtb infected model organisms. Direct manipulation was achieved by

controlling the partial pressure of oxygen within inhaled air. Guinea pigs infected with Mtb and maintained in 10% oxygen contained reduced colony forming units (CFUs) within their lungs and spleens as compared to their Mtb-infected counterparts maintained under room air (21%) oxygen) (37). Alternatively, indirect manipulation of oxygen by re-routing a percentage of the blood supply away from otherwise well-ventilated alveoli created an artificially elevated partial pressure of oxygen within the associated alveolar sacs, as oxygen was no longer carried off into the circulation by hemoglobin. Monkeys that received a surgical intervention to re-route blood from lung tissue prior to infection with Mtb, for example by pulmonary artery ligation, developed increased mycobacterial numbers within the associated lung tissue (38). Thus these initial experiments supported the centrality of oxygen in regulating mycobacterial growth in vivo. Further supporting evidence arose from the observation that TBassociated lesions tend to localize within lung tissue containing the greatest oxygen tension.

The diagnosis of TB in humans is made in part by capturing radiographic images of the lungs. This led to the observation that in adults, pathologic lesions associated with chronic pulmonary TB are often localized within the lung apices (39). As a result of gravity, the apices receive less blood flow than more inferior lung tissue, and therefore contain greater oxygen tensions, 132 mmHg as compared to 89 mmHg in more inferior tissue (40, 41). In order to determine whether the higher oxygen tensions of the lung apices contributed to the localization of TB-associated pathology, Medlar and Sasano infected rabbits with Mtb and maintained a subset of infected animals in a vertical position to simulate

the erect posture of humans. Typically, Mtb infected (horizontal) rabbits develop tissue pathology in the posterior dorsal basal lung segments that contain the highest oxygen tensions. However, vertically oriented rabbits developed Mtb-associated lesions in the lung apices (42). While the results of this seminal experiment supported a relationship between oxygen tension and progressive TB lung pathology, an alternative explanation posited that impaired lymphatic clearance of antigens within the lung apices could account for these observations (40).

Although these historic experiments suggested an association between oxygen and mycobacterial growth *in vivo*, it is not possible to conclude from them that Mtb infected hosts are hypoxic, or that oxygen regulates the growth of Mtb in the host. In the modern era, however, investigators have developed tools to directly address whether Mtb infected host tissues are hypoxic. Molecular markers of hypoxia, such as the nitroimidazoles, mycobacterial hypoxic response gene expression analysis and direct measurement of tissue oxygen tensions with an oxygen probe have been used to demonstrate that Mtb infected model hosts, such as the guinea pig, rabbit, macaque develop hypoxic tissue pathology (30). These same techniques have also indicated that the C3HeB/FeJ mouse model, but not BALB/c (43) or C57BL/6 (31, 32) mouse strains, develop hypoxic TB-associated lesions. Therefore, within model hosts that have demonstrated hypoxic, TB-associated tissue pathology, Mtb likely experiences hypoxia. That the availability of oxygen regulates mycobacterial growth in vitro is clear. Mycobacteria no longer replicate once oxygen tensions fall to, or below 1% (26, 27). Thus oxygen likely regulates mycobacterial replication in vivo as well.

When oxygen is scarce, Mtb respires the alternative electron acceptor nitrate.

In response to hypoxia, Mtb induces the expression of the nitrate/proton symporter *narK2*, which allows for the rapid entry of nitrate into the cell (44). Nitrate is subsequently reduced to nitrite by the constitutively active nitrate reductase encoded by *narGHJI* (45). This process reportedly supports hypoxic ATP synthesis and redox balancing (46, 47). Moreover, the upregulation of this alternative respiratory pathway enhances the survival of Mtb challenged with sudden anaerobiosis (48). Human tissues contain abundant quantities of nitrate, which arise primarily from the diet and nitric oxide synthase (NOS) enzymes. NOS enzymes produce nitric oxide that auto-oxidize to form both nitrate and nitrite (49). Over the course of infection with Mtb, patients accumulate increasing concentrations of plasma nitrite (50, 51). This likely arises from the two NOS isoforms, endothelial and inducible (eNOS and iNOS respectively) that are expressed in the granulomatous tissue of TB patients (52-54). Mycobacterial nitrate reduction, which is responsible for the production of 2.5 mM nitrite by axenic cultures of Mtb (55), may also contribute to this pool of plasma nitrite.

### *Mtb respires nitrate within primary human macrophages.*

Our laboratory recently adopted an alternative protocol to differentiate primary human macrophages. There are several features that distinguish this protocol from others (56). In particular, monocytes are differentiated in 10% oxygen in the gas phase over 2 weeks in a medium

containing 40% human plasma, 60% RPMI and the cytokines GMCSF and TNF $\alpha$ . The resultant macrophages better restricted the replication of Mtb over a period of weeks (57), as compared to macrophages differentiated using more traditional protocols, which are incubated under room air oxygen tensions of a period of days, in a medium composed of DMEM, 10% fetal bovine serum and MCSF (56). Initial experiments revealed that, despite culture under non-hypoxic oxygen tensions, macrophage-resident Mtb produced significant amounts of nitrite. This was immediately of interest as nitrate and nitrite can be reduced to form nitric oxide *in vivo* (49).

The nitrate-nitrite-nitric oxide pathway begins with the reduction of nitrate to nitrite by oral commensal bacteria, which can then be further reduced to nitric oxide through an array of physiologic processes, many of which are non-enzymatic (49). The reduction of nitrate or nitrite to form nitric oxide has been shown to reduce blood pressure (58, 59), enhance the efficacy of mitochondrial oxidative phosphorylation (60), impair platelet aggregation (61), and to exert carcinogenic (62) and antimicrobial effects (49, 63, 64). These broad effects arise once nitrate or nitrite is reduced to nitric oxide. However, nitrite has also been proposed to directly react with hemes and thiols without the intermediary formation of nitric oxide (65).

### Conclusions

In the context of tuberculosis, nitrite is generally overlooked as simply a degradation product of NOS enzymes. This is in part due to the fact that nitrate is not a component of the standard cell medium used to culture Mtb axenically. In addition, in contrast to our protocol, traditional monocyte differentiation techniques typically involve the use of medium containing low levels of nitrate and so mycobacterial nitrate reduction is rarely appreciated in either axenic or co-culture experiments.

Although high concentrations of nitrite restrict the growth of Mtb without any associated lethality (55), this has remained mechanistically unexplained. Therefore we hypothesized that Mtb-derived nitrite might impact Mtb itself. Taken together, two central areas of investigation emerged from our observation that Mtb produced abundant amounts of infected macrophages: first, what had nitrite within prompted nitrate respiration, were macrophage-resident Mtb mycobacterial functionally hypoxic? Second, given the broad impact of the nitrate and nitrite on human physiology and the observation that nitrite impairs mycobacterial replication, we proposed to characterize additional consequences of Mtb-derived nitrite on the pathogen itself and to determine to what extent these effects resemble the well-documented effects of nitric oxide.

## References

- World Health Organization (2011) Global tuberculosis control: WHO report 2011.
- 2. Zumla A, Raviglione M, Hafner R, Fordham von Reyn C (2013) Tuberculosis. *N Engl J Med* 368:745–755.
- 3. Gutierrez MC et al. (2005) Ancient origin and gene mosaicism of the progenitor of *Mycobacterium tuberculosis*. *PLoS Pathog* 1:e5.
- 4. Nathan C (2009) Taming tuberculosis: a challenge for science and society. *Cell Host Microbe* 5:220–224.
- 5. Fennelly KP et al. (2004) Cough-generated aerosols of *Mycobacterium tuberculosis*: a new method to study infectiousness. *Am J Respir Crit Care Med* 169:604–609.
- 6. Fennelly KP et al. (2012) Variability of infectious aerosols produced during coughing by patients with pulmonary tuberculosis. *Am J Respir Crit Care Med* 186:450–457.
- 7. Schaible UE, Sturgill-Koszycki S, Schlesinger PH, Russell DG (1998) Cytokine activation leads to acidification and increases maturation of *Mycobacterium avium*-containing phagosomes in murine macrophages. *J Immunol* 160:1290–1296.
- 8. Nathan C, Shiloh MU (2000) Reactive oxygen and nitrogen intermediates in the relationship between mammalian hosts and microbial pathogens. *Proc Natl Acad Sci USA* 97:8841–8848.
- 9. Ehrt S, Schnappinger D (2009) Mycobacterial survival strategies in the phagosome: defence against host stresses. *Cell Microbiol* 11:1170–1178.
- 10. Lundberg JO, Weitzberg E, Gladwin MT (2008) The nitrate-nitritenitric oxide pathway in physiology and therapeutics. *Nat Rev Drug Discov* 7:156–167.

- Bryk R, Lima CD, Erdjument-Bromage H, Tempst P, Nathan C (2002) Metabolic enzymes of mycobacteria linked to antioxidant defense by a thioredoxin-like protein. *Science* 295:1073–1077.
- 12. Joseph SV, Madhavilatha GK, Kumar RA, Mundayoor S (2012) Comparative analysis of mycobacterial truncated hemoglobin promoters and the groEL2 promoter in free-living and intracellular mycobacteria. *Appl Environ Microbiol* 78:6499–6506.
- 13. Lee WL et al. (2009) *Mycobacterium tuberculosis* expresses methionine sulphoxide reductases A and B that protect from killing by nitrite and hypochlorite. *Mol Microbiol* 71:583–593.
- 14. Darwin KH, Nathan CF (2005) Role for nucleotide excision repair in virulence of *Mycobacterium tuberculosis*. *Infect Immun* 73:4581– 4587.
- 15. Newton GL, Buchmeier N, Fahey RC (2008) Biosynthesis and Functions of Mycothiol, the Unique Protective Thiol of Actinobacteria. *Microbiol Mol Biol Rev* 72:471–494.
- 16. Hood MI, Skaar EP (2012) Nutritional immunity: transitionmetals at the pathogen–host interface. *Nat Rev Microbiol* 10:525–537.
- 17. Byrd TF, Horwitz MA (1993) Regulation of transferrin receptor expression and ferritin content in human mononuclear phagocytes. Coordinate upregulation by iron transferrin and downregulation by interferon gamma. *J Clin Invest* 91:969–976.
- 18. Byrd TF, Horwitz MA (1989) Interferon gamma-activated human monocytes downregulate transferrin receptors and inhibit the intracellular multiplication of *Legionella pneumophila* by limiting the availability of iron. *J Clin Invest* 83:1457–1465.
- 19. Abel L et al. (1998) Susceptibility to leprosy is linked to the human NRAMP1 gene. *J Infect Dis* 177:133–145.
- 20. Bellamy R et al. (1998) Variations in the NRAMP1 gene and

susceptibility to tuberculosis in West Africans. *N Engl J Med* 338:640–644.

- 21. Roger M, Levee G, Chanteau S, Gicquel B, Schurr E (1997) No evidence for linkage between leprosy susceptibility and the human natural resistance-associated macrophage protein 1 (NRAMP1) gene in French Polynesia. *Int J Lepr Other Mycobact Dis* 65:197–202.
- 22. Vidal SM, Malo D, Vogan K, Skamene E, Gros P (1993) Natural resistance to infection with intracellular parasites: isolation of a candidate for Bcg. *Cell* 73:469–485.
- 23. Wolschendorf F et al. (2011) Copper resistance is essential for virulence of *Mycobacterium tuberculosis*. *Proc Natl Acad Sci* 108:1621–1626.
- 24. White C, Lee J, Kambe T, Fritsche K, Petris MJ (2009) A role for the ATP7A copper-transporting ATPase in macrophage bactericidal activity. *J Biol Chem* 284:33949–33956.
- 25. Macomber L, Imlay JA (2009) The iron-sulfur clusters of dehydratases are primary intracellular targets of copper toxicity. *Proc Natl Acad Sci* 106:8344–8349.
- 26. Sherman DR et al. (2001) Regulation of the *Mycobacterium tuberculosis* hypoxic response gene encoding alpha -crystallin. *Proc Natl Acad Sci USA* 98:7534–7539.
- 27. Wayne LG, Sohaskey CD (2001) Nonreplicating persistence of *Mycobacterium tuberculosis. Annu Rev Microbiol* 55:139–163.
- James PE, Grinberg OY, Michaels G, Swartz HM (1995) Intraphagosomal oxygen in stimulated macrophages. J Cell Physiol 163:241–247.
- 29. Ramakrishnan L (2012) Revisiting the role of the granuloma in tuberculosis. *Nat Rev Immunol* 12:352–366.
- 30. Via LE et al. (2008) Tuberculous granulomas are hypoxic in guinea

pigs, rabbits, and nonhuman primates. Infect Immun 76:2333-2340.

- 31. Aly S et al. (2006) Oxygen status of lung granulomas in *Mycobacterium tuberculosis*-infected mice. *J Pathol* 210:298–305.
- 32. Tsai MC et al. (2006) Characterization of the tuberculous granuloma in murine and human lungs: cellular composition and relative tissue oxygen tension. *Cell Microbiol* 8:218–232.
- 33. Wayne LG, Hayes LG (1996) An *in vitro* model for sequential study of shiftdown of *Mycobacterium tuberculosis* through two stages of nonreplicating persistence. *Infect Immun* 64:2062–2069.
- 34. Wayne LG (1977) Synchronized replication of *Mycobacterium* tuberculosis. Infect Immun 17:528–530.
- 35. Wayne LG, Sramek HA (1994) Metronidazole is bactericidal to dormant cells of *Mycobacterium tuberculosis*. *Antimicrob Agents Chemother* 38:2054–2058.
- 36. Meylan PR, Richman DD, Kornbluth RS (2010) Oxygen tensions and mycobacterial infections. *Clin Infect Dis* 15:372–373.
- 37. Rich AR, Follis RH (1942) The effect of low oxygen tension upon the development of experimental tuberculosis. *Bull Johns Hopkins Hospitl* 71:345–363.
- 38. Olson JB, Scott WH, Hanlon RC, Mattern TCF (1952) Experimental tuberculosis. III. Further observations on the effects of alteration of the pulmonary arterial circulation on tuberculosis in monkeys. *Am Rev Tuberc* 65:48–63.
- 39. Leung AN (1999) Pulmonary tuberculosis: the essentials. *Radiology* 210:307–322.
- 40. Goodwin RA (1983) Apical localization of pulmonary tuberculosis, chronic pulmonary histoplasmosis, and progressive massive fibrosis of the lung. *CHEST* 83:801.

- 41. West JB, Dollery CT (1960) Distribution of blood flow and ventilation-perfusion ratio in the lung, measured with radioactive carbon dioxide. *J Appl Physiol* 15:405–410.
- 42. Medlar EM, Sasano KT (1935) A study of the pathology of experimental pulmonary tuberculosis in the rabbit. *Am Rev Tuberc* 34:456–475.
- 43. Harper J et al. (2012) Mouse model of necrotic tuberculosis granulomas develops hypoxic lesions. *Journal of Infectious Diseases* 205:595–602.
- 44. Giffin MM, Raab RW, Morganstern M, Sohaskey CD (2012) Mutational analysis of the respiratory nitrate transporter NarK2 of *Mycobacterium tuberculosis*. *PLoS ONE* 7:e45459.
- 45. Sohaskey CD, Wayne LG (2003) Role of *narK2X* and *narGHJI* in hypoxic upregulation of nitrate reduction by *Mycobacterium tuberculosis*. *Journal of Bacteriology* 185:7247–7256.
- 46. Richardson DJ (2000) Bacterial respiration: a flexible process for a changing environment. *Microbiology (Reading, Engl)* 146 ( Pt 3):551–571.
- 47. Tan MP et al. (2010) Nitrate respiration protects hypoxic *Mycobacterium tuberculosis* against acid- and reactive nitrogen species stresses. *PLoS ONE* 5:e13356.
- 48. Sohaskey CD (2008) Nitrate enhances the survival of *Mycobacterium tuberculosis* during inhibition of respiration. *Journal of Bacteriology* 190:2981–2986.
- 49. Lundberg JO, Weitzberg E, Gladwin MT (2008) The nitrate–nitrite– nitric oxide pathway in physiology and therapeutics. *Nat Rev Drug Discov* 7:156–167.
- 50. Dlugovitzky D et al. (2000) Influence of disease severity on nitrite and cytokine production by peripheral blood mononuclear cells (PBMC) from patients with pulmonary tuberculosis (TB). *Clin Exp*

Immunol 122:343–349.

- 51. Lamsal M et al. (2007) Evaluation of lipid peroxidation product, nitrite and antioxidant levels in newly diagnosed and two months follow-up patients with pulmonary tuberculosis. *Southeast Asian J Trop Med Public Health* 38:695–703.
- 52. Choi H-S, Rai PR, Chu HW, Cool C, Chan ED (2002) Analysis of nitric oxide synthase and nitrotyrosine expression in human pulmonary tuberculosis. *Am J Respir Crit Care Med* 166:178–186.
- 53. MacMicking JD et al. (1997) Identification of nitric oxide synthase as a protective locus against tuberculosis. *Proc Natl Acad Sci USA* 94:5243–5248.
- 54. Nicholson S et al. (1996) Inducible nitric oxide synthase in pulmonary alveolar macrophages from patients with tuberculosis. *J Exp Med* 183:2293–2302.
- 55. Wayne LG, Hayes LG (1998) Nitrate reduction as a marker for hypoxic shiftdown of *Mycobacterium tuberculosis*. *Tuber Lung Dis* 79:127–132.
- Jordao L, Bleck CKE, Mayorga L, Griffiths G, Anes E (2008) On the killing of mycobacteria by macrophages. *Cell Microbiol* 10:529– 548.
- 57. Vogt G, Nathan C (2011) *In vitro* differentiation of human macrophages with enhanced antimycobacterial activity. *J Clin Invest* 121:3889–3901.
- 58. Carlstrom M et al. (2011) Dietary nitrate attenuates oxidative stress, prevents cardiac and renal injuries, and reduces blood pressure in salt-induced hypertension. *Cardiovascular Research* 89:574–585.
- 59. Larsen FJ, Ekblom B, Sahlin K, Lundberg JO, Weitzberg E (2006) Effects of dietary nitrate on blood pressure in healthy volunteers. *N Engl J Med* 355:2792–2793.

- 60. Larsen FJ et al. (2011) Dietary inorganic nitrate improves mitochondrial efficiency in humans. *Cell Metab* 13:149–159.
- 61. Webb AJ et al. (2008) Acute blood pressure lowering, vasoprotective, and antiplatelet properties of dietary nitrate via bioconversion to nitrite. *Hypertension* 51:784–790.
- 62. Tannenbaum SR, Correa P (1985) Nitrate and gastric cancer risks. *Nature* 317:675–676.
- 63. Dykhuizen RS et al. (1996) Antimicrobial effect of acidified nitrite on gut pathogens: importance of dietary nitrate in host defense. *Antimicrob Agents Chemother* 40:1422–1425.
- 64. Reddy D, Lancaster JR, Cornforth DP (1983) Nitrite inhibition of *Clostridium botulinum*: electron spin resonance detection of ironnitric oxide complexes. *Science* 221:769–770.
- 65. Bryan NS et al. (2005) Nitrite is a signaling molecule and regulator of gene expression in mammalian tissues. *Nat Chem Biol* 1:290–297.

### **CHAPTER 2**

# *MYCOBACTERIUM TUBERCULOSIS* GENERATES NITRITE IN HUMAN MACROPHAGES AT PHYSIOLOGIC OXYGEN TENSIONS: IMPACT ON MYCOBACTERIAL ATP CONSUMPTION AND GENE EXPRESSION<sup>\*</sup>

### Summary

In high enough concentrations, such as produced by inducible nitric oxide synthase (iNOS), reactive nitrogen species (RNS) can kill *Mycobacterium* tuberculosis (Mtb). Lesional macrophages in macaques and humans with tuberculosis (TB) express iNOS, and mice need iNOS to avoid succumbing rapidly to TB. However, Mtb's own ability to produce RNS is rarely considered, perhaps because nitrate reduction to nitrite is only prominent in axenic Mtb cultures at oxygen tensions  $\leq 1\%$ . Here we found that cultures of Mtb-infected human macrophages cultured at physiologic oxygen tensions produced copious nitrite. Surprisingly, the nitrite arose from the Mtb, not the macrophages. Mtb responded to nitrite by ceasing growth; elevating levels of ATP through reduced consumption; and altering the expression of 120 genes associated with adaptation to acid, hypoxia, nitric oxide, oxidative stress, and iron deprivation. The transcriptomic effect of endogenous nitrite was distinct from that of nitric oxide. Thus, whether or not Mtb is hypoxic, the host expresses iNOS, or hypoxia impairs the action of iNOS, Mtb in vivo is likely to encounter RNS by producing nitrite. Endogenous nitrite may slow Mtb's growth and prepare it to resist host stresses while the pathogen waits for

<sup>\*</sup> Amy Cunningham-Bussel, Tuo Zhang, and Carl Nathan, "Mycobacterium Tuberculosis Generates

immunopathology to promote its transmission.

#### Introduction

The ability to produce RNS is widely distributed among animals, plants and eubacteria. The high-output pathway of RNS production by host iNOS contributes to control of diverse infections (1, 2), including tuberculosis (TB) (3, 4), via impacts on both the pathogen and the host (5). Macrophages in the lungs of humans (3, 6, 7) and macaques (8) with TB express functional iNOS. Correlative evidence suggests that the action of iNOS may contribute to human control of TB (1, 2, 9, 10).

Contributions of bacterial RNS production to host-pathogen interactions have only recently come under study. Bacterial NO synthases or nitrate reductases are sources of RNS that can contribute to antibiotic resistance (3, 4, 11, 12) and increase virulence (5, 13). Some commensal bacteria exploit host iNOS during colonic inflammation by using iNOS-derived nitrate ( $NO_3^{-1}$ ) as a terminal electron acceptor (3, 6, 7, 14). Nitrate, along with nitrite ( $NO_2^{-1}$ ), is an auto-oxidation product of nitric oxide ( $NO_3$ ), the product of iNOS.

In the host, Mtb is thought to survive over decades in hypoxic sites (8, 15, 16). When oxygen is scarce, Mtb reduces nitrate  $(NO_3^{-})$  to nitrite  $(NO_2^{-})$  as a means to maintain redox homeostasis and energy production (17). The mycobacterial respiratory nitrate reductase, encoded by *narGHJI*, is constitutively expressed (17), and functions at a low level under aerobic conditions when nitrate is taken up by passive diffusion (18). Hypoxia promotes the induction of *narK2*, a nitrate transporter and a member of the dormancy survival (*DOS*) regulon (17, 19-21), such that

Mtb cultured at  $\leq 1\%$  oxygen in the presence of nitrate rapidly produces nitrite up to a level of 2.5 mM (22). Further production is restricted in association with non-lethal cessation of bacterial growth (22).

Transcripts from narG, which encodes a subunit of the nitrate reductase, and *narX*, which lies downstream of *narK2* and encodes a nonfunctional nitrate reductase, were identified in granulomas from humans with TB (23, 24). This suggests that populations of mycobacteria residing within the human host are exposed to low oxygen tensions, and likely respire nitrate to adapt. Nitrate is a physiologic component of human body fluids, where it arises from the diet and from auto-oxidation of the NO produced not only by iNOS (NOS2) but also by the constitutively expressed enzymes NOS1 and NOS3. Studies in which nitrate was furnished *in vitro* revealed that Mtb's ability to respire nitrate allows it to better withstand acid, nitrosative stress (25) or sudden anaerobiosis (26). The evolutionary success of "modern" Mtb strains, which are more prevalent than "ancestral" strains, has been attributed to their enhanced nitrate reductase activity (27). Although nitrate and nitrite reproduce the bioactivity of nitric oxide *in vivo* in a wide array of physiologic processes (28), reduction of nitrate to nitrite is rarely considered in studies of the host-pathogen relationship in TB. In fact, nitrate was omitted when the standard growth media were formulated in which almost all in vitro experiments with Mtb are currently conducted.

During the course of modifying standard culture conditions for Mtb-infected human monocyte-derived macrophages (29), we observed abundant accumulation of nitrite in the supernatant. Herein, we report the identification of Mtb as both the source of the nitrite and a highly responsive target.

#### Results

*Mtb respires nitrate in human macrophages cultured at non-hypoxic oxygen tensions.* 

We differentiated normal donors' monocytes into macrophages under 10% oxygen for 2 weeks, activated them with interferon-g and infected them with Mtb, as described (29). Cultures of macrophages from multiple donors consistently accumulated nitrite in a time-dependent manner to a median level of 25 µM over 5 days (Figure 1a). This suggested that the macrophages may have expressed one of the three NOS isoforms. However, we detected no iNOS protein by immunoblot (Supplementary Figure S1, see Appendix 1) and no NOS1, NOS2 or NOS3 transcripts by microarray or RNA Seq (not shown). To see if Mtb might be the source, we infected macrophages with a strain of Mtb lacking narG. The mutant strain grows normally in aerobic culture and is as virulent as wild type Mtb in mice (15), but cannot respire nitrate. Macrophage cultures infected with  $\Delta narG$  Mtb accumulated no nitrite, while infection with a mutant strain complemented with the wild type *narG* allele led to accumulation of as much nitrite as infection with wild type (WT) Mtb (Figure 1B). Accumulation of nitrite was nearly abolished when the cultures were washed gently to replace most of the standard culture medium with medium containing negligible nitrate, and was restored when the nitratedeficient cultures were repleted with 5 mM nitrate (Figure 1C). Even when we cultured infected macrophages in conventional, hyperoxic levels of oxygen (21%), we observed *narG*-dependent nitrite accumulation in

the supernatant (Figure 1D). To further confirm that the resultant nitrite was a mycobacterial, rather than a host product, we infected macrophages with Mtb in the presence of the bactericidal antibiotic, rifampicin. Addition of rifampicin to macrophages 2 days prior to infection completely abolished nitrite production after infection. Treatment with rifampicin even 24 or 48 hours post infection markedly reduced the production of nitrite (Figure 1E).

We wondered what levels of Mtb-derived nitrite might be attained in tissue sites such as granulomas, where the number of macrophages per unit volume vastly exceeds the non-physiologic ratio attainable in monolayer culture. To better simulate the cell densities of tissues, we cultured macrophages on Cytodex microcarrier beads. The *narG*dependent accumulation of nitrite in bead cultures reached 800  $\mu$ M (Figure 1F). In both monolayer and bead culture, the colony-forming units (CFU) of Mtb recovered from the cultures were not significantly different for *narG*-deficient and WT Mtb (Cunningham-Bussel, *MicrobiologyOpen*, in press). Figure 1. Nitrite generation by Mtb within primary human macrophages cultured in non-hypoxic levels of oxygen. (A) Accumulation of nitrite in supernatant of macrophages in monolayer culture in 10% oxygen infected with wild type (Wt) Mtb (MOI:5;  $5x10^5$ bacteria) for the indicated times. Box and whisker plots depict median and 5<sup>th</sup>-95<sup>th</sup> percentiles of results from 6 donors. (B) As in (A), but macrophages were uninfected or infected with *narG*-deficient Mtb (NarG) or Wt or complemented (NarGc) strains. Results for uninfected macrophages and those infected with NarG Mtb overlap near zero. Means  $\pm$  SD from one experiment representative of > 10. (C) Lack of accumulation of nitrite in nitrate-depleted medium. Macrophages were infected in the presence (\*) or absence of 5 mM nitrate. Means  $\pm$  SD for one experiment representative of 2. (D) As in (B), but with macrophages maintained under either 21% or 10% oxygen. Means ± SD for one experiment representative of 2. (E) Impact of rifampicin (RIF) on nitrite production. Where indicated, macrophages were infected with Wt Mtb (MOI: 5) for 3 days in 10% oxygen in the presence or absence of rifampicin (1 µg mL<sup>-1</sup>). Rifampicin was added 2 days before, 24 hrs after or 48 hrs after infection. Uninfected macrophages were treated with rifampicin for the duration of the experiment. Nitrite was measured on day 3 after infection. Means  $\pm$  SD, n= 2 experiments. (F) As in (B) but using ~20-fold higher numbers of macrophages grown on Cytodex surface microcarrier beads in 10% oxygen and then infected with Mtb (MOI: 5;  $10^7$  bacteria). Means  $\pm$  SD, n= 2 experiments.

Figure 1 (continued)









Oxygen tension within Mtb-infected human macrophages functionally approximates 1%.

Abundant mycobacterial nitrate reduction within infected macrophages incubated under ambient oxygen tensions of 10% and 21%suggested that Mtb experienced an intracellular oxygen tension that approximated 1%. To test this, we compared the expression of DOS regulon members hspX and fdxA by Mtb cultured axenically or within infected macrophages. As reported (20), we found that Mtb cultured axenically in standard medium under 1% oxygen but not 21% oxygen induced both genes by ~100-fold (Figure 2A). Mtb residing within macrophages also highly induced the two DOS genes, particularly under 10% oxygen (Figure 2B) and less so under 21% oxygen (Figure 2C). To confirm this surprising finding, we treated Mtb-infected macrophages with pimonidazole. At oxygen tensions  $\leq 1\%$  pimonidazole, is reduced to form covalently modifies that proteins, detected a as by immunocytochemistry (30). Macrophages cultured at an oxygen tension of 10% showed intense pimonidazole staining, whether or not they were Mtb-infected (Figure 2D). Even under hyperoxic conditions (21%) oxygen), considerable pimonidazole staining was detected (Figure 2D). Thus, despite incubation under ambient tensions of 10% or even 21%, intracellular oxygen tensions in the human macrophages functionally approximated 1%.


Figure 2. Functional evidence for intracellular hypoxia in primary human macrophages cultured at physiologic tissue-level (10%) or hyperoxic (room air) oxygen tensions. (A) Expression of DOS regulon genes *hspX* and *fdxA* in axenic cultures of Wt or *narG*-deficient (NarG) Mtb in 21% or 1% oxygen in standard medium without nitrate,  $OD_{580}$  0.1 (means  $\pm$  SD, n=2 experiments). (B, C) Expression of *hspX* and *fdxA* by Wt or NarG strains within human macrophages incubated in 21% or 10% oxygen for 10 hours (MOI: 40). Means  $\pm$  SD, n=2 experiments. (D) Pimonidazole staining of macrophages infected or not with Wt Mtb and 5) incubated the oxygen tensions indicated. (MOI: at Photomicrographs: Representative images. Blue: DAPI- stained nuclei, Red: pimonidazole adducts. Graph: Quantification of pimonidazole staining intensity. 0 PIMO, uninfected cells treated with neither pimonidazole nor DMSO. DMSO, uninfected cells treated with only the vehicle. Each dot represents the average intensity in  $\sim 500$  macrophages. Horizontal lines indicate the means of 2 independent experiments.

Endogenously generated nitrite represses mycobacterial growth and ATP consumption.

As noted by Wayne and Hayes (22), nitrite, whether endogenously produced or exogenously supplied, suppressed the growth of Mtb under 21% oxygen (Figure 3A). The concentration-dependence of this effect was similar for WT, *narG*-deficient and complemented Mtb (Figure 3A). In contrast, when we incubated the 3 strains in 1% oxygen for 3 days with 5 mM nitrate, followed by return to 21% oxygen, only the WT and complemented strains failed to resume growing, reflecting that they generated endogenous nitrite from nitrate, which *narG*-deficient Mtb could not (Figure 3B). The failure of cultures of WT Mtb to increase in turbidity after generating nitrite reflected bacteriostasis, not cell death, because the number of CFU recovered was unaffected (Figure 3C).



Figure 3. Nitrite represses mycobacterial growth. (A, B) Strains of Mtb designated as in Figure 1 ( $OD_{580}$  0.1, as marked by horizontal dotted lines) were incubated in the presence of the indicated concentrations of nitrite (A) or nitrate (B) for 3 days in 1% oxygen and then transferred to 21% oxygen for 5 days before growth was assessed by measuring  $OD_{580}$ . Means  $\pm$  SD, n=2 experiments. (C) As in (A) and (B), except that CFU were determined at the end of the 3-day incubation in 1% oxygen for wild type Mtb. Means  $\pm$  SD, n=2 experiments.

It is not known how nitrite impairs aerobic mycobacterial growth. We suspected an impact of nitrate respiration on intrabacterial ATP, because nitrite has been shown to inhibit bacterial respiration and prevent ATP synthesis (31). To test this, we cultured WT, *narG*-deficient and

complemented strains of Mtb axenically for 3 days under 1% oxygen either at high density (OD: 0.1, ~  $5 \times 10^7$  cells mL<sup>-1</sup>) or low density (OD: 0.01, ~  $5 \times 10^6$  cells mL<sup>-1</sup>) in the presence of 5 mM nitrate and measured ATP in their lysates. As expected, in the low-density cultures the concentration of nitrite remained low. Under these conditions, ATP levels in WT and *narG*-deficient Mtb were not significantly different (Figure 4A). In contrast, high-density cultures of the WT and complemented strains contained significantly more ATP than the *narG*-deficient strain, confirming an observation of Tan et al. (25) (Figure 4A).

Tan et al. (25) attributed this result to increased ATP production by Mtb using nitrate as an electron acceptor when oxygen was limiting. However, when we added 2.5 mM exogenous nitrite to low-density cultures of Mtb in 1% oxygen, ATP levels were increased even farther than seen with WT Mtb given twice as much nitrate (Figure 4B). The ATP-elevating effect of exogenous nitrite was not dependent on the presence of *narG* or *nirBD*, which encodes a nitrite reductase (32). Thus, it appeared that nitrite, a product of nitrate respiration, led to accumulation of ATP in Mtb in a manner that did not depend on Mtb's donation of electrons to nitrate.

To test if nitrite itself was responsible for this effect, as opposed to nitric oxide, to which nitrite can be converted, we incubated low-density Mtb cultures with 5 mM nitrate as a control or with diethylenetriamine-NO adduct (DETANO), a compound that decomposes to release nitric oxide with a half-life of ~22 hours at pH 7.4. DETANO also elevated intrabacterial ATP levels in WT, *narG*-deficient and the complemented strains. Both nitrite and DETANO increased intrabacterial ATP in a

concentration-dependent manner. DETANO was ~25-fold more potent than nitrite (Figure 4C, D). Thus, either nitrite or nitric oxide potentially arising from it appeared to increase ATP.

We next considered whether nitrite or its possible conversion products might increase ATP production or decrease ATP consumption. Nitrite repressed the growth of Mtb; therefore, we considered it unlikely that nitrite enhanced the synthesis of ATP. Thus, we compared the consumption of exogenous ATP in lysates from nitrite-treated and untreated Mtb, after first demonstrating that neither nitrite nor nitric oxide interfered with the ATP assay (not shown). WT Mtb was incubated with or without nitrite as described above for 3 days in 1% oxygen. Whole cell lysates were prepared on ice, moved to room temperature and allowed to consume endogenous ATP until it was no longer detectable, which required less than 20 minutes. We then added various amounts of the lysates and immediately measured its exogenous ATP to concentration. We consistently recovered greater amounts of exogenous ATP in lysates from nitrite-treated Mtb than untreated Mtb (Figure 4E). Time-course measurements of the disappearance of exogenous ATP (Supplementary Figure S2, see Appendix 1) supported the interpretation that consumption of ATP was slower in lysates from nitrite-treated Mtb than from untreated Mtb.

Figure 4. Endogenously generated or exogenously supplied nitrite elevates ATP in Mtb and suppresses ATP consumption in its lysates. (A) Intrabacterial ATP content measured in wild type (Wt), narGdeficient (NarG) or complemented (NarGc) strains of Mtb incubated at high (0.1) or low  $OD_{580}$  (0.01) for 3 days in 1% oxygen in the presence of 5 mM nitrate. Results are expressed per mg protein in the lysate and are means ± SEM from 2 experiments.) (B) Impact of exogenous nitrite or nitrate on ATP content in Wt, narG-deficient (NarG), nirB-deficient (NirB) and respective complemented strains (NarGc, NirBc) of Mtb incubated for 3 days in 1% oxygen. (Means  $\pm$  SEM, n=3 experiments). (C) As in (B), but comparing the impact of single concentrations of nitrate (5 mM) and DETANO (200  $\mu$ M). Means ± SEM, n=3 experiments. (D) As in (C), but comparing the impact of varied concentrations of nitrate (yaxis on the right) and DETANO (y-axis on the left). Means  $\pm$  SEM, n=2 experiments. (E) Recovery of exogenous ATP added to lysates from Wt Mtb that had been incubated in the presence or absence of 2.5 mM nitrite for 3 days at 1% oxygen. ATP was added only after cell lysates had consumed endogenous ATP. Means  $\pm$  SEM, n=2 experiments. P values were determined by unpaired t tests.

## Figure 4 (continued)



#### Endogenously generated nitrite regulates the Mtb transcriptome.

Having demonstrated an effect of nitrite on ATP metabolism, we characterized the impact of endogenously generated nitrite more broadly by examining its effect on Mtb gene expression as assessed by RNA Seq. Samples of Mtb were prepared following the same protocol that revealed that mycobacterial nitrate respiration repressed subsequent bacterial growth in air (Figure 3a-b). In 3 independent experiments, we treated axenic cultures of Mtb with or without 5 mM nitrate for 3 days at 1% oxygen to allow for the generation of endogenous nitrite and then transferred the cultures to 21% oxygen overnight. Figure 5 refers to these samples as "untreated" and "nitrate", respectively. Two further control cultures were prepared as above, except that one was treated with 2.5 mM nitrite in place of nitrate and the other was exposed neither to nitrate nor nitrite but its RNA was collected at the conclusion of the 3 day incubation in 1% oxygen, without an  $\sim$ 18 hour culture in 21% oxygen. Figure 5 refers to these two samples as "nitrite" and "untreated 1% oxygen," respectively. Ribosomal RNA was removed from the total RNA by oligonucleotide-based hybridization. Sequences determined in three separate sequencing runs were aligned to the Mtb H37Rv genome (Accession NC\_000962.3). After culling sequences encoding ribosomal RNA, ~3x10<sup>6</sup> reads per sample were mapped to coding regions, covering ~99.9% of the coding genome. Gene expression was reported as reads per kilobase per million mapped reads (RPKM).

We found that 120 genes were significantly regulated by provision of nitrate. Those induced are listed in Supplementary Table S1 and those repressed are listed in Supplementary Table S2, see Appendix 1.

32

Inspection of the list suggested that the most informative comparisons would be with reports of Mtb's transcriptional adaptation to host-imposed stresses (33), such as hypoxia (20), iron restriction (34), acid (pH 5.5) (35), hydrogen peroxide (10 mM) (36), nitric oxide (50 mm) (21), and residence in mouse macrophages (35, 37)

We first analyzed expression of the *DOS* regulon, which is induced by three of these conditions—hypoxia (20), nitric oxide (21), and residence in macrophages (35, 37). The average expression of all 47 *DOS* regulon genes was nearly the same for the "untreated" and "untreated 1% oxygen" samples (Figure 5A, first two columns). The close correspondence of *DOS* induction in these two conditions was also evident for each *DOS* gene considered individually (Figure 5B).

Surprisingly, in comparison to the untreated sample, the average expression of the *DOS* regulon was significantly repressed by provision of nitrate (p< 0.0004) (Figure 5A). Individually, 27 *DOS* genes were significantly repressed in nitrate-treated Mtb (Figure 5C). (To achieve statistical significance for all 47 *DOS* genes considered individually, we would likely need to sequence more samples.) This effect of nitrate could be attributed to generation of nitrite, because the effect was qualitatively and quantitatively the same when nitrite itself was provided (Figure 5D). Finally, the effect of nitrite could not be ascribed to conversion to nitric oxide, in that nitric oxide has the opposite effect: it induces the *DOS* regulon, rather than repressing it. This is shown in Figure 5E, which plots the present results for 28 *DOS* genes: 27 that were significantly repressed by nitrite and reported to be induced by hypoxia (20) and 28 that were significantly repressed by nitrite and reported to be induced by 50 mM

nitric oxide (21). Hypoxic cultures of Mtb have been reported to accumulate up to 2.5 mM nitrite and no more (22). These results offer an explanation: nitrite negatively regulates its own accumulation by limiting the expression of the DOS regulon and thus *narK2* (Supplementary Figure 3, see Appendix 1).

Figure 5. Respiration of nitrate represses expression of the DOS regulon in Mtb. (A-D) RPKM for DOS genes as determined by RNA Seq for Wt Mtb incubated in 1% oxygen for 3 days ("untreated 1% oxygen") or incubated in 1% oxygen for 3 days in the absence ("untreated") or presence of 5 mM nitrate ("nitrate") or 2.5 mM nitrite ("nitrite"), as indicated and transferred to 21% oxygen overnight. (A) The average expression of all 47 DOS genes across all samples (means  $\pm$  SD) for the "untreated" and "nitrate" samples in 3 independent samples per condition, for the "untreated 1% oxygen" and "nitrite" samples (n=1 experiment per condition; see Methods). The p value was determined by an unpaired t test. (B) Gene-by-gene comparison of expression in the "untreated 1% oxygen" sample and the "untreated" sample. (C) Expression of the DOS genes that were significantly regulated by nitrate respiration compared to expression of each such gene in the untreated sample. (D) Expression of each DOS gene from Wt type Mtb treated with 5 mM nitrate as compared to a sample treated with 2.5 mM nitrite. (B-D) Red lines indicate 1:1 correlation; dashed lines indicate a 20% deviation from a 1:1 correlation. (E) Comparison of the expression of individual genes significantly regulated by nitrate in these experiments, with their reported regulation in hypoxia (20) or upon exposure to DETANO as a source of nitric oxide (21). The x-axis indicates the regulation of shared genes by the condition of interest and the y-axis indicates their regulation by nitrate respiration. All y values above or below the grey bar were regulated greater than 2fold.

**Figure 5 (continued)** 



Of the 120 genes that were significantly regulated by nitrate respiration, many overlapped with genes that were regulated by Mtb residing within activated mouse macrophages (35, 37) (Figure 6AB). In addition, Schnappinger *et al.* reported a subset of genes that were specifically regulated by iNOS. However, there was no correlation between regulation of genes by Mtb's respiration of nitrate and those regulated specifically by Mtb's residence in macrophages that expressed iNOS (37) (Figure 6A). This provided further evidence that the effect of endogenously generated nitrite was not solely mediated by conversion to nitric oxide.

Many Mtb genes regulated by nitrate respiration were reported to be similarly regulated in Mtb exposed to iron restriction (34) (Figure 6C), pH 5.5 (35) (Figure 6D), or treatment with 10 mM hydrogen peroxide (36) (Figure 6E). Table 1 presents the total number of genes regulated by nitrate respiration and reported to be regulated by these and other physiologic stimuli. After standardization to account for gene lists of differing size, the comparison indicates the number of genes expected to overlap by chance and whether the overlap is statistically significant. This analysis suggested that nitrate regulates the expression of genes that are also regulated when Mtb is exposed to several host-imposed stresses.



Figure 6. Comparison of the impact of nitrate respiration on the Mtb transcriptome with the impact of (A, B) residence within macrophages, (C) iron restriction, (D) exposure to pH 5.5, or (E) treatment with 10 mM hydrogen peroxide. Results of the present experiments and literature reports are graphed as in Figure 5E. The literature results were from the following studies. (A) (37). (B) (56). (C) (34) (D) (56). (E) reported by (36).

Table 1. Comparison of regulation of expression of Mtb genes by nitrate respiration observed here and by physiologic situations or stresses as reported in the literature. The total number of genes induced or repressed by nitrate respiration is listed along with the total number reported to be regulated by the indicated condition. The number regulated in common is compared to the number that would be expected to show a similar regulation by chance calculated by a Fisher's exact test. The columns labeled "number induced/repressed by nitrate respiration" refer only to the number of overlapping genes. "Percent of total gene list that overlaps" indicates the percent of all genes regulated by nitrate respiration that are also regulated by the condition of interest. The associated significance of this overlap, listed as a p value in the final column, was calculated using a Fisher's exact test. Of note, genes regulated by mitomycin C-induced DNA damage and starvation did not significantly overlap with genes regulated by nitrate respiration.

# Table 1 (continued)

Condition	Regulation	Genes regulated by the indicated condition	Total number that overlap (by chance)	Number induced by nitrate respiration	Number repressed by nitrate respiration	Percent of total gene list that overlaps	Significance of the overlap, p value
Iron depletion <sup>1</sup>	Induced	66	14 (2)	. 13		11.67	4.17E-08
(Low iron culture: 2 μM FeCl <sub>3</sub> vs High iron culture: 50 μM FeCl <sub>3</sub> , H37Rv)	Repressed	27	2 (1)	0	2	1.67	2.05E-01
<b>Hypoxia<sup>2</sup></b> (2 hrs at 0.2% oxygen, H37Rv)	Induced	47	27 (1)	0	27	22.50	< 2.2e-16
<b>Enduring Hypoxic Response<sup>3</sup></b> (Incubation at 0.2% oxygen over 4 days, H37Rv)	) Induced	230	17 (7)	12	IJ	14.17	6.53E-04
Nitric oxide <sup>4</sup> (50 µM DETA/NO for 40 minutes, H37Rv)	Induced	48	28 (1)	0	28	23.33	< 2.2e-16
Hydrogen peroxide <sup>5</sup> (10 mM, 40 minutes, clinical isolate 1254)	Induced Repressed	166 48	22 (5) 10 (1)	22 10	0 0	18.33 8.33	1.11E-08 4.65E-06
<b>pH 5.5</b> <sup>6</sup> (pH 5.5 vs pH 7, H37Rv)	Induced Repressed	261 375	16 (8) 18 (11)	15 1	1 17	13.33 15.00	5.72E-03 3.27E-02
Intraphagosomal <sup>7</sup> (24 hrs, H37Rv)	Induced Induced, iNOS dependent Repressed Repressed, iNOS dependent	454 60 147 8	65 (14) 32 (2) 11 (4) 3 (1)	35 0 0	30 23 11	54.17 26.67 9.17 2.50	<pre>&lt; 2.2e-16</pre> <pre>&lt; 2.2e-16</pre> <pre>5.63E-03</pre> <pre>3.31E-03</pre>
lntraphagosomal <sup>8</sup> (24 hrs, CDC 1551)	Induced Repressed	266 111	32 (8) 1 (3)	6 0	23 1	26.67 0.83	2.39E-11 9.65E-01
DNA damage <sup>°</sup> (Induced by mitomycin C, H37Rv)	Induced Repressed	113 26	1 (3) 2 (1)	1 2	0 0	0.83 1.67	9.67E-01 1.94E-01
<b>Starvation<sup>10</sup></b> (24 hrs after growth in PBS, H37Rv)	Induced Repressed	252 301	6 (8) 12 (9)	99	0	5.00 10.00	7.69E-01 1.94E-01
Nitrate respiration (Genes regulated by nitrate respiration and no other physiologic stimuli reported above)	I	I	1	9	5	10.83	ı
<sup>1</sup> Rodriguez 2002 <sup>6</sup> Rohde 2007 <sup>2</sup> Sherman 2001 <sup>7</sup> Schnappinger 2003 <sup>8</sup> Rustad 2008 <sup>8</sup> Rohde 2007 <sup>4</sup> Voskuil 2003 <sup>9</sup> Rand 2003 <sup>5</sup> Voskuil 2011 <sup>10</sup> Beste 2007							

#### Discussion

Within the host, both host NOSs and mycobacterial nitrate reductase can serve as sources of nitrite. However, while mycobacterial nitrate reduction is markedly induced by hypoxia (17), NOSs require oxygen as a substrate (38), and at oxygen tensions of 1%, iNOS loses 80-90% of its activity (39). Therefore, the activities of host iNOS and mycobacterial nitrate reductase could be spatially and temporally segregated, depending on their access to molecular oxygen.

The studies reported here led to four inter-related findings. First, there can be a striking dissociation between the oxygen tension in culture and the functional oxygen concentration in macrophage phagosomes. Although a dissociation was previously noted between oxygen concentration in extracellular fluid and in mouse macrophage phagosomes (40), to our knowledge this is the first report that the level of oxygen in the phagosomes of primary human macrophages cultured at physiologic tissue oxygen levels of 10% or even 21% approximates 1% functionally, as reflected by Mtb's induction of DOS genes and copious reduction of nitrate. The low functional level of intramacrophage oxygen was not dependent on ingestion of Mtb. In agreement with James et al. (29, 40), we speculate that the steep gradient between atmospheric and intracellular oxygen reflects a rate of mitochondrial oxygen consumption that outpaces diffusion from the gas phase through the extracellular fluid and across the plasma membrane. If Mtb likewise experiences functional hypoxia within human macrophages at physiologic oxygen tensions in vivo, then Mtb's respiration of nitrate to nitrite is likely to be much more widespread than previously appreciated, and the convention of culturing Mtb in room air levels of oxygen in the absence of nitrate is likely to be misleading with regard to Mtb's physiology.

Second, nitrite, whether endogenously produced or exogenously supplied, affects Mtb profoundly, although we may have only scratched the surface in characterizing its effects. Besides halting Mtb's growth, impairing its consumption of ATP and altering its transcriptome, mycobacterial nitrite also markedly reduces Mtb's susceptibility to isoniazid (Cunningham-Bussel, *MicrobiologyOpen*, in press). We do not know the major ATP-consuming pathway(s) in Mtb that function(s) in whole cell lysates and that nitrite inhibits. This contrasts with the reported effect of nitrite to reduce the amount of intrabacterial ATP in *P. aeruginosa* (31, 41).

Third, the effects of nitrite and nitric oxide partially overlap but in some respects are diametrically opposed. The effect of nitrite is also distinct from the effect of the respiration of nitrate, from which the nitrite arises. The ability of nitrite to affect biologic systems in its own right, rather than through conversion to more reactive forms, has been proposed (42, 43), and is supported by these results. For example, nitrite can oxidize ferrous iron to the ferric state and displace iron from iron-sulfur clusters (44, 45). The induced expression of mycobactin synthase genes suggests that Mtb-derived nitrite inactivated iron-containing enzymes, rendering the Mtb functionally iron-deficient despite the high iron content of the medium. *DOS* regulon expression is coordinated by the heme-containing kinases DosS and DosT, which function as redox and hypoxia sensors, respectively. Hypoxia, carbon monoxide and nitric oxide are physiologic ligands that induce DOS regulon expression by modulating

the oxidation and ligation states of the iron atoms in DosS and DosT (46). Nitrite may repress *DOS* regulon expression by oxidizing the iron of these sensor kinases in a manner that prevents their regulation by other physiologic ligands. Oxidation of iron in additional enzymes, such as ribonucleotide reductase, may also contribute to the ability of nitrite to slow or halt Mtb's growth (47-49).

Last, the shared regulation of genes by nitrate and by the potentially mycobactericidal conditions of iron restriction and exposure to acid and hydrogen peroxide suggests that nitrate respiration may benefit Mtb *in vivo* by preparing it to withstand the latter stresses. Nitrite is highly diffusible, as evidenced by its accumulation in the plasma of Mtb-infected individuals (50-52). Endogenously produced nitrite may diffuse within infected tissue and contribute to a pre-emptive induction of Mtb genes that benefit Mtb's survival in a hostile host environment.

To date, *in vivo* experiments to investigate the consequence of mycobacterial nitrate respiration have been limited to the mouse, where "granulomas" are areas of pneumonitic consolidation lacking the layered, fibrous-capped structure found in the granulomas of humans and non-human primates with TB. Mtb-infected granulomatous tissue in the mouse had a measured oxygen tension averaging ~40 mm Hg (or 2-9% oxygen) (15, 53) and did not stain with a hypoxia-sensitive 2-nitroimidazole probe (20, 54). Furthermore, Mtb in mice did not express the only DOS regulon member tested, *narK2*, until the onset of the adaptive immune response almost three weeks after infection, such that DOS regulon expression may have depended upon the activation of iNOS (55) rather than hypoxia. When mouse macrophages cultured in 21% oxygen were infected with

Mtb *in vitro*, *DOS* regulon induction depended on expression of iNOS (21, 37). In contrast, in the present studies, exposure of human macrophages to Mtb over an interval long enough to allow for phagocytosis but too brief to allow for nitrite accumulation led to marked induction of the *DOS* regulon in the absence of any NOS. Therefore, to the extent that *DOS* regulon induction is an indicator of hypoxia, human macrophages as cultured here appear to achieve a greater degree of functional intracellular hypoxia *in vitro* than mouse macrophages as usually cultured. The same may be true *in vivo*, insofar as iNOS expressed in mouse macrophages during Mtb infection appears to access sufficient oxygen to restrain Mtb growth (3, 37). Therefore, experiments with NarG-deficient Mtb in the mouse may not reflect the functional relevance of mycobacterial nitrate respiration in the human host.

RNS can act as double-edged swords, depending on their level (56, 57). The present results bring this truism into sharp relief with respect to Mtb. High levels of RNS can kill Mtb, but low levels may do the pathogen a service. First, nitrite of host or mycobacterial origin may induce genes with whose help Mtb can withstand other host stresses. Second, nitrite slows or stops Mtb's aerobic replication. Mtb may thus avoid killing a host that has not yet destroyed enough lung tissue to provide Mtb with a path back to the air. Third, nitrite increases intrabacterial ATP. Once Mtb is expelled from the host within aerosolized droplets, this reserve of metabolic energy may facilitate colonization of new hosts. Drugs that inhibit Mtb's defenses against RNS may therefore be useful even in settings where immunodeficiency impairs the expression of host iNOS or hypoxia limits its action.

#### Methods

#### Isolation and differentiation of primary human monocytes

Heparinized peripheral blood was collected by venepuncture from healthy human donors who provided informed consent under an IRB approved protocol. Monocytes were isolated and differentiated as described (29, 34). Briefly, mononuclear cells isolated by centrifugation over Ficoll-Paque (GE Healthcare) were subject to positive selection using magnetic beads coupled to anti-CD14 antibodies (Milteyi Biotec) and cultured in 60% RPMI, supplemented with 1% glutamax, 40% human plasma and GM-CSF and TNF $\alpha$  (0.5 ng ml<sup>-1</sup> each) at 5x10<sup>5</sup> cells ml<sup>-1</sup>. 30% of the total culture volume was replaced with fresh medium and cytokines twice a week. For Cytodex1 bead culture, Cytodex 1 beads were prepared according to the instructions provided by the manufacturer (GE Healthcare). The cells were plated at a density of  $2x10^6$  cells ml<sup>-1</sup> per well of a 24 well plate and 50% of the medium was replaced three times a week. After 2 weeks at 10% or 21% O<sub>2</sub> and 5% CO<sub>2</sub> at 37°C in a humidified atmosphere, the cells were activated with IFN $\gamma$  (5 ng ml<sup>-1</sup>) and infected with Mtb the following day. No antibiotics were used at any point.

To remove most nitrate from the medium, 1 day prior to infection the cells were washed 3 times with PBS and the medium was replaced with DMEM with 10% human plasma, GM-CSF, 0.5 ng mL<sup>-1</sup> TNF $\alpha$  and 5 ng mL<sup>-1</sup> IFN $\gamma$ .

#### Measurement of nitrite

Nitrite levels in the supernatants of macrophage cultures were measured by the Griess assay (41, 56) using nitrite standards prepared in the same medium used to culture the macrophages.

#### Preparation of Mtb

M. tuberculosis H37Rv was grown in Middlebrook 7H9 broth supplemented with 0.2% glycerol, 0.5% BSA, 0.2% dextrose and 0.085% NaCl (ADNaCl) with 0.05% Tween 80. The *narG*, *nirB* and complemented strains (36, 42) were generously provided by Dr. F. Bange. Single-cell suspensions were collected in the supernatant after centrifugation at 120 G for 10 minutes. For macrophage infection, ~2x10<sup>7</sup> bacteria were pelleted by centrifugation and resuspended in PBS. CFUs were determined by serial dilution in 0.1% TritonX-100, and plating on Middlebrook 7H11 agar with 10% oleic acid-albumin-dextrose-catalase (OADC) enrichment (Difco, BD) supplemented with 0.5% glycerol for 2-3 weeks at 37°C, 5% CO<sub>2</sub>.

#### *Pimonidazole staining*

Uninfected or Mtb-infected (MOI: 5) macrophages were treated with or without 200  $\mu$ M pimonidazole for 24 hours, washed and stained antipimonidazole antibody coupled to allophycocyanin (APC) (Hypoxyprobe RedAPC Kit) as recommended by the manufacturer and counterstained with the nuclear stain 4',6-diamidino-2-phenylindole (DAPI). APC and DAPI signals were quantified at the Cell Screening Core Facility, Weill Cornell Medical College and analyzed using MetaXpress High Content Image Acquisition & Analysis Software (Molecular Devices). The APC signal was normalized to the number of nuclei per image.

#### Intrabacterial ATP

Single-cell suspensions of Mtb ( $7x10^8$  per condition) were incubated in 7H9/ADNaCl with 0.05% Tween 80 at an OD<sub>580</sub> of 0.07 in 1% O<sub>2</sub>, 5% CO<sub>2</sub> at 37 °C in a humidified atmosphere, pelleted by centrifugation, washed once in PBS, pelleted and processed immediately for ATP measurement or frozen at -80°C for later use. For a single endpoint measurement, the pellet was resuspended in 220  $\mu$ l of standard lysis buffer: 100 mM Tris-OH and 1 mM PMSF, pH 7.4, with 1x protease inhibitor (Roche: cOmplete ULTRA Tablets, Mini, EDTA-free, EASY pack catalogue number: 05892791001) added immediately prior to bacterial lysis in a bead-beating homogenizer with silica beads for 2 cycles with incubation on ice between them. After brief centrifugation to pellet silica beads, 1:1:2 (v/v/v) ratio of cell lysate (50  $\mu$ L), lysis buffer and ATP reagent were added in triplicate to a black 96-well opaque plate for the BacTiter-Glo Microbial Cell Viability Assay (Promega). immediately using Luminescence was read а SpectraMax L Microplate Reader (Molecular Devices). For rate Luminescence measurements, the lysate was filter-sterilized (0.22 µm Corning Costar Spin-X Plastic Centrifuge Tube Filters at 10,000G for 20-30 minutes). ATP was measured in an aliquot and reagent ATP was added to the remaining lysate from which triplicate 10 µL aliquots were immediately dispensed to the assay plate. At the indicated times, 90  $\mu$ L of lysis buffer and 100 µL of ATP reagent were added and luminescence measured. ATP standards diluted in lysis buffer were prepared fresh for each experiment. The intrabacterial ATP for each sample was normalized to the protein concentration of each sample (Bio-Rad Protein Assay).

#### Quantitative RT-PCR

Macrophages ( $5x10^6$  per T25 flask) were cultured in 10% or 21% oxygen and infected with Mtb (MOI: 40) for 10 hours. The monolayer was washed twice with PBS. Trizol (2 mL) was added to each flask and the cells detached and using a rubber scraper. Alternatively,  $\geq 2 \times 10^9$  bacteria were incubated per condition in 1% or 21% oxygen for 3 days in standard medium, in the absence of nitrate. An equal volume of buffer containing 5 M guanidinium thiocyanate, 25 mM sodium citrate, 20 mM N-laurylsarcosine, 0.7% v/v  $\beta$ -mercaptoethanol was added. The bacteria were pelleted by centrifugation, 1 mL of Trizol added per sample and the suspension beaten with silica beads. RNA extraction was conducted using an RNeasy kit (Qiagen) in accordance with the manufacturer's instructions, except that off-column DNAse digestion was performed for 2 hrs at 37 °C. One mg of total RNA was reverse-transcribed using the GeneAmp RNA PCR Kit (Applied Biosystems). Quantitative RT-PCR was performed using gene specific primers (TaqMan Gene Expression Assay, Life Technologies) and the SuperScript III Platinum Two Step qRT-PCR Kit (Life Technologies) with a 7900HT Fast Real Time PCR System (Applied Biosystems). Each experiment was performed in triplicate. Values within the linear range of the primers were normalized to values for 16S rRNA.

#### RNA Sequencing

At least  $5x10^9$  bacteria per sample were incubated in 1% oxygen and left untreated or treated with 5 mM nitrate or 2.5 mM nitrite. After 3 days RNA was collected from one flask of untreated Mtb. The remaining flasks (untreated or nitrate- or nitrite- treated) were transferred to 21% oxygen for an additional overnight incubation before RNA collection. RNA samples with an RNA Integrity Number (RIN) (Bioanalyzer (Agilent Technologies 2100) value > 8 were processed further. Ribosomal RNA hybridization with magnetic was removed by bead-coupled oligonucleotides (MicrobExpress Kit, Life Technologies). Libraries were prepared according to the manufacturer's instructions and resultant RNA sequenced (HiSeq2000/1000, Illumina). Single-ended sequencing reads were aligned to the reference genome of Mtb H37Rv using the Burrows-Wheeler Alignment tool (20, 50) (Accession NC\_000962.3). Cufflinks (53) was used to measure transcript abundances in reads per kilobase of exon per million mapped reads (RPKM) as well as to find differentially expressed genes. Regulation was considered significant for changes with a p value corrected for multiple comparisons ( $p_{corr}$ ) < 0.05 (Supplementary Tables 1 and 2, see Appendix 1). The samples labeled "nitrate" and "untreated," were prepared in 3 independent experiments. Only one sample was available for the "nitrite" and "untreated 1% oxygen" conditions. These samples were only used to evaluate the expression all 47 genes of the DOS regulon as a unit, with the expression of each gene serving as a control for the expression of the others.

## Statistical analysis

Statistical analysis was performed as indicated in the figure legends using Prism 5.0f for Macintosh (GraphPad Software, San Diego, CA).

### References

- 1. Nathan C, Shiloh MU (2000) Reactive oxygen and nitrogen intermediates in the relationship between mammalian hosts and microbial pathogens. *Proc Natl Acad Sci USA* 97:8841–8848.
- 2. MacMicking J, Xie QW, Nathan C (1997) Nitric oxide and macrophage function. *Annu Rev Immunol* 15:323–350.
- 3. MacMicking JD et al. (1997) Identification of nitric oxide synthase as a protective locus against tuberculosis. *Proc Natl Acad Sci USA* 94:5243–5248.
- 4. Chan J, Xing Y, Magliozzo RS, Bloom BR (1992) Killing of virulent *Mycobacterium tuberculosis* by reactive nitrogen intermediates produced by activated murine macrophages. *J Exp Med* 175:1111–1122.
- 5. Mishra BB et al. (2013) Nitric oxide controls the immunopathology of tuberculosis by inhibiting NLRP3 inflammasome-dependent processing of IL-1β. *Nat Immunol* 14:52–60.
- 6. Nicholson S et al. (1996) Inducible nitric oxide synthase in pulmonary alveolar macrophages from patients with tuberculosis. *J Exp Med* 183:2293–2302.
- 7. Choi H-S, Rai PR, Chu HW, Cool C, Chan ED (2002) Analysis of nitric oxide synthase and nitrotyrosine expression in human pulmonary tuberculosis. *Am J Respir Crit Care Med* 166:178–186.
- 8. Mattila JT et al. (2013) Microenvironments in tuberculous granulomas are delineated by distinct populations of macrophage subsets and expression of nitric oxide synthase and arginase isoforms. *J Immunol* 191:773–784.
- 9. Schön T et al. (2003) Arginine as an adjuvant to chemotherapy improves clinical outcome in active tuberculosis. *Eur Respir J* 21:483–488.
- 10. Schön T et al. (2011) Effects of a food supplement rich in arginine

in patients with smear positive pulmonary tuberculosis--a randomised trial. *Tuberculosis (Edinb)* 91:370–377.

- 11. Gusarov I, Shatalin K, Starodubtseva M, Nudler E (2009) Endogenous nitric oxide protects bacteria against a wide spectrum of antibiotics. *Science* 325:1380–1384.
- 12. McCollister BD, Hoffman M, Husain M, Vázquez-Torres A (2011) Nitric oxide protects bacteria from aminoglycosides by blocking the energy-dependent phases of drug uptake. *Antimicrob Agents Chemother* 55:2189–2196.
- 13. van Sorge NM et al. (2013) Methicillin-resistant *Staphylococcus aureus* bacterial nitric-oxide synthase affects antibiotic sensitivity and skin abscess development. *Journal of Biological Chemistry* 288:6417–6426.
- 14. Winter SE et al. (2013) Host-Derived Nitrate Boosts Growth of *E. coli* in the Inflamed Gut. *Science* 339:708–711.
- 15. Aly S et al. (2006) Oxygen status of lung granulomas in *Mycobacterium tuberculosis*-infected mice. *J Pathol* 210:298–305.
- 16. Via LE et al. (2008) Tuberculous granulomas are hypoxic in guinea pigs, rabbits, and nonhuman primates. *Infect Immun* 76:2333–2340.
- 17. Sohaskey CD, Wayne LG (2003) Role of *narK2X* and *narGHJI* in hypoxic upregulation of nitrate reduction by *Mycobacterium tuberculosis*. *Journal of Bacteriology* 185:7247–7256.
- Wayne LG, Doubek JR (1965) Classification and identification of mycobacteria. II. Tests employing nitrate and nitrite as substrate. *Am Rev Respir Dis* 91:738–745.
- Giffin MM, Raab RW, Morganstern M, Sohaskey CD (2012) Mutational analysis of the respiratory nitrate transporter NarK2 of *Mycobacterium tuberculosis*. *PLoS ONE* 7:e45459.
- 20. Sherman DR et al. (2001) Regulation of the *Mycobacterium tuberculosis* hypoxic response gene encoding alpha -crystallin. *Proc Natl Acad Sci USA* 98:7534–7539.

- 21. Voskuil MI et al. (2003) Inhibition of respiration by nitric oxide induces a *Mycobacterium tuberculosis* dormancy program. *Journal of Experimental Medicine* 198:705–713.
- 22. Wayne LG, Hayes LG (1998) Nitrate reduction as a marker for hypoxic shiftdown of *Mycobacterium tuberculosis*. *Tuber Lung Dis* 79:127–132.
- 23. Fenhalls G et al. (2002) *In situ* detection of *Mycobacterium tuberculosis* transcripts in human lung granulomas reveals differential gene expression in necrotic lesions. *Infect Immun* 70:6330–6338.
- 24. Rachman H et al. (2006) Unique transcriptome signature of *Mycobacterium tuberculosis* in pulmonary tuberculosis. *Infect Immun* 74:1233–1242.
- 25. Tan MP et al. (2010) Nitrate respiration protects hypoxic *Mycobacterium tuberculosis* against acid- and reactive nitrogen species stresses. *PLoS ONE* 5:e13356.
- 26. Sohaskey CD (2008) Nitrate enhances the survival of *Mycobacterium tuberculosis* during inhibition of respiration. *Journal of Bacteriology* 190:2981–2986.
- 27. Goh KS, Rastogi N, Berchel M, Huard RC, Sola C (2005) Molecular evolutionary history of tubercle bacilli assessed by study of the polymorphic nucleotide within the nitrate reductase (narGHJI) operon promoter. *J Clin Microbiol* 43:4010–4014.
- 28. Lundberg JO, Weitzberg E, Gladwin MT (2008) The nitrate–nitrite– nitric oxide pathway in physiology and therapeutics. *Nat Rev Drug Discov* 7:156–167.
- 29. Vogt G, Nathan C (2011) *In vitro* differentiation of human macrophages with enhanced antimycobacterial activity. *J Clin Invest* 121:3889–3901.
- 30. Arteel GE, Thurman RG, Raleigh JA (1998) Reductive metabolism of the hypoxia marker pimonidazole is regulated by oxygen tension independent of the pyridine nucleotide redox state. *Eur J Biochem*

253:743-750.

- 31. Rowe JJ, Yarbrough JM, Rake JB, Eagon RG (1979) Nitrite inhibition of aerobic bacteria. *Current Microbiology* 2:51–54.
- 32. Malm S et al. (2009) The roles of the nitrate reductase NarGHJI, the nitrite reductase NirBD and the response regulator GlnR in nitrate assimilation of *Mycobacterium tuberculosis*. *Microbiology* 155:1332–1339.
- 33. Ehrt S, Schnappinger D (2009) Mycobacterial survival strategies in the phagosome: defence against host stresses. *Cell Microbiol* 11:1170–1178.
- 34. Rodriguez GM, Voskuil MI, Gold B, Schoolnik GK, Smith I (2002) ideR, an essential gene in *Mycobacterium tuberculosis*: role of IdeR in iron-dependent gene expression, iron metabolism, and oxidative stress response. *Infect Immun* 70:3371–3381.
- 35. Rohde KH, Abramovitch RB, Russell DG (2007) *Mycobacterium tuberculosis* invasion of macrophages: linking bacterial gene expression to environmental cues. *Cell Host Microbe* 2:352–364.
- 36. Voskuil MI, Bartek IL, Visconti K, Schoolnik GK (2011) The response of *Mycobacterium tuberculosis* to reactive oxygen and nitrogen species. *Front Microbiol* 2:105.
- 37. Schnappinger D et al. (2003) Transcriptional adaptation of *Mycobacterium tuberculosis* within macrophages: insights into the phagosomal environment. *J Exp Med* 198:693–704.
- Alderton WK, Cooper CE, Knowles RG (2001) Nitric oxide synthases: structure, function and inhibition. *Biochem J* 357:593– 615.
- 39. McCormick CC, Li WP, Calero M (2000) Oxygen tension limits nitric oxide synthesis by activated macrophages. *Biochem J* 350 Pt 3:709–716.
- 40. James PE, Grinberg OY, Michaels G, Swartz HM (1995) Intraphagosomal oxygen in stimulated macrophages. *J Cell Physiol*

163:241-247.

- 41. Ding AH, Nathan CF, Stuehr DJ (1988) Release of reactive nitrogen intermediates and reactive oxygen intermediates from mouse peritoneal macrophages. Comparison of activating cytokines and evidence for independent production. *J Immunol* 141:2407–2412.
- 42. Stermann M, Bohrssen A, Diephaus C, Maass S, Bange F-C (2003) Polymorphic nucleotide within the promoter of nitrate reductase (NarGHJI) is specific for *Mycobacterium tuberculosis*. *J Clin Microbiol* 41:3252–3259.
- Bryan NS et al. (2005) Nitrite is a signaling molecule and regulator of gene expression in mammalian tissues. *Nat Chem Biol* 1:290– 297.
- 44. Wright RO, Lewander WJ, Woolf AD (1999) Methemoglobinemia: etiology, pharmacology, and clinical management. *Ann Emerg Med* 34:646–656.
- 45. Reddy D, Lancaster JR, Cornforth DP (1983) Nitrite inhibition of *Clostridium botulinum*: electron spin resonance detection of ironnitric oxide complexes. *Science* 221:769–770.
- 46. Kumar A, Toledo JC, Patel RP, Lancaster JR, Steyn AJC (2007) Mycobacterium tuberculosis DosS is a redox sensor and DosT is a hypoxia sensor. *Proc Natl Acad Sci USA* 104:11568–11573.
- 47. Weinberg ED (1974) Iron and susceptibility to infectious disease. *Science* 184:952–956.
- 48. Harris AB (1967) Inhibition of growth and nucleic acid synthesis in iron-deficient Mycobacterium smegmatis. *J Gen Microbiol* 47:111–119.
- 49. De Voss JJ, Rutter K, Schroeder BG, Barry CE (1999) Iron acquisition and metabolism by mycobacteria. *Journal of Bacteriology* 181:4443–4451.
- 50. Li H, Durbin R (2010) Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* 26:589–595.

- Dlugovitzky D et al. (2000) Influence of disease severity on nitrite and cytokine production by peripheral blood mononuclear cells (PBMC) from patients with pulmonary tuberculosis (TB). *Clin Exp Immunol* 122:343–349.
- 52. Lamsal M et al. (2007) Evaluation of lipid peroxidation product, nitrite and antioxidant levels in newly diagnosed and two months follow-up patients with pulmonary tuberculosis. *Southeast Asian J Trop Med Public Health* 38:695–703.
- 53. Trapnell C et al. (2013) Differential analysis of gene regulation at transcript resolution with RNA-seq. *Nat Biotechnol* 31:46–53.
- 54. Tsai MC et al. (2006) Characterization of the tuberculous granuloma in murine and human lungs: cellular composition and relative tissue oxygen tension. *Cell Microbiol* 8:218–232.
- 55. Shi L et al. (2005) Changes in energy metabolism of Mycobacterium tuberculosis in mouse lung and under in vitro conditions affecting aerobic respiration. *Proc Natl Acad Sci USA* 102:15629–15634.
- 56. Rohde K, Yates RM, Purdy GE, Russell DG (2007) *Mycobacterium tuberculosis* and the environment within the phagosome. *Immunol Rev* 219:37–54.
- 57. Nathan C (1992) Nitric oxide as a secretory product of mammalian cells. *FASEB J* 6:3051–3064.

#### **CHAPTER 3**

# NITRITE IMPACTS THE SURVIVAL OF *MYCOBACTERIUM TUBERCULOSIS* IN RESPONSE TO ISONIAZID AND HYDROGEN PEROXIDE \*

#### Summary

When access to molecular oxygen is restricted, *Mycobacterium tuberculosis* (Mtb) can respire an alternative electron acceptor, nitrate. We found that Mtb within infected primary human macrophages *in vitro* at physiologic tissue oxygen tensions respired nitrate, generating copious nitrite. A strain of Mtb lacking a functioning nitrate reductase was more susceptible than wild type Mtb to treatment with isoniazid during infection of macrophages. Likewise, nitrate reductase-deficient Mtb was more susceptible to isoniazid than wild type Mtb in axenic culture, and more resistant to hydrogen peroxide. These phenotypes were reversed by the addition of exogenous nitrite. Further investigation suggested that nitrite might inhibit the bacterial catalase. To the extent that Mtb itself is the most relevant source of nitrite acting within Mtb, these findings suggest that inhibitors of Mtb's nitrate transporter or nitrate reductase could enhance the efficacy of isoniazid.

#### Introduction

Nitrate  $(NO_3^{-})$  and nitrite  $(NO_2^{-})$  arise in the mammalian host from three major sources—dietary, bacterial and host. Dietary nitrate is absorbed into the blood. Some is reduced by oral and gastrointestinal

<sup>&</sup>lt;sup>\*</sup> Amy Cunningham-Bussel, Franz-Christoph Bange, and Carl Nathan, "Nitrite Impacts the Survival of *Mycobacterium Tuberculosis* In Response to Isoniazid and Hydrogen Peroxide," *MicrobiologyOpen* (*Manuscript in press*)

commensals, producing nitrite, which is also absorbed. Nitric oxide synthases (NOSs), including those that are constitutively expressed, produce nitric oxide, which autoxidizes to nitrate and nitrite. Even in healthy individuals, in whom relatively little of the high-output NOS isoform (iNOS) is expressed (1), nitrate is present in plasma at levels of 20-40  $\mu$ M (2). In patients with tuberculosis (TB), iNOS and other NOSs are expressed in granulomas (1, 3).

At oxygen tensions of 1% or lower, Mtb itself can produce nitrite as it maintains ATP synthesis and redox homeostasis by reducing an alternative electron acceptor, nitrate. Mtb constitutively expresses the *narGHJI* operon, which encodes Mtb's only functional nitrate reductase (4, 5). However, transport of nitrate into the bacterium depends on transcription of a nitrate transporter encoded by *narK2*, a member of the dormancy survival (*DOS*) regulon (6-8). The *narK2* gene is expressed in axenically cultured Mtb only when oxygen levels fall below 1%. Induction of *narK2* markedly enhances mycobacterial nitrate respiration (4). RNA transcripts from *narG*, which encodes a subunit of the nitrate reductase, and *narX*, an inactive fused nitrate reductase downstream of *narK2* in a shared operon, have been detected in lungs of patients with TB (4, 9, 10), and RNA transcripts encoding the nitrate transporter, *narK2*, were elevated in the lungs of Mtb infected mice (11). Thus, mycobacterial nitrate reduction likely occurs in the infected host.

Our laboratory recently converted our culture system for primary human macrophages from conventional gas-phase conditions of 21%  $O_2$ , 5%  $CO_2$  to 10%  $O_2$ , 5%  $CO_2$  on the grounds that 10%  $O_2$  is physiologic for tissue macrophages (12). To our surprise, infection of the macrophages *in vitro* with wild type Mtb under 10% oxygen resulted in extensive accumulation of nitrite in the supernatant. Evidence will be presented elsewhere that macrophages were not the source of nitrite in this system. In contrast, as shown below, Mtb's NarG was the source, despite the non-hypoxic level of oxygen in the gas phase.

Wayne has proposed that nitrate respiration supports the establishment of a non-replicating persistent state (NRP) in response to hypoxia (13). We hypothesized that Mtb lacking the capacity for nitrate respiration might demonstrate an impaired entrance into NRP and therefore remain more susceptible to drugs that target replicating Mtb. The goal of the present study was to assess the contribution of nitrate respiration to Mtb's sensitivity to the drugs isoniazid (INH), ethambutol, streptomycin and rifampicin. Compared to wild type Mtb, *narG*-deficient Mtb, which is unable to produce nitrite, was selectively hyper-susceptible to INH, as well as hyper-resistant to hydrogen peroxide. INH requires oxidative activation by the mycobacterial catalase/peroxidase KatG in order become cidal. The resultant isonicotinoyl radical adducts with cellular pyridine nucleotides and potently inhibits InhA, a member of type II fatty acid synthesis pathway involved in cell wall mycolic acid biosynthesis (14, 15). These findings imply that the action of INH on Mtb may be blunted by nitrite contributed by the host, the pathogen or both.

#### Results

Nitrate respiration enhanced mycobacterial resistance to isoniazid within primary human macrophages.

We observed no difference in survival of wild type (Wt Mtb) and

*narG*-deficient Mtb (NarG Mtb) during infection of primary human macrophages under 10%  $O_2$  and 5%  $CO_2$  (Figure 1a). During the course of infection, large quantities of nitrite arose in the supernatant of macrophages infected with wild type Mtb, which was not observed in the supernatants of macrophages infected with *narG*-deficient Mtb (Figure 1b). To test the role of nitrate reduction on the susceptibility of Mtb to antibiotics, we infected macrophages with wild type, *narG*-deficient and complemented strains (NarGc Mtb) and treated the cultures with drugs. Mtb deficient in *narG* was hyper-susceptible to INH, with a 10-fold decrease in the MBC99 compared to wild type (Figure 1c-d). In contrast, *narG* made no observable difference to the antibacterial actions of streptomycin, rifampicin or ethambutol on Mtb within macrophages (Supplementary Figure 1a-f, see Appendix 2).
#### INFECTED HUMAN MACROPHAGES



Figure 1. Effect of nitrate respiration on mycobacterial resistance to isoniazid within primary human macrophages. (A) Survival of wild type and *narG*-deficient (NarG) Mtb within macrophages (MOI: 0.1, corresponding to 10,000 bacteria). (B) Nitrite levels arising in the supernatant of macrophages 3 days after infection with wild type Mtb (Wt Mtb), NarG Mtb and the complemented strain (NarGc Mtb) (MOI: 5). (C) Susceptibility of Mtb to INH within infected macrophages (MOI: 0.1) over 3 days. Means ±SEM from one experiment representative of 3 independent experiments using cells from different donors. (D) As in (C), pooling results from the 3 donors (means ± SEM) at a single INH concentration administered to the culture for 3 days (0.1 µg mL<sup>-1</sup>). The p value was determined by an unpaired t test.

Addition of nitrite reversed the hyper-susceptibility of narG-deficient Mtb to isoniazid.

The hyper-susceptibility of *narG*-deficient Mtb to INH may have resulted from loss of the metabolic energy generated by nitrate reduction from failure to produce nitrite. To distinguish between these or possibilities, we added exogenous nitrite to the medium of infected macrophage cultures treated with INH. Because the standard culture medium consisted of 40% human plasma and 60% RPMI 1640, it contained high concentrations of nitrate. Therefore, to limit the accumulation of nitrite produced by wild type Mtb, we replaced the medium immediately prior to infection with a low-nitrate medium (to minimize perturbation we refrained from washing the cells thoroughly; thus, medium exchange was not complete). Addition of exogenous nitrite enhanced bacterial survival in the presence of INH in a dose-dependent manner (Figure 2a-c). That this effect was less striking in the case of wild type Mtb was likely due to residual nitrate in the medium, allowing for low-level nitrite production (Figure 2a).

INFECTED HUMAN MACROPHAGES



Figure 2. Impact of exogenous nitrite on the hyper-susceptibility of narG-deficient Mtb to isoniazid. (A, B) The effect of nitrite on the survival of (A) wild type or (B) *narG*-deficient (NarG) Mtb within INH-treated infected macrophages (MOI: 0.1). The standard culture medium was replaced with low-nitrate medium. Where indicated by the dashed lines, exogenous sodium nitrite (final concentration 1 mM) was added to the culture medium along with the indicated concentrations of INH. CFU were harvested 3 days later. (C) Survival of wild type and *narG*-deficient (NarG) Mtb within macrophages treated with increasing concentrations of nitrite in the presence of a single concentration of INH (0.1  $\mu$ g mL<sup>-1</sup>). The standard culture medium was replaced with low-nitrate medium and exogenous sodium nitrite was added at the indicated final concentrations. Each result is representative of at least two independent experiments (means ± SEM).

Mtb deficient in narG Mtb was hyper-susceptible to isoniazid in axenic culture.

Mtb respires nitrate when infecting human macrophages under 10% oxygen, but in axenic culture, as noted, nitrate respiration is barely detectable until oxygen is lowered to 1% or less. This implied that within macrophage phagosomes, the functional level of oxygen perceived by

Mtb was far lower than the level present in the gas phase. The nitrate reductase assay is routinely used in clinical laboratories to identify drug resistant Mtb (16). Although the assay is carried out under room air (21%)oxygen), nitrate reductase activity is measured at the end of a prolonged incubation period, which may indicate that the medium in which Mtb is incubated becomes hypoxic. Alternatively, while aerobic cultures of Mtb do not express the nitrate transporter, *narK2*, nitrate may gradually diffuse across the plasma membrane and be reduced to nitrite. In order to test the role of nitrate respiration in Mtb's susceptibility to INH in the absence of macrophages, we cultured wild type and *narG*-deficient Mtb in 21% oxygen or 1% oxygen. No measurable survival differences were observed for the two strains in room air in response to isoniazid, streptomycin, or rifampicin (Figure 3a-c). Under 1% oxygen, however, the survival of *narG*-deficient INH-treated Mtb dropped by 2-3  $\log_{10}$  as compared to wild type and complemented strains (Figure 3d), which was due to the induction of the nitrate reductase system. Next, we examined survival as a function of time rather than concentration. A single application of INH led to a drop of  $2 \log_{10}$  in CFU of the *narG*-deficient strain over 9 days, while the CFU of the wild type and complemented strains remained constant (Figure 3e). Exogenous nitrite enhanced the survival of the *narG*-deficient and wild type strains cultured in the absence of nitrate and treated with INH (Figure 3f).

Following INH treatment and its subsequent activation by KatG, Mtb induces the expression of *kasA* and *ahpC* (17). The *narG*-deficient strain demonstrated greater induction of *kasA* and *ahpC* RNA transcripts than the wild type strain in response to 0.1  $\mu$ g mL<sup>-1</sup> INH, and induction was attenuated following treatment with nitrite (Figure 3g). INH requires oxidative activation by the mycobacterial catalase/peroxidase KatG in order to become bioactive. These results suggest that when presented with the same concentration of INH under conditions that could sustain nitrate respiration by wild type Mtb, *narG*-deficient Mtb converted a larger proportion of the INH prodrug to its active form than wild type Mtb. This implied that the production of nitrite during respiration of nitrate by wild type Mtb might partially inhibit KatG.

Figure 3. Impact of narG-deficiency on susceptibility of Mtb to isoniazid in axenic culture. (A-C) Survival of wild type and narGdeficient Mtb in the presence of the indicated concentrations of (A) isoniazid, (B) rifampicin or (C) streptomycin in 21% oxygen over 5 days, initial (OD 0.1). Published MICs are provided for reference (46). (D, E) Survival of wild type, *narG*-deficient (NarG) and complemented (NarGc) strains cultured in 1% oxygen (D) for 3 days with the indicated concentrations of INH or (E) as a function of time over 9 days with 0.1 µg  $mL^{-1}$  INH. (F) Survival of wild type and *narG*-deficient Mtb incubated with increasing concentrations of exogenous nitrite or one fixed dose of nitrate, as indicated, over 3 days in 1% oxygen with 0.1  $\mu$ g mL<sup>-1</sup> INH. Results for (A-F) are means  $\pm$  SEM representative of 2 independent experiments. (G) Expression of transcripts of *ahpC* and *kasA* by wild type and *narG*-deficient Mtb treated with 0.1  $\mu$ g mL<sup>-1</sup> INH and 5 mM nitrate for 16 hrs at 1% oxygen relative to 16S RNA. Where indicated, 2.5 mM nitrite was also added. Bars indicate means  $\pm$  SEM from 3 independent experiments analyzed by unpaired t tests.

## Figure 3 (continued)

BROTH CULTURE OF MTB IN THE ABSENCE OF HUMAN MACROPHAGES





narG-deficient Mtb was not hyper-susceptible to treatment with ethionamide.

To further investigate the role of KatG in mediating the hypersusceptibility of *narG*-deficient Mtb to INH, we infected human macrophages with Mtb and treated the cultures with ethionamide (ETH). ETH and INH are close chemical analogues. Both are prodrugs that when activated inhibit the same target, InhA. However, ETH is activated by the monoxygenase EtaA rather than by KatG (18, 19). Survival of wild type and *narG*-deficient Mtb treated with ETH within infected human macrophages did not differ (Figure 4a). Therefore, the INH susceptibility of *narG*-deficient Mtb is unlikely to be due to differential dependence on InhA and may instead be due to differences in the activation of INH by KatG.

#### Nitrate reduction enhanced the lethality of hydrogen peroxide.

KatG is the mycobacterial catalase peroxidase that is also responsible for the cellular activation of INH. Levels of KatG mRNA determined by qRT-PCR and KatG protein detected by western blot did not differ among wild type, *narG*-deficient and complemented strains of Mtb (Supplementary figure 2b-c, see Appendix 2). Moreover, we were unable to measure differences in the extent of KatG dependent oxidation of INH via nitroblue tetrazolium staining of polyacrylamide gel electrophoregrams of lysates obtained from the three strains (data not shown). Nonetheless, to help assess whether the catalytic activity of KatG might differ between wild type and *narG*-deficient Mtb, the strains were incubated for several hours with nitrite or nitrate at 1% oxygen to allow for nitrate respiration and then treated with hydrogen peroxide, the natural substrate of KatG. We reasoned that if the activity of KatG were attenuated by nitrite, the wild type and complemented strains might be more susceptible to hydrogen peroxide than *narG*-deficient Mtb. In accord with this hypothesis, *narG*-deficient Mtb was more resistant to hydrogen peroxide than the other two strains (Figure 4b). The enhanced mycobactericidal effect of hydrogen peroxide on wild type Mtb incubated with nitrate was also observed following the addition of increasing concentrations of exogenous nitrite in the absence of nitrate (Figure 4c and Supplementary Figure 2b, see Appendix 2).





Figure 4. Impact of narG-deficiency on susceptibility of Mtb to ethionamide and hydrogen peroxide. (A) Survival of wild type and *narG*-deficient Mtb within infected primary human macrophages (MOI: 0.1) treated with the indicated concentrations of ethionamide for 3 days at 10% oxygen. (B) Survival of Mtb in axenic culture treated with nitrate or nitrite for 8 hours in 1% oxygen, to allow for nitrate respiration to occur, and then treated with hydrogen peroxide (500  $\mu$ M) and incubated in 1% oxygen overnight (initial OD<sub>580</sub>: 0.1). These are pooled data from 2 independent experiments and were analyzed by an unpaired t test. (C) Survival of wild type, *narG*-deficient (NarG Mtb) and complemented strains (NarGc Mtb) (initial OD580: 0.1) following treatment with both hydrogen peroxide (500  $\mu$ M) and increasing concentrations of nitrite as indicated. Individual experiments representative of at least 2 independent experiments. Bars indicate means ± SEM.

#### Discussion

we report that deficiency of *narG* increases Herein the susceptibility of Mtb to INH and decreases its susceptibility to hydrogen peroxide. Prior to this work, the knockout of only one gene, cydC, had been reported to increase the susceptibility of Mtb to INH (20). The mechanism in the present study most likely involves nitrite-mediated impairment of the activity of KatG that takes place when wild type Mtb reduces nitrate to nitrite, which is absent in *narG*-deficient Mtb. Although nitrate is usually omitted from mycobacterial culture media, it is a physiologic constituent of human body fluids. Although Mtb does not reduce nitrate to nitrite in axenic culture unless the culture is hypoxic, we found that Mtb does reduce nitrate to nitrite when cultured within macrophages at physiologic oxygen tensions. Thus, phenotypic INH resistance arising from Mtb's nitrate respiration is likely to occur in the Mtb-infected host.

Interactions of catalase and other peroxidases with nitrite are complex (21-24). When we incubated Mtb with nitrite, lysed the cells, separated the proteins by non-reducing polyacrylamide gel electrophoresis and incubated the gel with INH and a tetrazolium dye as described (25), we did not see impairment in the activity of catalase compared to lysates from Mtb not exposed to nitrite (not shown). However, the biochemical environment within intact mycobacteria differs markedly from that of gels in room air, and tetrazolium is not a physiologic oxidant for INH. Moreover, *katG* mutations that cause INH resistance do not necessarily impact the rate of INH-NAD adduct formation by KatG relative to wild type Mtb, pointing to the potential for dissociation between the functional effects of KatG mutations and the results of biochemical assays for KatG activity (26-29).

The following evidence supports the hypothesis that nitrite interfered with the INH-activating function of catalase in the intact mycobacterium: Mtb deficient in *narG* was both more susceptible to INH and more resistant to hydrogen peroxide than wild type Mtb. Second, *narG*-deficient Mtb was not hyper-susceptible to treatment with ETH. Third, compared to wild type and complemented strains, *narG*-deficient Mtb demonstrated higher expression of ahpC and kasA, two genes induced by treatment with INH, and their expression was reduced by the addition of nitrite to the cell culture. Taken together, these results suggest that Mtb lacking *narG* achieves greater KatG-mediated activation of INH than wild type Mtb. Lastly, in agreement with this hypothesis, narGdeficient Mtb better resisted treatment with hydrogen peroxide. The mutation of genes other than katG can also cause INH resistance. For example, mutations that impair the catalytic activities of Ndh, a type II NADH dehydrogenase, and MshA and MshC, which contribute to mycothiol biosynthesis, can cause INH resistance. However, inactivating mutations in any one of these genes also cause ETH resistance (30-32), which was not observed in this study.

Nitric oxide or nitrite at low pH, which generates nitric oxide (33), can potentiate the bactericidal activity of hydrogen peroxide towards *E*. *coli* (2, 21, 34, 35). However, to our knowledge, this is the first report that nitrite at neutral pH can synergistically enhance the bactericidal action of hydrogen peroxide. Production of hydrogen peroxide by phagocytic cells is an important component of host antibacterial defense

72

(36). Mycobacterial nitrate respiration may enhance the lethality of host reactive oxygen species *in vivo*. This may result from inhibition of KatG by nitrite or from inhibition of a variety of antioxidant defense proteins, including KatG, by products more reactive than nitrite to which nitrite can give rise. In *E. coli*, nitrate reductase is an important source of species reactive enough to nitrosylate proteins, as evidenced by an 80% reduction in protein S-nitrosylation in *narG*-deficient as compared to wild type bacteria (37-39). Rhee *et al.* identified 29 mycobacterial proteins that were S-nitrosylated when Mtb was treated with nitrite at low pH, including KatG (40). In addition, heme peroxidases such as KatG oxidize nitrite to more reactive species, such as nitrogen dioxide ( $^{\circ}NO_2$ ) and nitryl chloride ( $NO_2$ Cl) (21-24).

INH is widely reported to lose mycobactericidal activity when Mtb is cultured in conditions that prevent it from replicating. However, when the oxygen in an axenic, nitrate-containing culture of Mtb was reduced below 1%, *narG*-deficient Mtb, which was not replicating, became highly INH sensitive. We confirmed that effect and demonstrated its reversal with exogenous nitrite. To the extent that a relevant source of nitrite for Mtb residing in a macrophage in the human host may be Mtb itself, the question arises whether the cidality of INH for non-replicating *narG*-deficient Mtb could be phenocopied in non-replicating wild type Mtb by an inhibitor of Mtb's nitrate transporter or nitrate reductase. If so, such agents might significantly reduce the time required to cure latent TB with INH or active TB with INH-containing regimens, reducing the incidence of emergent drug resistance associated with incomplete treatment (41, 42). InhA is essential to aerobically and perhaps also hypoxically cultured

mycobacteria as two compounds that targeted both InhA as well as fatty acid synthase type 1 were cidal to non-replicating Mtb (43). Nevertheless, it remains unclear whether the reduced survival of *narG*-deficient Mtb treated with INH resulted only from the inactivation of InhA or by inhibition of additional mycobacterial target(s). A mechanistic understanding of the hyper-susceptibility of *narG*-deficient Mtb to INH may reveal new INH target(s) amenable to inhibition by other compounds.

#### Methods

#### *Isolation and differentiation of primary human monocytes*

Isolation of human monocytes and their differentiation into macrophages was as described (12). In brief, heparinized peripheral blood was collected by venipuncture from healthy human donors who provided informed consent under an IRB approved protocol. Peripheral blood mononuclear cells were first isolated by centrifugation of whole blood over Ficoll-Paque (GE Healthcare). The buffy coats were collected and monocytes were isolated by positive selection using magnetic beads conjugated to anti-CD14 antibodies (Milteyi Biotec). Following isolation, the human monocytes were plated at a density of 500,000 cells ml<sup>-1</sup> per well of a 96 well plate. The culture medium consisted of 60% RPMI 1640, supplemented with 1% glutamax, 40% human plasma and GMCF and TNF $\alpha$  (0.5 ng ml<sup>-1</sup> each). 30% of the total culture volume was replaced with fresh medium and cytokines every 3 to 4 days. Replacing fresh medium was conducted as rapidly as possible in room air before the cells were placed back in the low oxygen incubator. Differentiation of the

monocytes was conducted over 2 weeks in our standard medium at 10%  $O_2$  and 5%  $CO_2$  at 37°C in a humidified atmosphere in a chamber flushed with  $N_2$  under the control of a PRoOX sensor and ProCO<sub>2</sub> regulator (BioSpherix). They were then activated with IFN $\gamma$  (5 ng ml<sup>-1</sup>) before infection with *M. tuberculosis* the following day at the desired multiplicity of infection (MOI).

## Preparation of Mtb and infection of human macrophages with Mycobacterium tuberculosis

*M. tuberculosis* H37Rv was grown in Middlebrook 7H9 broth supplemented with 0.2% glycerol, 0.5% BSA, 0.2% dextrose and 0.085% NaCl with 0.05% Tween 80. The *narG*-deficient and complemented strains were generated as described (44). Cultures were started with 1 mL stock originally frozen at -80°C in log phase and then grown over 4 to 5 days to optical densities (OD) of 0.5-1.25 before the start of an experiment. A single-cell suspension was generated by centrifugation at 120 g for 10 minutes. For macrophages infection, roughly 200x10<sup>6</sup> bacteria were centrifuged, in order to be able to observe a pellet. The 7H9 was completely removed and the cells were washed with PBS. The bacteria were then resuspended in culture medium and the desired number of Mtb was added to the macrophage culture.

Certain experiments required that we remove the nitrate-containing medium from the cell culture. In this case, 1 day prior to infection the cells were washed with room temperature PBS 3 times and a low nitrate formulation of DMEM with 10% human plasma was added along with 0.5 ng mL<sup>-1</sup> GMCSF and TNF $\alpha$  and 5 ng mL<sup>-1</sup> IFN $\gamma$ .

#### *Measurement of nitrite*

Nitrite levels produced in the co-culture supernatants of infected macrophages were measured by the Griess assay. Briefly, 100  $\mu$ l of supernatant was removed and to this were added 50  $\mu$ l of 2% sulfanilamide with 5% phosphoric acid and 50  $\mu$ l of 0.2% N-1-napthylethylenediamine dihydrochloride. Nitrite standards were prepared in the same medium used to culture the macrophages. Absorbance was measured at 550 nm immediately after addition of the reagents to the supernatant samples.

#### Treatment of infected macrophages

Macrophages were incubated with Mtb at the indicated MOI for 4-5 hours under 10% oxygen. The supernatant was removed and the macrophages were washed 3 times with room temperature PBS to remove extracellular bacteria. Fresh medium was added to the cells, followed by the addition of nitrite, nitrate and/or an antimycobacterial drug such as INH. The cultures were then returned to the low oxygen incubator and incubated for a subsequent 3 days with the desired compound. Drugs were freshly dissolved before each experiment in DMSO, except for streptomycin, which was dissolved in water.

#### CFU determination from infected macrophages

Following incubation with Mtb for 3 days, the macrophages were inspected by microscopy to ensure continued confluence and adherence to the well. The supernatant was collected and the cells were gently washed 3 times with room temperature PBS to wash away remaining compounds. Following the wash steps, the cells were again visualized to confirm that the monolayer was intact and then lysed by incubation in 0.5% Triton X-100 for 10 minutes at 37°C. The lysate was serially diluted in 0.1% Triton X-100 and plated on Middlebrook 7H11 agar with 10% oleic acidalbumin-dextrose-catalase (OADC) enrichment (Difco) supplemented with 0.5% glycerol. Colonies were enumerated following 3 weeks of culture at 37 °C, in 21% O<sub>2</sub>, 5% CO<sub>2</sub>.

#### Treatment of Mtb in axenic culture

Mtb was added at the indicated OD to 96 well plates in 7H9 broth with or without 5 mM nitrate or the indicated concentration of nitrite, supplemented with 10 ADNaCl (0.2% glycerol, 0.5% BSA, 0.2% dextrose, 0.085% NaCl) and 0.05% Tween 80. The indicated antibiotic was added immediately and the plate was incubated in 1% oxygen for 3 days or at 21% oxygen for 5 days, as indicated. The Mtb was resuspended, serially diluted in 0.1% Triton-X 100 and plated for CFU as above. Alternatively, for OD measurements, the Mtb was resuspended in the culture medium and the absorbance was read at 580 nm on the fifth day of incubation with the indicated compound. In the case of exposure to hydrogen peroxide, the plate was incubated in 1% oxygen for 8 hours with or without 1 mM nitrate or the indicated concentration of nitrite. At this point, the indicated concentration of hydrogen peroxide was added and the cells were incubated overnight at 1% oxygen before plating for CFU.

#### KatG western blot and measurement of KatG dependent oxidation of INH

A minimum of  $4x10^9$  bacteria per condition were suspended in 7H9, 10% ADNaCl at an OD of 0.2 in the presence or absence of 2.5 mM nitrite. Cultures were incubated for 1 day in 1% oxygen. The cultures were collected by centrifugation and resuspended in 200  $\mu$ l of lysis buffer containing 50 mM NaPO<sub>4</sub>, 1 mM PMSF and 4x complete protease inhibitor cocktail (Roche). The cells were lysed in a bead-beating homogenizer with silica beads 3 times with intermittent incubation on ice. The lysate was collected and filter sterilized and concentrated at 4°C to ~100 µl using Amicon Ultra-0.5 mL centrifugal filters from Millipore with a molecular weight cut-off of 3 kDa. Protein concentration was measured using the Bio-Rad assay and equal amounts of protein from each sample were loaded into 7.5% reducing gels. After electrophoresis, proteins were transferred onto a polyvinylidene difluoride membrane, blocked with Odyssey blocking buffer from LI-COR Biosciences and stained with anti-KatG antibody (obtained through BEI Resources, NIAID, NIH: Monoclonal Anti-Mycobacterium tuberculosis KatG, gene Rv1908c, clone IT-57 (CDA4) (culture supernatant), NR-13793) and anti-DlaT antibody (1:1000) and then visualized with fluorophore-coupled secondary antibodies (45). To test the degree of KatG dependent oxidation of INH, 50 µg of total cell lysate was loaded into 7.5% nonreducing gels. After electrophoresis, the gel was incubated at room temperature for 15 - 30 minutes in 50 µLs of NaPO<sub>4</sub> buffer containing 62 mg INH, and 2 mM nitroblue tetrazolium (NBT) at pH 7 to which hydrogen peroxide was added to a final concentration of 800  $\mu$ M (25).

# Quantitative PCR of RNA from Mtb growing in macrophages or broth culture

Five million human macrophages were cultured per T25 flask in 10% oxygen and infected with wild type, narG-deficient or the complemented strains of Mtb at an MOI of 40 for 10 hours. Supernatant was removed and the monolayer was washed 2 times with room temperature PBS. Trizol (2 mL) was added to each flask and the cells were detached using a rubber policeman. For broth cultures, at least 2.5x10<sup>9</sup> bacteria per condition were incubated in 1% oxygen for 3 days with or without 2.5 mM nitrite and 0.1  $\mu$ g mL<sup>-1</sup> INH. Then an equal volume of 5M GTC buffer containing 5M guanidine thiocyanate, 25 mM sodium citrate, 20 mM N-lauryl-sarcosine, 0.7% v/v  $\beta$ - mercaptoethanol, was added. The bacteria were pelleted by centrifugation and 1 mL of Trizol was added per sample. The suspension was beaten with silica beads, 3 times. RNA was extracted using an RNeasy kit (Qiagen) in accordance with the manufacturer's instructions with the exception that off column DNAse digestion was performed for 2 hrs at 37 °C. To generate cDNA, total RNA (900 ng) was reverse transcribed using the GeneAmp RNA PCR Kit (Applied Biosystems). Quantitative RT-PCR was performed using gene specific primers (Life Technologies) and the SuperScript III Platinum Two Step qRT-PCR Kit (Life Technologies) with a 7900HT Fast Real Time PCR System (Applied Biosystems). Each experiment contained experimental triplicates. All reported values were within the linear range of the primers and the experimental results were normalized to 16S rRNA values.

### Statistical analysis

Statistical analysis was performed as indicated in the figure legends using the standard statistical software Prism version 5.0f for Macintosh, GraphPad Software, San Diego California USA

## References

- 1. Nicholson S et al. (1996) Inducible nitric oxide synthase in pulmonary alveolar macrophages from patients with tuberculosis. *J Exp Med* 183:2293–2302.
- 2. Lundberg JO, Weitzberg E (2010) NO-synthase independent NO generation in mammals. *Biochem Biophys Res Commun* 396:39–45.
- 3. Choi H-S, Rai PR, Chu HW, Cool C, Chan ED (2002) Analysis of nitric oxide synthase and nitrotyrosine expression in human pulmonary tuberculosis. *Am J Respir Crit Care Med* 166:178–186.
- 4. Sohaskey CD, Wayne LG (2003) Role of *narK2X* and *narGHJI* in hypoxic upregulation of nitrate reduction by *Mycobacterium tuberculosis*. *Journal of Bacteriology* 185:7247–7256.
- 5. Wayne LG, Sohaskey CD (2001) Nonreplicating persistence of *Mycobacterium tuberculosis*. *Annu Rev Microbiol* 55:139–163.
- 6. Giffin MM, Raab RW, Morganstern M, Sohaskey CD (2012) Mutational analysis of the respiratory nitrate transporter NarK2 of *Mycobacterium tuberculosis*. *PLoS ONE* 7:e45459.
- 7. Voskuil MI et al. (2003) Inhibition of respiration by nitric oxide induces a *Mycobacterium tuberculosis* dormancy program. *Journal of Experimental Medicine* 198:705–713.
- 8. Sherman DR et al. (2001) Regulation of the *Mycobacterium tuberculosis* hypoxic response gene encoding alpha -crystallin. *Proc Natl Acad Sci USA* 98:7534–7539.
- 9. Rachman H et al. (2006) Unique transcriptome signature of *Mycobacterium tuberculosis* in pulmonary tuberculosis. *Infect Immun* 74:1233–1242.
- 10. Fenhalls G et al. (2002) *In situ* detection of *Mycobacterium tuberculosis* transcripts in human lung granulomas reveals

differential gene expression in necrotic lesions. *Infect Immun* 70:6330–6338.

- 11. Shi L et al. (2005) Changes in energy metabolism of *Mycobacterium tuberculosis* in mouse lung and under in vitro conditions affecting aerobic respiration. *Proc Natl Acad Sci USA* 102:15629–15634.
- 12. Vogt G, Nathan C (2011) *In vitro* differentiation of human macrophages with enhanced antimycobacterial activity. *J Clin Invest* 121:3889–3901.
- 13. Wayne LG, Hayes LG (1998) Nitrate reduction as a marker for hypoxic shiftdown of *Mycobacterium tuberculosis*. *Tuber Lung Dis* 79:127–132.
- 14. Winder FG, Collins PB (1970) Inhibition by isoniazid of synthesis of mycolic acids in *Mycobacterium tuberculosis*. J Gen Microbiol 63:41–48.
- 15. Rozwarski DA (1998) Modification of the NADH of the isoniazid target (InhA) from *Mycobacterium tuberculosis*. *Science* 279:98–102.
- 16. Angeby KAK, Klintz L, Hoffner SE (2002) Rapid and inexpensive drug susceptibility testing of *Mycobacterium tuberculosis* with a nitrate reductase assay. *J Clin Microbiol* 40:553–555.
- 17. Wilson M et al. (1999) Exploring drug-induced alterations in gene expression in Mycobacterium tuberculosis by microarray hybridization. *Proc Natl Acad Sci USA* 96:12833–12838.
- 18. Baulard AR et al. (2000) Activation of the pro-drug ethionamide is regulated in mycobacteria. *J Biol Chem* 275:28326–28331.
- DeBarber AE, Mdluli K, Bosman M, Bekker LG, Barry CE (2000) Ethionamide activation and sensitivity in multidrug-resistant *Mycobacterium tuberculosis*. *Proc Natl Acad Sci USA* 97:9677– 9682.

- 20. Dhar N, McKinney JD (2010) *Mycobacterium tuberculosis* persistence mutants identified by screening in isoniazid-treated mice. *Proceedings of the National Academy of Sciences* 107:12275– 12280.
- Klebanoff SJ (1993) Reactive nitrogen intermediates and antimicrobial activity: role of nitrite. *Free Radic Biol Med* 14:351– 360.
- 22. van der Vliet A, Eiserich JP, Halliwell B, Cross CE (1997) Formation of reactive nitrogen species during peroxidase-catalyzed oxidation of nitrite. A potential additional mechanism of nitric oxide-dependent toxicity. *J Biol Chem* 272:7617–7625.
- 23. Eiserich JP et al. (1998) Formation of nitric oxide-derived inflammatory oxidants by myeloperoxidase in neutrophils. *Nature* 391:393–397.
- 24. Battistuzzi G, Bellei M, Bortolotti CA, Sola M (2010) Redox properties of heme peroxidases. *Archives of Biochemistry and Biophysics* 500:21–36.
- 25. Saint-Joanis B et al. (1999) Use of site-directed mutagenesis to probe the structure, function and isoniazid activation of the catalase/peroxidase, KatG, from *Mycobacterium tuberculosis*. *Biochem J* 338 (Pt 3):753–760.
- 26. Musser JM et al. (1996) Characterization of the catalase-peroxidase gene (katG) and inhA locus in isoniazid-resistant and -susceptible strains of *Mycobacterium tuberculosis* by automated DNA sequencing: restricted array of mutations associated with drug resistance. *J Infect Dis* 173:196–202.
- 27. Wengenack NL et al. (2004) Purification and characterization of *Mycobacterium tuberculosis* KatG, KatG(S315T), and *Mycobacterium bovis* KatG(R463L). *Protein Expression and Purification* 36:232–243.
- 28. Ghiladi RA, Medzihradszky KF, Rusnak FM, Ortiz de Montellano

PR (2005) Correlation between isoniazid resistance and superoxide reactivity in *Mycobacterium tuberculosis* KatG. *J Am Chem Soc* 127:13428–13442.

- 29. Cade CE, Dlouhy AC, Medzihradszky KF, Salas-Castillo SP, Ghiladi RA (2010) Isoniazid-resistance conferring mutations in *Mycobacterium tuberculosis* KatG: catalase, peroxidase, and INH-NADH adduct formation activities. *Protein Science*:458–474.
- 30. Vilcheze C et al. (2011) Coresistance to isoniazid and ethionamide maps to mycothiol biosynthetic genes in *Mycobacterium bovis*. *Antimicrob Agents Chemother* 55:4422–4423.
- 31. Vilchèze C et al. (2005) Altered NADH/NAD+ ratio mediates coresistance to isoniazid and ethionamide in mycobacteria. *Antimicrob Agents Chemother* 49:708–720.
- 32. Hazbon MH et al. (2006) Population genetics study of isoniazid resistance mutations and evolution of multidrug-resistant *Mycobacterium tuberculosis*. *Antimicrob Agents Chemother* 50:2640–2649.
- 33. Lundberg JO, Weitzberg E, Gladwin MT (2008) The nitrate–nitrite– nitric oxide pathway in physiology and therapeutics. *Nat Rev Drug Discov* 7:156–167.
- 34. Pacelli R et al. (1995) Nitric oxide potentiates hydrogen peroxideinduced killing of *Escherichia coli*. J Exp Med 182:1469–1479.
- 35. Woodmansee AN, Imlay JA (2003) A mechanism by which nitric oxide accelerates the rate of oxidative DNA damage in *Escherichia coli*. *Mol Microbiol* 49:11–22.
- 36. Nathan C, Shiloh MU (2000) Reactive oxygen and nitrogen intermediates in the relationship between mammalian hosts and microbial pathogens. *Proc Natl Acad Sci USA* 97:8841–8848.
- 37. Seth D, Hausladen A, Wang YJ, Stamler JS (2012) Endogenous protein S-nitrosylation in *E. coli*: regulation by OxyR. *Science*

336:470-473.

- Ralt D, Wishnok JS, Fitts R, Tannenbaum SR (1988) Bacterial catalysis of nitrosation: involvement of the nar operon of *Escherichia coli*. Journal of Bacteriology 170:359–364.
- Corker H (2003) Nitric oxide formation by *Escherichia coli*: dependence on nitrite reductase, the NO-sensing regulator Fnr, and flavohemoglobin Hmp. *Journal of Biological Chemistry* 278:31584– 31592.
- 40. Rhee KY, Erdjument-Bromage H, Tempst P, Nathan CF (2005) Snitroso proteome of *Mycobacterium tuberculosis*: enzymes of intermediary metabolism and antioxidant defense. *Proc Natl Acad Sci USA* 102:467–472.
- 41. van den Boogaard J et al. (2009) Community vs. facility-based directly observed treatment for tuberculosis in Tanzania's Kilimanjaro Region. *Int J Tuberc Lung Dis* 13:1524–1529.
- 42. Zumla A, Raviglione M, Hafner R, Fordham von Reyn C (2013) Tuberculosis. *N Engl J Med* 368:745–755.
- 43. Vilcheze C et al. (2011) Novel inhibitors of InhA efficiently kill *Mycobacterium tuberculosis* under aerobic and anaerobic conditions. *Antimicrob Agents Chemother* 55:3889–3898.
- 44. Stermann M, Bohrssen A, Diephaus C, Maass S, Bange F-C (2003) Polymorphic nucleotide within the promoter of nitrate reductase (NarGHJI) is specific for *Mycobacterium tuberculosis*. J Clin Microbiol 41:3252–3259.
- 45. Venugopal A et al. (2011) Virulence of *Mycobacterium tuberculosis* depends on lipoamide dehydrogenase, a member of three multienzyme complexes. *Cell Host Microbe* 9:21–31.
- 46. Wanger A, Mills K (1996) Testing of *Mycobacterium tuberculosis* susceptibility to ethambutol, isoniazid, rifampin, and streptomycin by using Etest. *J Clin Microbiol* 34:1672–1676.

## CHAPTER 4 CONCLUSIONS

Within Mtb-infected humans, nitrite levels rise to such substantial levels as to be measured within the plasma (1, 2). This nitrite is likely a product of the host's immune response as well as Mtb's hypoxic respiratory pathway. However, these enzymatic processes are inversely regulated by oxygen tension (3, 4). Therefore, in the host whether Mtb faces normoxia or hypoxia it will encounter nitrite and the RNS that generate or arise from it. That Mtb resides within hypoxic tissue in the human host is inferred from experiments with infected model hosts, which develop hypoxic lesions containing Mtb (5-7). In addition, lung tissue resected from patients with TB contains RNA transcripts that encode proteins essential to nitrate respiration (8, 9). Therefore, Mtb likely respires nitrate within the human host.

The majority of infected individuals will never eradicate Mtb, though most successfully sequester the pathogen. Subpopulations of mycobacteria residing within the human host are thought to enter into a dormant, non-replicating state and to survive for the lifetime of the host (10). Persisting mycobacteria will serve as a nidus for disease reactivation and infection of new hosts (10, 11). The extent to which nitrate respiration contributes to Mtb's persistence *in vivo* remains unknown. However, the host is comprised of heterogeneous microenvironments in which Mtb may transiently be exposed to concentrations of oxygen sufficient to support active replication (12). While renewed growth may be advantageous and contribute to the colonization of new hosts, ill-timed growth may disrupt the dynamic equilibrium that exists between the pathogen and host and compromise bacterial dissemination. Once a granulomatous lesion containing Mtb erodes into a bronchus, droplet nuclei containing Mtb are expelled from the diseased individual to enter and infect a naïve host (13). Therefore, in the setting of sustained normoxia, Mtb-derived nitrite may limit untimely growth and provide a store of metabolic energy to provision the colonization of naïve hosts.

Nitrate respiration regulated the expression profile of genes that are also regulated by exposure of Mtb to pH 5.5, hydrogen peroxide, nitric oxide, hypoxia, iron-depleted medium and residence within mouse macrophages. Therefore, nitrate respiration may support the survival of Mtb faced with hostile host stresses. In fact, this has already been demonstrated for Mtb exposed to low pH, RNS (14) and sudden anaerobiosis (15). We have reported that the combination of Mtb-derived nitrite and hydrogen peroxide is more lethal to Mtb than treatment with hydrogen peroxide alone (Cunningham-Bussel, manuscript in preparation, see chapter 2). However, while short incubations with nitrite synergized with hydrogen peroxide to kill Mtb, longer incubations seemed to promote the survival of hydrogen peroxide treated Mtb, although this remains to be confirmed (data not shown). Therefore in regards to hydrogen peroxide, nitrite may exert opposing effects depending both upon the concentration and length of exposure of Mtb to nitrite prior to treatment with hydrogen peroxide.

We also observed that in the absence of nitrate respiration, as compared to bacteria that continued to respire nitrate, Mtb became hyper-

87

susceptible to INH. Therefore, inhibitors of the nitrate reductase or nitrate/proton symporter narK2 may synergize with INH to significantly reduce the length of time required to treat TB. That INH remained bactericidal even under non-replicating conditions suggested that the hyper-susceptibility of narG-deficient Mtb to INH might have been due to the inhibition of a cellular protein(s) other than the recognized protein target of INH, InhA. Furthermore, regardless of any synergy with INH, nitrate respiration may prove to more broadly support the survival of Mtb *in vivo*. Therefore, compounds that prevent nitrate respiration may prove to be valuable additions to the arsenal of antitubercular antibiotics.

#### Future directions

Future experiments will need to first address whether nitrate respiration benefits the survival of Mtb in a model organism. Similarly, the observation that nitrate respiration allows Mtb to resist INH should also be tested *in vivo*. High-throughput screening of chemical libraries against axenic cultures of Mtb may identify novel compounds that inhibit the mycobacterial nitrate reductase and nitrate/proton symporter *narK2*. In addition, it may be possible to identify the protein target(s) of INH in non-replicating Mtb lacking a functional nitrate reductase by screening for spontaneous revertants of *narG*-deficient Mtb that lose their hypersusceptibility to INH. Alternatively, this may be accomplished by screening a library of transposon insertion mutants generated in *narG*-deficient Mtb for variants that have altered susceptibility to INH. The mutants that demonstrate either increased or reduced sensitivity may

contain transposon insertions in genes that regulate the susceptibility of non-replicating Mtb to INH.

*In vivo* experiments to determine the impact of nitrate respiration on Mtb within the host will require a model organism that develops hypoxic, TB-associated lesions, such as the guinea pig, rabbit or macaque (5). These organisms better mimic human tissue pathology as compared to the standard mouse model, which is not hypoxic (5-7). Recently, an alternative mouse model was reported to develop hypoxic lesions in response to infection with Mtb (16). This mouse strain may therefore be a more suitable model system in which to investigate the contribution of nitrate respiration to the persistence of Mtb *in vivo*.

Both the host and pathogen are potential sources of nitrite *in vivo*. Therefore, it will be important to determine the amount of nitrite individually produced by the host and pathogen and to selectively abolish these catalytic activities during the course of infection. The production of host-nitrite can be limited by the use of selective or non-selective NOS inhibitors (17, 18) or greatly repressed by genetic deletion of iNOS (19). Pathogen derived nitrite production is abolished by the genetic deletion of *narG* (4). In addition to the selective abrogation of nitrite generated by the host or pathogen, nitrite can be supplemented by adding nitrate or nitrite supplementation can greatly increase the concentration of plasma and tissue nitrite as well as the concentration of tissue S-nitrosylation (20). Therefore, an additional experiment might involve infection of iNOS deficient host organisms with *narG*-deficient Mtb followed by the supplemental administration of exogenous nitrate or nitrite, which may

contribute to determining the impact of nitrite on Mtb in vivo.

Mycobacterial nitrate respiration caused a non-heritable resistance to INH, which remains mechanistically unexplained. Nitrite and the RNS that arise from it are known to post-translationally modify cellular proteins and interfere with their function (22-25). Drug resistance is often attributed to the acquisition of heritable genetic mutations that protect bacterial populations from cidal compounds. However, non-heritable survival has also been described to account for the persistence of genetically susceptible bacteria treated with bactericidal antibiotics (26, 27). Mtb persists in vivo, despite treatment with antimycobacterial antibiotics such as INH (28). Therefore, despite an underlying genetic sensitivity to INH, the survival of subpopulations of Mtb treated with INH may be due to the resistance afforded by mycobacterial nitrate respiration. Nitrite is highly diffusible and therefore could non-heritably manipulate array of cellular targets across a population of bacteria. Therefore, INH resistance is but one functional consequence of mycobacterial nitrate respiration; others likely remain to be characterized.

Nitrite has previously been demonstrated to regulate mammalian gene expression (23, 29). However, we did not observe any impact of mycobacterial nitrate respiration on the transcriptome of Mtb infected human macrophages. However, unlike the transcriptional profiles of groups of genetically identical mice or bacteria, the transcriptional profile of Mtb infected macrophages was analyzed using RNA collected from three genetically dissimilar individuals. We propose that the lack of an observable impact of nitrate respiration on the transcriptome of infected primary human macrophages may be due the reportedly variable gene expression that exists between unrelated humans (30).

In summary, we have demonstrated that mycobacterial nitrite reduction impacts a variety of previously described mycobacterial cellular processes. However, we anticipate that Mtb-derived nitrite broadly impacts not only the cellular processes of Mtb itself, but also the host. Nitrate respiration may therefore play a central role in the interaction between the host and mycobacterial pathogen, the full extent of which remains to be characterized.

## References

- 1. Lamsal M et al. (2007) Evaluation of lipid peroxidation product, nitrite and antioxidant levels in newly diagnosed and two months follow-up patients with pulmonary tuberculosis. *Southeast Asian J Trop Med Public Health* 38:695–703.
- Dlugovitzky D et al. (2000) Influence of disease severity on nitrite and cytokine production by peripheral blood mononuclear cells (PBMC) from patients with pulmonary tuberculosis (TB). *Clin Exp Immunol* 122:343–349.
- 3. McCormick CC, Li WP, Calero M (2000) Oxygen tension limits nitric oxide synthesis by activated macrophages. *Biochem J* 350 Pt 3:709–716.
- 4. Sohaskey CD, Wayne LG (2003) Role of *narK2X* and *narGHJI* in hypoxic upregulation of nitrate reduction by *Mycobacterium tuberculosis*. *Journal of Bacteriology* 185:7247–7256.
- 5. Via LE et al. (2008) Tuberculous granulomas are hypoxic in guinea pigs, rabbits, and nonhuman primates. *Infect Immun* 76:2333–2340.
- 6. Aly S et al. (2006) Oxygen status of lung granulomas in *Mycobacterium tuberculosis*-infected mice. *J Pathol* 210:298–305.
- 7. Tsai MC et al. (2006) Characterization of the tuberculous granuloma in murine and human lungs: cellular composition and relative tissue oxygen tension. *Cell Microbiol* 8:218–232.
- 8. Fenhalls G et al. (2002) *In situ* detection of *Mycobacterium tuberculosis* transcripts in human lung granulomas reveals differential gene expression in necrotic lesions. *Infect Immun* 70:6330–6338.
- 9. Rachman H et al. (2006) Unique transcriptome signature of *Mycobacterium tuberculosis* in pulmonary tuberculosis. *Infect Immun* 74:1233–1242.

- 10. Wayne LG, Sohaskey CD (2001) Nonreplicating persistence of *Mycobacterium tuberculosis. Annu Rev Microbiol* 55:139–163.
- 11. Flynn JL, Chan J (2001) Tuberculosis: latency and reactivation. *Infect Immun* 69:4195–4201.
- 12. Boshoff HIM (2004) The transcriptional responses of *Mycobacterium tuberculosis* to inhibitors of metabolism: novel insights into drug mechanisms of action. *Journal of Biological Chemistry* 279:40174–40184.
- 13. Nathan C (2009) Taming tuberculosis: a challenge for science and society. *Cell Host Microbe* 5:220–224.
- 14. Tan MP et al. (2010) Nitrate respiration protects hypoxic *Mycobacterium tuberculosis* against acid- and reactive nitrogen species stresses. *PLoS ONE* 5:e13356.
- 15. Sohaskey CD (2008) Nitrate enhances the survival of *Mycobacterium tuberculosis* during inhibition of respiration. *Journal of Bacteriology* 190:2981–2986.
- Harper J et al. (2012) Mouse model of necrotic tuberculosis granulomas develops hypoxic lesions. *Journal of Infectious Diseases* 205:595–602.
- Alderton WK, Cooper CE, Knowles RG (2001) Nitric oxide synthases: structure, function and inhibition. *Biochem J* 357:593– 615.
- 18. Wolff DJ, Lubeskie A (1995) Aminoguanidine is an isoformselective, mechanism-based inactivator of nitric oxide synthase. *Archives of Biochemistry and Biophysics* 316:290–301.
- 19. MacMicking JD et al. (1997) Identification of nitric oxide synthase as a protective locus against tuberculosis. *Proc Natl Acad Sci USA* 94:5243–5248.
- 20. Carlström M et al. (2010) Dietary inorganic nitrate reverses features

of metabolic syndrome in endothelial nitric oxide synthase-deficient mice. *Proceedings of the National Academy of Sciences* 107:17716–17720.

- 21. Carlström M et al. (2011) Dietary nitrate attenuates oxidative stress, prevents cardiac and renal injuries, and reduces blood pressure in salt-induced hypertension. *Cardiovascular Research* 89:574–585.
- 22. Rhee KY, Erdjument-Bromage H, Tempst P, Nathan CF (2005) Snitroso proteome of *Mycobacterium tuberculosis*: enzymes of intermediary metabolism and antioxidant defense. *Proc Natl Acad Sci USA* 102:467–472.
- Bryan NS et al. (2005) Nitrite is a signaling molecule and regulator of gene expression in mammalian tissues. *Nat Chem Biol* 1:290–297.
- 24. Lundberg JO, Weitzberg E (2010) NO-synthase independent NO generation in mammals. *Biochem Biophys Res Commun* 396:39–45.
- 25. Lundberg JO, Weitzberg E, Gladwin MT (2008) The nitrate–nitrite– nitric oxide pathway in physiology and therapeutics. *Nat Rev Drug Discov* 7:156–167.
- 26. Wakamoto Y et al. (2013) Dynamic persistence of antibioticstressed mycobacteria. *Science* 339:91–95.
- 27. Balaban NQ, Gerdes K, Lewis K, McKinney JD (2013) A problem of persistence: still more questions than answers? *Nat Rev Microbiol* 11:587–591.
- 28. Zumla A, Raviglione M, Hafner R, Fordham von Reyn C (2013) Tuberculosis. *N Engl J Med* 368:745–755.
- 29. Gusarov I et al. (2013) Bacterial nitric oxide extends the lifespan of *C. elegans. Cell* 152:818–830.
- 30. Cheung VG et al. (2003) Natural variation in human gene expression assessed in lymphoblastoid cells. *Nat Genet* 33:422–425.

## PUBLICATIONS

- 1. Cunningham-Bussel A, Zhang T, Nathan C *Mycobacterium tuberculosis* generates nitrite in human macrophages at physiologic oxygen tensions: impact on mycobacterial ATP consumption and gene expression. (*manuscript submitted*).
- 2. Cunningham-Bussel A, Bange F-C, Nathan C Nitrite impacts the survival of *Mycobacterium tuberculosis* in response to isoniazid and hydrogen peroxide. *Microbiology Open, in press.*
- 3. Nathan C, Cunningham-Bussel A (2013) Beyond oxidative stress: an immunologist's guide to reactive oxygen species. *Nat Rev Immunol* 13:349–361.
- 4. Gold B et al. (2012) Nonsteroidal anti-inflammatory drug sensitizes Mycobacterium tuberculosis to endogenous and exogenous antimicrobials. *Proceedings of the National Academy of Sciences* 109:16004–16011.
- 5. Penack O et al. (2009) NOD2 regulates hematopoietic cell function during graft-versus-host disease. *J Exp Med* 206:2101–2110.
- 6. Root JC et al. (2009) Frontolimbic function and cortisol reactivity in response to emotional stimuli. *Neuroreport* 20:429–434.
- 7. Cunningham-Bussel AC et al. (2009) Diurnal cortisol amplitude and fronto-limbic activity in response to stressful stimuli. *Psychoneuroendocrinology* 34:694–704.
- 8. Butler T et al. (2007) Sex specificity of ventral anterior cingulate cortex suppression during a cognitive task. *Hum Brain Mapp* 28:1206–1212.
- 9. Butler T et al. (2007) Human fear-related motor neurocircuitry. *Neuroscience* 150:1–7.

## **APPENDIX 1**


Significance of

the overlap,

p value

4.17E-08

2.05E-01

< 2.2e-16

6.53E-04

< 2.2e-16

1.11E-08

4.65E-06

5.72E-03

3.27E-02

< 2.2e-16

< 2.2e-16

5.63E-03

3.31E-03

2.39E-11

9.65E-01 9.67E-01

1.94E-01

7.69E-01

1.94E-01

t of total gene

hat overlaps

11.67 1.67

22.50

14.17

23 33

18.33

8.33 13.33

15.00

54.17

26.67

9.17

2.50

26.67

0.83

0.83 1.67

5.00

10.00

10.83

Supplementary Figure 1. iNOS was not detected within Mtb-infected primary human macrophages. Macrophages were either infected with Mtb (MOI: 5, indicated by a "+" sign) or remained uninfected for 3 days at 10% oxygen. Results are shown for 6 individual donors. Lysates were separated by reducing SDS polyacrylamide gel electrophoresis and probed with anti-iNOS antibody (Santa Cruz Biotechnology sc-8310) and anti- $\beta$ actin (abcam 8226). The positive control (+ con) was generated by lipofectamine-assisted transfection of HEK293 cells with a pcDNA3.1 plasmid construct containing human iNOS. To generate the negative control, HEK293 cells were transfected with an empty plasmid.



Supplementary Figure 2. Consumption of exogenous ATP added to cell lysates from nitrite treated or untreated bacterial cultures. Wt Mtb was incubated in the presence or absence of 2.5 mM nitrite for 3 days in 1% oxygen; cell lysates were incubated for 20 minutes at room temperature to allow for the complete metabolism of endogenous ATP. Reagent ATP (10  $\mu$ M) was then added to the cell lysates and ATP content measured at intervals. Results are from 1 experiment representative of 3.



Supplementary Figure S3. Average expression of *narK2* by untreated and nitrate treated wild type Mtb. Means  $\pm$  SD, n=3 independent experiments. The p value was determined by an unpaired t test.

## Supplementary Table 1. Genes significantly induced by nitrate. $(p_{\rm corr}$ <0.05), (n=3)

	PATIO						
ConcilD	KALLU Gene (Nitrate/Untroated) n corr <0.0						
Dv2780	Gene	(Nitrate/Untreated)	2 58E-03				
RV2700	aiu cmtP	9.30	2.30E-03				
RV19940	cinick	4.22	2.392-02				
Rv3743c	ctn /	5.82	2.72E-04				
Rv3019c	cipu esvP	6.10	4.23E-03				
RV30190	esxr	6.86	1.28E-03				
RV30200	fadD0	5.04	3 425 02				
Rv2390	fadE0	1 23	3.42E-02				
By1000c	furA	7.23	2 41 = 02				
Dv2384	mht A	5.65	2.41E-03				
Dv2383c	mbtR	30.00	5.50E-04				
RV2380c	mbtE	13 40	1.02E-02				
RV2300C	mbt	10.40	2 405 05				
RV2300C	mbt l	12.52	5.40E-03				
Rv2305	moeB1	4.00	1.07E-03				
RV32000		0.42	4.90E-02				
RV0747	PE_PGRS10	4.50	1.392-03				
RV0034C	PE_PGRS14	11.01	1.94E-02				
RV0978C	PE_PGRS17	6.20	1.19E-05				
RV2015C		0.30	1.07E-02				
RV3746C	PE34	7.84	3.40E-05				
RV 1402	priA Du0140	0.01	1.22E-03				
RV0142	RV0142	0.10	1.80E-03				
RV0396C	RV0396C	J.11 A 96	2.01E-02				
RV0403C	RV0405C	4.00	5.200-02				
RV0575C	RV0575C	10.5	5.07 E-00				
RV0041	RV0041	5.91	4.912-02				
RV1049	RV1049	0.40	1.09E-03				
RV1050	RV1050	3.90	4.04E-02				
RV1007	RV1037	4.40	2.100-02				
Rv1403c	RV1403c	15 01	2.33E-03				
Rv1405c	Rv1405c	14 53	4 71E-05				
Rv1461	Rv1461	34 70	1.71E-03				
Rv1578c	Rv1578c	3 83	4 89E-02				
Rv1670	Rv1670	4 18	3.09E-02				
Rv1671	Rv1671	4.10	2 39E-02				
Rv1767	Rv1767	4.05	3.51E-02				
Rv1907c	Rv1907c	4 14	2 08E-02				
Rv1986	Rv1986	3 93	1.32E-02				
Rv2324	Rv2324	4 03	3.58E-02				
Rv2616	Rv2616	100.96	1.00E-11				
Rv2617c	Rv2617c	137 41	4 23E-10				
Rv2618	Rv2618	26.9	4.08E-04				
Rv2619c	Rv2619c	20.75	3 11E-05				
Rv2621c	Rv2621c	21.63	1 78E-06				
Rv2622	Rv2622	10.26	1 21E-06				
Rv2650c	Rv2650c	5 21	6.03E-03				
Rv2651c	Rv2651c	4 00	2.39E-02				
Rv2729c	Rv2729c	4 28	1 14E-02				
Rv2781c	Rv2781c	6.21	7.53F-04				
Rv3289c	Rv3289c	8 99	3 29F-04				
Rv3402c	Rv3402c	6.94	2.36F-04				
Rv3403c	Rv3403c	4.70	2.51F-02				
Rv3832c	Rv3832c	6.02	2.94F-02				
Rv3839	Rv3839	28.77	4.23F-10				
Rv3840	Rv3840	4.57	1.05E-02				
Rv3288c	usfY	6.32	7.30E-03				

## Supplementary Table 2. Genes significantly repressed by nitrate. $(p_{\mbox{\tiny corr}}$

<0.05), (n=3) Genes labeled with an asterix are members of the DOS regulon as defined by (20).

		RATIO					
	Gene ID	Gene	(Nitrate/Untreated)	p corr <0.05			
*	Rv0569	Rv0569	0.06	4.01E-10			
*	Rv2007c	fdxA	0.05	4.67E-10			
*	Rv2627c	Rv2627c	0.06	7.49E-10			
*	Rv2626c	hrp1	0.06	1.08E-09			
*	Rv3841	bfrB	0.05	4.29E-09			
*	Rv0570	nrdZ	0.08	2.76E-08			
*	Rv1733c	Rv1733c	0.09	0.00000036			
*	Rv2031c	hspX	0.07	0.000000192			
*	Rv2625c	Rv2625c	0.09	0.000000408			
*	Rv1739c	Rv1739c	0.10	0.000000643			
*	Rv2032	acg	0.10	0.00000105			
*	Rv2628	Rv2628	0.11	0.00000134			
*	Rv1738	Rv1738	0.10	0.0000014			
*	Rv0572c	Rv0572c	0.12	0.00000188			
*	Rv2623	TB31.7	0.09	0.00000203			
*	Rv3134c	Rv3134c	0.10	0.00000203			
*	Rv1997	ctpF	0.11	0.00000272			
<i>*</i>	Rv2030c	Rv2030c	0.09	0.00000429			
_ _	RV3130C	tgs1	0.09	0.0000295			
Ĵ	Rv1998c	Rv1998c	0.16	0.0000559			
<u>^</u>	RV3127	Rv3127	0.08	0.0000559			
<u>^</u>	Rv2624C	Rv2624c	0.15	0.000211			
- +	RV2029C	рткв	0.09	0.00027			
*	RV3131	RV3131	0.04	0.0018			
*	RV3120C	RV3126C	0.18	0.00482			
*	RV2000	OISB /	0.21	0.00698			
*	RV1990	RV1996	0.21	0.0076			
*	RV17300	nark2	0.10	0.00991			
*	RV1737C	Dv2028c	0.09	0.0358			
	11020200	1120200	0.15	0.0400			
	Rv1195	PE13	0.10	0.00000123			
	Rv0700	rpsJ	0.16	0.000454			
	Rv2190c	Rv2190c	0.17	0.000626			
	Rv0315	Rv0315	0.18	0.000646			
	Rv1196	PPE18	0.17	0.00278			
	Rv0227c	Rv0227c	0.20	0.00306			
	Rv0892	Rv0892	0.19	0.00508			
	Rv0714	rpIN	0.22	0.0052			
	Rv0237	lpql	0.22	0.0063			
	Rv0718	rpsH	0.23	0.00635			
	Rv2952	Rv2952	0.21	0.00789			
	Rv2033c	Rv2033c	0.22	0.0102			
	Rv2563	Rv2563	0.23	0.0108			
	RV0/1/	rpsN1	0.25	0.0109			
	RV0009	ppiA	0.23	0.0131			
	RV1884C	rprC	0.23	0.0148			
	Dv24500	yillQ rof⊑	0.20	0.015			
	Dv18060	ipie fbpB	0.19	0.0209			
	DVU033	iope nete	0.10	0.0239			
	Rv2816c	PV28160	0.27	0.025			
	Rv0710		0.20	0.0209			
	Rv1515c	Rv1515c	0.27	0.0270			
	Rv20/5c	InnY	0.30	0.0342			
	Rv2959c	1222 Rv2959c	0.23	0.0335			
	Rv2416c	eis	0.31	0.0431			

		RATIO	
Gene ID	Gene	(Nitrate/Untreated)	p corr <0.05
Rv2917	Rv2917	0.31	0.0431
Rv0934	pstS1	0.27	0.0475
Rv3628	рра	0.30	0.0475
Rv0935	pstC1	0.30	0.0476
Rv0165c	mce1R	0.28	0.0492
Rv0867c	rpfA	0.28	0.0494
Rv2431c	PE25	0.27	0.0494

Supplementary Table 3. Genes that are significantly regulated by nitrate and no other tested stimuli.  $(p_{corr} < 0.05)$  The expression of genes regulated only by nitrate and no other stimuli reported in Table 1.

-	-	Entrez	- Untreated	Nitrate	RATIO (Nitrate/Un			
Gene	Gene ID	Gene ID	(RPKM)	(RPKM)	treated)	p <sub>corr</sub> <0.05	Function	Functional category
Rv0978c	PE_PGRS17	885077	0.38	4.25	11.21	0.000012	Function unknown	PE/PPE Intermediary
Rv2781c	Rv2781c	888488	4.24	26.37	6.21	0.0008	Function unknown	metabolism and respiration
Rv1402	priA	886716	0.31	1.88	6.01	0.0012	Functions as a helicase within the primosome	Information pathways
Rv3743c	ctpJ	885106	0.50	2.89	5.82	0.0042	Cation-transporting ATPase	Cell wall and cell processes
Rv0237	lpql	886693	3.11	0.69	0.22	0.0063	Probable conserved lipoprotein	Cell wall and cell processes
Rv0747	PE_PGRS10	888662	1.61	7.33	4.56	0.0074	Function unknown	PE/PPE, lipid metabolism
Rv2615c	PE_PGRS45	888215	1.35	8.50	6.30	0.0107	Function unknown	PE/PPE
Rv2563	Rv2563	887516	3.78	0.85	0.23	0.0108	Active transport of glutamine	Cell wall and cell processes
Rv0752c	fadE9	888684	2.85	12.08	4.23	0.0334	Function unknown	Lipid metabolism
Rv1515c	Rv1515c	886459	1.94	0.57	0.30	0.0342	Function unknown	Conserved hypothetical
Rv2917	Rv2917	887758	1.34	0.42	0.31	0.0431	Function unknown, but involved in lipid degradation	Conserved hypothetical, cell wall and cell processes
Rv0841	Rv0841	3205068	9.10	35.58	3.91	0.0491	Function unknown	Cell wall and cell processes
Rv0165c	mce1R	886818	4.50	1.27	0.28	0.0492	Probable transcriptional regulatory protein Mce1R (probably GntR-family)	Regulatory proteins

## **APPENDIX 2**



1. Effect of nitrate respiration Supplementary Figure on mycobacterial resistance to streptomycin, rifampicin and ethambutol within primary human macrophages. (A-C) Survival of wild type, narG-deficient (NarG) and complemented strains (NarGc) within macrophages (MOI: 0.1, corresponding to 10,000 bacteria) treated with the indicated concentrations of (A) rifampicin (B) streptomycin or (C) ethambutol over 3 days in 10% oxygen. The results shown in A-C are single experiments that are representative of at least two independent experiments conducted using cells isolated from at least two distinct human donors. (D-F) contain the pooled CFU data from at least two independent experiments conducted with cells from at least 2 individual human donors shown for one drug concentration as follows: (D) rifampicin (0.2  $\mu$ g mL<sup>-1</sup>) (E) streptomycin (0.6  $\mu$ g mL<sup>-1</sup>) or (F) ethambutol  $(0.1 \ \mu g \ mL^{-1})$ . Bars indicate means  $\pm$  SEM.



Supplementary Figure 2. Nitrite treatment hyper-sensitized Mtb to hydrogen peroxide treatment without an observed difference in the expression of *katG* or a measurably reduced degree of its peroxidase activity. (A) The survival of axenic cultures of Mtb (OD 0.1) following treatment with the indicated concentration of hydrogen peroxide at 1% oxygen incubated overnight with 1 mM nitrite. This is representative of 2 independent experiments. (B) The mRNA expression of katG by wild type, *narG* -deficient (NarG) and the complemented (NarGc) strains within infected human macrophages following incubation for three days at 10% oxygen. The given experiment is the pooled result of 2 independent experiments. (C) Axenic cultures of wild type, narGdeficient and the complemented strains of Mtb were incubated over one day at 1% oxygen and lysed to collect total cell protein. Individual proteins from the total cell lysate were separated by native gel electrophoresis and KatG and the housekeeping gene DLAT were visualized by fluorophore coupled antibody staining. This is representative of three independent experiments. Bars indicate means ± SEM.